

Rensselaer Polytechnic Institute
Department of Electrical, Computer, and Systems Engineering
ECSE 2500: Engineering Probability, Spring 2023
Homework #7 Solutions

1. (a) Each value of X in $[5, 7]$ has a $\frac{1}{3}$ probability of occurring since the distribution is uniform. If we let E be the number of eggs available, the distribution of E given X is a binomial with probability of success $\frac{1}{2}$. The number of eggs the carton Y will then be $\max(6, E)$. That is, once we have 6 eggs, we don't collect any more. So the joint PMF table looks like:

		X		
		5	6	7
Y	0	$\frac{1}{3} \cdot \left(\frac{1}{2}\right)^5$	$\frac{1}{3} \cdot \left(\frac{1}{2}\right)^6$	$\frac{1}{3} \cdot \left(\frac{1}{2}\right)^7$
	1	$\frac{1}{3} \cdot 5 \cdot \left(\frac{1}{2}\right)^5$	$\frac{1}{3} \cdot 6 \cdot \left(\frac{1}{2}\right)^6$	$\frac{1}{3} \cdot 7 \cdot \left(\frac{1}{2}\right)^7$
	2	$\frac{1}{3} \cdot \binom{5}{2} \cdot \left(\frac{1}{2}\right)^5$	$\frac{1}{3} \cdot \binom{6}{2} \cdot \left(\frac{1}{2}\right)^6$	$\frac{1}{3} \cdot \binom{7}{2} \cdot \left(\frac{1}{2}\right)^7$
	3	$\frac{1}{3} \cdot \binom{5}{3} \cdot \left(\frac{1}{2}\right)^5$	$\frac{1}{3} \cdot \binom{6}{3} \cdot \left(\frac{1}{2}\right)^6$	$\frac{1}{3} \cdot \binom{7}{3} \cdot \left(\frac{1}{2}\right)^7$
	4	$\frac{1}{3} \cdot \binom{5}{4} \cdot \left(\frac{1}{2}\right)^5$	$\frac{1}{3} \cdot \binom{6}{4} \cdot \left(\frac{1}{2}\right)^6$	$\frac{1}{3} \cdot \binom{7}{4} \cdot \left(\frac{1}{2}\right)^7$
	5	$\frac{1}{3} \cdot \left(\frac{1}{2}\right)^5$	$\frac{1}{3} \cdot \binom{6}{5} \cdot \left(\frac{1}{2}\right)^6$	$\frac{1}{3} \cdot \binom{7}{5} \cdot \left(\frac{1}{2}\right)^7$
	6	0	$\frac{1}{3} \cdot \left(\frac{1}{2}\right)^6$	$\frac{1}{3} \cdot \binom{7}{6} \cdot \left(\frac{1}{2}\right)^7 + \left(\frac{1}{2}\right)^7$


or getting everything into a common denominator,

		X		
		5	6	7
Y	0	$\frac{4}{384}$	$\frac{2}{384}$	$\frac{1}{384}$
	1	$\frac{20}{384}$	$\frac{12}{384}$	$\frac{7}{384}$
	2	$\frac{40}{384}$	$\frac{30}{384}$	$\frac{21}{384}$
	3	$\frac{40}{384}$	$\frac{40}{384}$	$\frac{35}{384}$
	4	$\frac{20}{384}$	$\frac{30}{384}$	$\frac{35}{384}$
	5	$\frac{4}{384}$	$\frac{12}{384}$	$\frac{21}{384}$
	6	0	$\frac{2}{384}$	$\frac{8}{384}$

Grading criteria: 10 points in total

- 1 point: 1-2 errors
- 2 point: 3-4 errors
- 3 point: 5-6 errors
- 4 point: 7-8 errors
- 5 point: 9-10 errors
- 6 point: 11-12 errors
- 7 point: 13-14 errors
- 8 point: 15-17 errors
- 9 point: 18-20 errors
- 10 point: Incorrect or no answer

- (b) The joint CDF looks like a staircase function that's easiest to represent as a grid. Remember that the joint CDF is the sum of all probability to the left and below a given point (indicated by small numbers at the dots); we accumulate probability every time we cross a line.



Grading criteria: 10 points in total

- 1 point: 1-2 errors
- 2 point: 3-4 errors
- 3 point: 5-6 errors
- 4 point: 7-8 errors
- 5 point: 9-10 errors
- 6 point: 11-12 errors
- 7 point: 13-14 errors
- 8 point: 15-17 errors
- 9 point: 18-20 errors
- 10 point: Incorrect or no answer

- (c) From the joint PMF in part (a), it's easy to compute the marginal PMF of Y just by summing up the rows; we obtain

	$p_Y(Y)$
0	$\frac{7}{384}$
1	$\frac{39}{384}$
2	$\frac{91}{384}$
3	$\frac{115}{384}$
4	$\frac{85}{384}$
5	$\frac{37}{384}$
6	$\frac{10}{384}$

As expected these probabilities sum to 1, since the marginal is a valid PMF.

Grading criteria: 5 points in total

-1 point: small error

-2 point: two errors

-3 point: 3-4 errors

-4 point: 5-6 errors

-2 point: 5-6 errors

-2 point: did not compute final answer

-10 point: Incorrect or no answer

2. (a) Let's integrate this function and see what we get:

$$\begin{aligned}
 \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g(x, y) \, dx \, dy &= k \int_0^1 \int_0^1 x(1-x)y \, dx \, dy \\
 &= k \left(\frac{1}{2}x^2 - \frac{1}{3}x^3 \right) \Big|_{x=0}^{x=1} \left(\frac{1}{2}y^2 \right) \Big|_{y=0}^{y=1} \\
 &= k \left(\frac{1}{6} \right) \left(\frac{1}{2} \right) \\
 &= \frac{k}{12}
 \end{aligned}$$

Since for a valid PDF we need this integral to equal 1, this means that $k = 12$.

Grading criteria: 5 points in total

- 0 point: Incorrect, but correct based on your part (a)
- 1 point: Small error
- 1.5 point: Computation error
- 2.5 point: The integration is incorrect
- 4 point: Incorrect, some efforts
- 5 point: Incorrect or no answer

- (b) The joint CDF in the “interesting” range $x \in [0, 1], y \in [0, 1]$ is computed as

$$\begin{aligned}
 F_{XY}(x, y) \, dx \, dy &= \int_0^x \int_0^y f_{XY}(x, y) \, dx \, dy \\
 &= 12 \left(\frac{1}{2}x^2 - \frac{1}{3}x^3 \right) \left(\frac{1}{2}y^2 \right) \\
 &= (3x^2 - 2x^3)y^2
 \end{aligned}$$

Grading criteria: 5 points in total

- 0 point: small error
- 2 point: Equation is correct, final answer is incorrect
- 3 point: Partially correct
- 4 point: Incorrect, some efforts
- 5 point: Incorrect or no answer
- 2 point: Incomplete process
- 1 point: Missing conclusion (what is the value of c?)

- (c) To compute the marginals, we integrate out the variable we don't care about. For $x \in [0, 1]$ we have

$$\begin{aligned}
 f_X(x) &= \int_0^1 12x(1-x)y \, dy \\
 &= (12x(1-x)) \left(\frac{1}{2}y^2 \right) \Big|_{y=0}^{y=1} \\
 &= (12x(1-x)) \left(\frac{1}{2} \right) \\
 &= 6x(1-x) \quad x \in [0, 1], 0 \text{ otherwise}
 \end{aligned}$$

Grading criteria: 5 points in total

- 1 point: Small error
- 2.5 point: The integration is incorrect
- 1.5 point: Did not specify the pdf at all values of x
- 4 point: Incorrect, some efforts
- 5 point: Incorrect or no answer

(d) Similarly for $y \in [0, 1]$ we have

$$\begin{aligned} f_Y(y) &= \int_0^1 12x(1-x)y \, dx \\ &= y \left(6x^2 - 4x^3 \right) \Big|_{x=0}^{x=1} \\ &= y(2) \\ &= 2y \quad y \in [0, 1], 0 \text{ otherwise} \end{aligned}$$

Grading criteria: 5 points in total

- 1 point: Small error
- 2.5 point: The integration is incorrect
- 1.5 point: Did not specify the pdf at all values of x
- 4 point: Incorrect, some efforts
- 5 point: Incorrect or no answer

(e) Yes, X and Y are independent since we can see that

$$\begin{aligned} f_{XY}(x, y) &= 12x(1-x)y \\ &= (6x(1-x))(2y) \\ &= f_X(x) f_Y(y) \end{aligned}$$

Grading criteria: 5 points in total

- 0 point: Incorrect, but correct based on your previous parts
- 2.5 point: X and Y are independent
- 2 point: No conclusion
- 4 point: Correct conclusion without reason
- 3.5 point: Correct conclusion with incorrect reason
- 4 point: Incorrect, some efforts
- 5 point: Incorrect or no answer

(f)

$$\begin{aligned} P(Y < \sqrt{X}) &= \int_0^1 \int_0^{\sqrt{x}} 12x(1-x)y \, dy \, dx \\ &= \int_0^1 (12x(1-x)) \left(\frac{1}{2}y^2 \right) \Big|_{y=0}^{y=\sqrt{x}} dx \\ &= \int_0^1 (12x(1-x)) \left(\frac{1}{2}x \right) dx \\ &= \int_0^1 6x^2(1-x) \, dx \\ &= 2x^3 - \frac{6}{4}x^4 \Big|_{x=0}^{x=1} \\ &= \frac{1}{2} \end{aligned}$$

Grading criteria: 5 points in total

-0 point: Incorrect, but correct based on your previous parts

-2 point: Computation error

-4 point: Incorrect, some efforts

-5 point: Incorrect or no answer

3. (a) We're told that X and Y are jointly Gaussian with the PDF

$$f_{XY}(x, y) = c \exp\left(-\frac{3}{64}(12x^2 - 80x + 3y^2 + 24y - 4xy)\right)$$

where c is an unknown constant. We are also told that the correlation coefficient $\rho = \frac{1}{3}$. We need to pattern-match this against the 2D Gaussian PDF of the form

$$f_{XY}(x, y) = \frac{1}{2\pi\sigma_x\sigma_y\sqrt{1-\rho^2}} \exp\left(\frac{-1}{2(1-\rho^2)} \left[\left(\frac{x-\mu_x}{\sigma_x}\right)^2 - 2\rho\left(\frac{x-\mu_x}{\sigma_x}\right)\left(\frac{y-\mu_y}{\sigma_y}\right) + \left(\frac{y-\mu_y}{\sigma_y}\right)^2 \right]\right)$$

We don't care about the constant in front; let's just expand the exponent, plugging in $\rho = \frac{1}{3}$:

$$\begin{aligned} &\frac{-1}{2(1-\rho^2)\sigma_x^2}x^2 - \frac{1}{2(1-\rho^2)\sigma_y^2}y^2 + \left(\frac{\mu_x}{(1-\rho^2)\sigma_x^2} - \frac{\rho\mu_y}{(1-\rho^2)\sigma_x\sigma_y}\right)x + \left(\frac{\mu_y}{(1-\rho^2)\sigma_y^2} - \frac{\rho\mu_x}{(1-\rho^2)\sigma_x\sigma_y}\right)y \\ &\quad + \frac{\rho}{(1-\rho^2)\sigma_x\sigma_y}xy + \text{constant} \\ &= \frac{-9}{16\sigma_x^2}x^2 - \frac{9}{16\sigma_y^2}y^2 + \left(\frac{9\mu_x}{8\sigma_x^2} - \frac{3\mu_y}{8\sigma_x\sigma_y}\right)x + \left(\frac{9\mu_y}{8\sigma_y^2} - \frac{3\mu_x}{8\sigma_x\sigma_y}\right)y + \frac{3}{8\sigma_x\sigma_y}xy + \text{constant} \end{aligned}$$

Matching up the coefficients on the x^2 and y^2 terms gives us

$$\frac{-9}{16\sigma_x^2} = \frac{-9}{16} \qquad \frac{-9}{16\sigma_y^2} = \frac{-9}{64}$$

which tells us that $\sigma_x = 1$ and $\sigma_y = 2$. As a sanity check we can see that the xy term also agrees.

Grading criteria: 10 points in total

- 2 point: Small error
- 2 point: Mess up variance and standard deviation
- 5 point: Incorrect, partial efforts for equation simplification
- 4 point: Missing final answer
- 7 point: Correct answer with no process shown
- 8 point: Incorrect, some efforts
- 9 point: Incorrect answer with no process
- 10 point: Incorrect or no answer

(b) Now that we know σ_x and σ_y we can look at the x and y terms:

$$\begin{aligned}\frac{9\mu_x}{8} - \frac{3\mu_y}{16} &= \frac{15}{4} \rightarrow 18\mu_x - 3\mu_y = 60 \\ \frac{9\mu_y}{32} - \frac{3\mu_x}{16} &= \frac{-9}{8} \rightarrow -6\mu_x + 9\mu_y = -36\end{aligned}$$

Solving this linear system gives $\mu_x = 3, \mu_y = -2$.

Grading criteria: 10 points in total

- 2 point: Small error
- 4 point: Missing final answer
- 5 point: Incorrect, partial efforts (some equations)
- 7 point: Correct answer with no process shown
- 7 point: Incomplete answer
- 8 point: Incorrect, some efforts
- 9 point: Incorrect answer with no process
- 10 point: Incorrect or no answer

4. (a) The correlation of X and Y is defined as $E(XY)$.

$$\begin{aligned}E(XY) &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xy f_{XY}(x, y) dx dy \\ &= \frac{6}{19} \int_0^2 \int_0^1 xy(x^2 + y^3) dy dx = \frac{6}{19} \int_0^2 \left[\frac{1}{2} x^3 y^2 + \frac{1}{5} x y^5 \right]_{y=0}^{y=1} dx \\ &= \frac{6}{19} \int_0^2 \left(\frac{1}{2} x^3 + \frac{1}{5} x \right) dx \\ &= \frac{6}{19} \left(\frac{1}{8} x^4 + \frac{1}{10} x^2 \right) \Big|_{x=0}^{x=2} \\ &= \frac{6}{19} \cdot \frac{12}{5} = \frac{72}{95}\end{aligned}$$

Grading criteria: 5 points in total

- 1 point: Small error
- 2.5 point: Incorrect, partial efforts
- 4 point: Incorrect, some efforts
- 5 point: Incorrect or no answer

- (b) The covariance of X and Y is defined as $E(XY) - E(X)E(Y)$, which means we first have to compute the marginal means $E(X)$ and $E(Y)$.

$$\begin{aligned}
 E(X) &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} x f_{XY}(x, y) \, dx \, dy \\
 &= \frac{6}{19} \int_0^2 \int_0^1 x(x^2 + y^3) \, dy \, dx = \frac{6}{19} \int_0^2 \left[x^3 y + \frac{1}{4} x y^4 \right]_{y=0}^{y=1} dx \\
 &= \frac{6}{19} \int_0^2 x^3 + \frac{1}{4} x \, dx \\
 &= \frac{6}{19} \left(\frac{1}{4} x^4 + \frac{1}{8} x^2 \right) \Big|_{x=0}^{x=2} \\
 &= \frac{6}{19} \cdot \frac{9}{2} = \frac{27}{19}
 \end{aligned}$$

$$\begin{aligned}
 E(Y) &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} y f_{XY}(x, y) \, dx \, dy \\
 &= \frac{6}{19} \int_0^2 \int_0^1 y(x^2 + y^3) \, dy \, dx = \frac{6}{19} \int_0^2 \left[\frac{1}{2} x^2 y^2 + \frac{1}{5} y^5 \right]_{y=0}^{y=1} dx \\
 &= \frac{6}{19} \int_0^2 \frac{1}{2} x^2 + \frac{1}{5} \, dx \\
 &= \frac{6}{19} \left(\frac{1}{6} x^3 + \frac{1}{5} x \right) \Big|_{x=0}^{x=2} \\
 &= \frac{6}{19} \cdot \frac{26}{15} = \frac{52}{95}
 \end{aligned}$$

Thus

$$\text{Cov}(X, Y) = \frac{72}{95} - \frac{27}{19} \cdot \frac{52}{95} = \frac{-36}{1805}$$

Grading criteria: 10 points in total

- 0 point: Incorrect, but correct based on your part (a)
- 2 point: Computation error
- 3 point: $E(X)$ is incorrect
- 3 point: $E(Y)$ is incorrect
- 4 point: Errors in expectations
- 4 point: Incorrect or missing calculation of covariance
- 5 point: Incorrect, partial efforts for integration
- 7 point: Incomplete answer
- 8 point: Incorrect, some efforts
- 9 point: Only provide a general equation
- 10 point: Incorrect or no answer

- (c) Now we use the properties of expected value and the numbers we already computed:

$$\begin{aligned}
 E(95(X(1 + Y) + 2Y(1 - X))) &= 95E(X + 2Y - XY) \\
 &= 95(E(X) + 2E(Y) - E(XY)) \\
 &= 95 \left(\frac{27}{19} + \frac{104}{95} - \frac{72}{95} \right) = 167
 \end{aligned}$$

Grading criteria: 5 points in total

-0 point: Incorrect, but correct based on your part (a)

-1 point: Small error

-2.5 point: Incorrect, partial efforts (equations)

-4 point: Incorrect, some efforts

-4 point: Incomplete answer

-5 point: Incorrect or no answer

(d) No, X and Y are not uncorrelated since $E(XY) \neq 0$.

Grading criteria: 5 points in total

-0 point: Incorrect, but correct based on your part (a)

-3 point: Correct with no justification

-3 point: Correct answer with wrong justification

-5 point: Incorrect or no answer