

Name: Hayden Fuller

RCS ID: fulleh @rpi.edu

CSCI 2200 — Foundations of Computer Science (FoCS) — Exam 1

Sign below to confirm that you will not copy and will not cheat on this exam, which also means you will not communicate with anyone under any circumstances about this exam until it is graded.

Signature: Hayden Fuller

Any copying or collaborating with others will result in a grade of ZERO on this exam.

Q1-18 (2 POINTS EACH) — Mark one answer for each question....

34

Q1	(a)	(b)	(c)	(d)	(e)
Q2	(a)	(b)	(c)	(d)	(e)
Q3	(a)	(b)	(c)	(d)	(e)
Q4	(a)	(b)	(c)	(d)	(e)
Q5	(a)	(b)	(c)	(d)	(e)
Q6	(a)	(b)	(c)	(d)	(e)
Q7	(a)	(b)	(c)	(d)	(e)
Q8	(a)	(b)	(c)	(d)	(e)
Q9	(a)	(b)	(c)	(d)	(e)

Q10	(a)	(b)	(c)	(d)	(e)
Q11	(a)	(b)	(c)	(d)	(e)
Q12	(a)	(b)	(c)	(d)	(e)
Q13	(a)	(b)	(c)	(d)	(e)
Q14	(a)	(b)	(c)	(d)	(e)
Q15	(a)	(b)	(c)	(d)	(e)
Q16	(a)	(b)	(c)	(d)	(e)
Q17	(a)	(b)	(c)	(d)	(e)
Q18	(a)	(b)	(c)	(d)	(e)

I forgot my log rules...

Q19 (4 POINTS)

Proof. we prove the claim by contradiction
 Assume $\log_2 9$ is a rational number, $\log_2 9 = \frac{a}{b}$ $a \neq 0, b \in \mathbb{N}$, a, b are integers with no common factors.
 $9 = 2^{\frac{a}{b}} = 2^a \cdot 2^{-\frac{a}{b}}$

Q20 (5 POINTS)

Proof. We use strong induction to prove $n^2 - 1$ is divisible by 8 for all odd $n \in \mathbb{N}$.

[base cases] For $P(1)$ we have $1^2 - 1 = 0$ is divisible by 8, which is true.

And for $P(3)$ we have $3^2 - 1 = 8$ is divisible by 8, which is true.

[induction step] We show $P(n) \rightarrow P(n+4)$ using a direct proof. We must prove $P(n+4)$, i.e. $(n+4)^2 - 1$ is divisible by 8. Assume $n^2 - 1$ is divisible by 8.

$$\text{LHS } (n+4)^2 - 1 = n^2 + 8n + 16 - 1$$

Since n is odd, let $n = 2k+1$ for $k \in \mathbb{N}_0$.

$$\text{LHS } (n+4)^2 - 1 = (2k+1)^2 + 8(2k+1) + 16 - 1 = 4k^2 + 4k + 1 + 16k + 8 + 16 - 1 = 4k^2 + 4k + 8(2k+1) + 16$$

$$\text{if } k \text{ is even, let } k = 2j. 4(2j)^2 + 4(2j) = 4 \cdot 4j^2 + 8j = 8(2j^2 + j)$$

$$\text{if } k \text{ is odd, let } k = 2j+1. 4(2j+1)^2 + 4(2j+1) = 4(4j^2 + 4j + 1) + 4(2j+1) = 4(4j^2 + 6j + 2) = 8(2j^2 + 3j + 1)$$

therefore $4k^2 + 4k$ is divisible by 8, and can be written as $8(m)$.

$$(n+4)^2 - 1 = 8m + 8(2k+1) + 16 = 8(m + 2k + 1 + 2), \text{ so } (n+4)^2 - 1 \text{ is divisible by 8, and } P(n) \rightarrow P(n+4)$$

therefore, by induction $n^2 - 1$ is divisible by 8 for all odd $n \in \mathbb{N}$. ■

Q21 (5 POINTS)

Proof. We use strong induction to prove $P(n)$.

[base case] For $P(2)$ we have $\frac{1}{2} + \frac{1}{2} + \frac{1}{2} = 1$.

[induction step]