

Binomial Distributions

Dr. Bashar Al-Shboul

The University of Jordan

- A **Bernoulli Trial** is a random. Experiment with the following features:
 - The outcome can be classified as either a success or a failure (only two options and each is mutually exclusive).
 - The probability of success is p and probability of failure is $q = 1 - p$.
- A **Bernoulli random variable** is a variable assigned to represent the successes in a Bernoulli trial.
- If we wish to keep track of the number of successes that occur in repeated Bernoulli trials, we use a **binomial random variable**.
- A **binomial experiment** occurs when the following conditions are met:
 - Each trial can result in one of only two mutually exclusive outcomes (success or failure).
 - There are a fixed number of trials.
 - Outcomes of different trials are independent.
 - The probability that a trial results in success is the same for all trials.

- The random variable X = number of successes of a binomial experiment is a binomial distribution with parameters p and n where p represents the probability of a success and n is the number of trials.
- The possible values of X are whole numbers that range from 0 to n . As an abbreviation, we say

$$X \sim B(n, p).$$

- Binomial probabilities are calculated with the following formula:

$$P(X = k) = \binom{n}{k} p^k (1-p)^{n-k} = {}_n C_k p^k (1-p)^{n-k}$$

Binomial Distribution using MATLAB

- Binopdf (X,N,P) – Binomial Probability Density Function
- Computes the binomial pdf at each of the values of X using the corresponding number of trials in N and probability of success for each trial in P.
- N must be positive integers
- P falls in the interval [0, 1]

- The mean and variance of a binomial distribution are computed using the following formulas :
- $\mu_x = E[X] = np$
- $\sigma_x^2 = np(1-p)$

A fair coin is flipped 30 times. $N=30$, $p = 1/2$

- a. What is the probability that the coin comes up heads exactly 12 times?
 - $p(X=12) = {}_{30}C_{12} (1/2)^{12} (1-1/2)^{(30-12)} = 0.0806$
- b. What is the probability the coin comes up heads less than 12 times ?
 - $P(X < 12) = P(X \leq 11) = 0.0509$
- c. What is the probability the coin comes up heads more than 12 times?
 - $P(X > 12) = 1 - P(X \leq 12) = 1 - \text{Binomial}(30, 0.5, 12) = 0.9194$
- d. What is the expected number of heads?
 - $E[X] = np = 30 (1/2) = 15$

Suppose it is known that 80% of the people exposed to the flu virus will contract the flu. Out of a family of five exposed to the virus, what is the probability that:

- No one will contract the flu ?
 - $P(X=0) = {}_5C_0 (0.8)^0 (1-0.8)^{(5-0)} = 0.00032$
- All will contract the flu ?
 - $P(X=5) = {}_5C_5 (0.8)^5 (1-0.8)^{(5-5)} = 0.3277$
- Exactly two will get the flu ?
 - $P(X=2) = {}_5C_2 (0.8)^2 (1-0.8)^{(5-2)} = 0.0512$
- At least two will get the flu ?
 - $P(X \geq 2) = 1 - P(X \leq 1) = 0.9936$

- Using the binomial distribution in example 2,
- a. Let X = number of family members contracting the flu. Create the probability distribution table of X .

X	0	1	2	3	4	5
P(X)	0.00032	0.0064	0.0572	0.2048	0.4096	0.32768

- b. Find the mean and variance of this distribution.
 - $E[X] = 0(0.00032) + 1(0.0064) + 2(0.0572) + 3(0.2048) + 4(0.4096) + 5(0.32768)$
 $= np = 5 (0.8) = 4$
 - $\sigma_x^2 = np(1-p) = 5 (0.8)(0.2) = 0.8$