

---

# SEASONALLY VARIABLE THERMAL PERFORMANCE CURVES PREVENT ADVERSE EFFECTS OF HEATWAVES

---

A PREPRINT

**Matthew C. Sasaki**

Department of Marine Sciences  
University of Connecticut  
Groton, CT 06340  
matthew.sasaki@uconn.edu

**Michael Finiguerra**

Ecology and Evolutionary Biology Department  
University of Connecticut  
Groton, CT 06340

**Hans G. Dam**

Department of Marine Sciences  
University of Connecticut  
Groton, CT 06340

August 30, 2022

## Abstract

The increasing frequency and intensity of heatwaves may represent a significant challenge for organisms in a warming ocean. The direct impacts of heatwaves on populations depend on the relative position of environmental temperatures to the thermal performance curve optima. The effects of heatwaves may therefore vary seasonally along the annual temperature cycle. However, this seasonal variation in the effects of heatwaves may be dampened by corresponding variation in thermal performance curves. In organisms with relatively short generation times, these changes may be driven by phenotypic plasticity as well as genetic differentiation. We investigate the effects of seasonal timing and duration on the impacts of heatwaves in the ecologically important copepod *Acartia tonsa*. We show that thermal performance curves are seasonally variable in the field, and that this variation buffers against negative effects of simulated heatwaves. Further, the offspring of individuals that experienced the simulated heatwaves were raised in the laboratory to examine trans-generational effects of heatwaves on body size and reproductive output. The lack of a clear pattern in the trans-generational effects may indicate that seasonal variation in thermal performance curves also buffers against indirect effects of heat waves by reducing the effects of parental stress on the offspring. Our results show that seasonal variation in thermal performance curves has the potential to limit the adverse effects of heatwaves on populations of short-lived organisms.

**Keywords** Heat wave · Climate change · Copepod · Transgenerational plasticity · Seasonal variation · Thermal performance

## Introduction

Heatwaves are increasing in frequency and intensity in marine ecosystems (Frölicher et al. 2018; Oliver et al. 2018, 2019). These periods of anomalously high temperatures present severe challenges to marine biota (Smale et al. 2019), and both lethal and non-lethal temperature effects on organisms have strong potential to alter community composition and ecosystem functions (REF). The effects on community dynamics are determined in part by the relative vulnerabilities of component taxa (REF). In order to predict how




Figure 1: Caption

communities may respond to both long-term warming and the increasing effects of heatwaves, there is an urgent need to understand what determines relative vulnerability to heatwaves.

Most previous studies have focused on the direct effects of increased temperature on organisms and population dynamics (REF). Shaped by resulting changes in feeding, competitive ability, reproduction, and survival, increased temperature during a heatwave may have profound effects on community dynamics (REF) and the distribution of organisms (REF). There may also be a wide range of indirect effects of heatwaves on populations, propagated by mechanisms like trans-generational plasticity and maternal effects (REF). As both the frequency and duration of heatwaves increases, these indirect effects may be especially important to consider (papers on the benefits of trans-generational effects for events that last a long time - REF).

In marine communities, autotrophs (e.g. phytoplankton) and primary consumers (e.g. copepods) typically have generation times on the order of days to weeks. As such, the response of these key groups to heatwaves may vary over seasonal timescales, reflecting either acclimatization or adaptation of thermal performance curves to different seasonal conditions (Sasaki and Dam 2020; Adaptive tracking paper). The presence or absence of seasonal variation in thermal performance curves will play an important role in determining the effects of heatwaves on a population (Figure 1). Much of the past work on the effects of heatwaves assumes performance curves are fixed (i.e. - there is no seasonal variation in TPCs). In this case, the effect of heatwaves will vary seasonally as the relative position of environmental temperatures to TPC optimum temperature and lethal thermal limits varies. Alternatively, variation in TPCs over seasonal timescales may act as a buffer against negative effects of heatwaves if optimum temperatures track environmental temperatures, preventing environmental temperatures from exceeding thermal optima. Differences in the magnitude of seasonal variation in TPCs may therefore play a key role in shaping patterns in relative vulnerability of different taxa within a community.

Seasonal variation in TPCs could modify both direct and indirect effects of heatwaves on marine organisms with short generation times, and thus shape the relative vulnerability of different taxa. It is therefore important to consider this temporal variation in predictions about how the increasing threat of heatwaves may affect marine communities. Copepods are an ideal model system to examine these dynamics, as they dominate the planktonic community in many temperate locations and typically have short generation times. By nature of their abundance, copepods are exceptionally ecologically important, and as a result there is pressing need to understand how these organisms may be affected by marine heatwaves. Here we investigate the potential for seasonal variation in thermal performance curves of the ecologically important copepod species *Acartia hudsonica* and *A. tonsa* to buffer populations against negative effects of heatwaves. We examined how thermal performance curves for key fitness related traits (egg production, hatching success, and survivorship)

varied over the course of the seasonal temperature cycle. We also quantified both direct and indirect (e.g. -trans-generational) effects of heatwaves on one of the species (*A. tonsa*) in the laboratory using a series of simulated heatwave experiments. We test two hypotheses: 1) there is seasonal variation in the thermal performance curves of key fitness related traits in *A. hudsonica* and *A. tonsa*; and 2) when TPC parameters track changes in environmental temperatures, the effects of simulated heatwaves do not vary over the course of the year. By integrating experiments with field-collected and lab-reared organisms, we show that TPCs of *A. tonsa* but not *A. hudsonica* vary across seasons and that this variation allows the population to maintain a relatively constant margin between optimum temperatures and ambient environmental temperatures. The mechanisms that produce seasonally variable TPCs, whether genetic or plastic, reduced the potential for heatwaves to adversely affect population dynamics and thus increase resilience in the face of climate change.

## Methods

### Generating Field TPCs

Copepods were collected from Long Island Sound, near the University of Connecticut, Avery Point Campus at several times throughout the year using surface tows of a 200 µm mesh plankton net. Mature *Acartia hudsonica* or *Acartia tonsa* females were isolated from the contents of the plankton tow and isolated individually in petri dishes for egg production and hatching success assays. During these assays, females were fed *Scenedesmus*. Females produced eggs for 3 days, after which the female was removed. Eggs were given three additional days before hatching frequency was assessed. These assays were performed across a range of temperatures (10-30 degrees C for *A. tonsa* and 4-24 degrees C for *A. hudsonica*).

**Will need substantial input from other people to fill in the methods section.**

Table 1: Caption

Month	Date	Temperature	Species
January	January 21st 2015	2.1	<i>Acartia hudsonica</i>
February	February 21st 2015	-0.4	<i>Acartia hudsonica</i>
March	March 16 2015	4.0	<i>Acartia hudsonica</i>
April	April 21st 2015	9.2	<i>Acartia hudsonica</i>
May	May 14th 2015	12.0	<i>Acartia hudsonica</i>
June	June ____ 2015	18.6	<i>Acartia hudsonica</i>
July	July 29th 2014	20.0	<i>Acartia tonsa</i>
August	August 13th 2014	18.0	<i>Acartia tonsa</i>
September	September 11th 2014	18.0	<i>Acartia tonsa</i>
October	October 22nd 2014	16.0	<i>Acartia tonsa</i>
November 1	November 4th 2014	14.5	<i>Acartia tonsa</i>
November 2	November 19th 2014	11.0	<i>Acartia tonsa</i>

TPCs for egg production, hatching success, and production were then generated using the framework outlined by Padfield et al. (2021). Egg production, hatching success, and production data were modeled using a Gaussian equation. Parameters (maximum values and thermal optima) were extracted from these model fits. Survivorship was modeled using a logistic regression of individual survivorship against stress temperature and used to estimate thermal tolerance (as LD50, the temperature of 50% mortality).

### Simulated Heat Waves

To test our second hypothesis (that varying thermal performance curve parameters reduce vulnerability to heatwaves) we collected *Acartia tonsa* again in 2015 for use in laboratory simulated heatwave experiments. In order to test the effects of a heatwave against the seasonally shifting baseline of ambient temperature, three collections were made before, during, and after peak environmental temperatures, corresponding to late June, late July, and early December (Table 2). These experiments examined both direct and indirect (e.g. transgenerational) effects of heatwaves. To examine the direct effects, egg production rate, hatching success, and production were measured for around 60 females per collection split into two groups (control and heatwave). These assays were performed as described in the Field TPC section, with females isolated in individual petri dishes and provided with food *ad libitum*. The control group remained at a temperature near

the current ambient temperature in Long Island Sound while the Heatwave group experienced temperatures 5 degrees C above the ambient temperature (Table 3). Traits were measured over two periods to examine the effects of short and long heatwave events - Days 1-3 and Days 5-7, respectively. Females were moved into petri dishes with fresh food solution on Days 3 and 5. Eggs produced in between the two time periods were not examined.

[STATS]

Table 2: Caption

Period	Date	Field Temp.	Control Temp.	Heatwave Temp.
Early	June 27th 2015	17.8	17	22
Mid	July 29th 2015	22.4	22	27
Late	December 1st 2015	12.0	12	17

### Transgenerational Experiments

In addition to the individuals maintained in petri dishes to measure direct effects of heatwaves, several hundred adult copepods were also placed into each of eight 4L buckets of filtered seawater and provided with food ad libitum (four each, maintained at control and heatwave temperatures). Water in each bucket was kept oxygenated using a small aquarium pump. Eggs were collected from each bucket following the same schedule as the direct effect experiments - eggs were collected on Day 3 and Day 7 for the short and long heatwave exposures, respectively. Eggs were discarded on Day 5 to ensure all individuals reflected the correct exposure periods. These eggs were then split into three groups which developed at one of three different temperatures (12, 17, or 22 degrees C). After these individuals matured, body size and the three reproductive traits (egg production, hatching success, and production) were measured at the temperature individuals developed at. Within developmental temperatures, differences between the Heatwave and Control treatment groups should reflect the indirect effects of heatwaves: these individuals represent eggs produced by parents experiencing the heatwave conditions, but that developed at the same temperatures as the Control individuals.

[STATS]

## Results

### Seasonal Variation in Field TPCs

There was abundant variation in TPCs for Egg Production Rate (EPR), Hatching Success, and production (the product of Hatching Success and Total Egg Production; the number of hatched offspring per female per day) for copepods collected throughout the year.

[EPR] *Acartia hudsonica* EPR TPCs tended to have lower optima and were less variable than *A. tonsa* EPR TPCs. Within *A. tonsa*, EPR optimum temperatures were higher in warmer months (July, August, and September) than in cooler months (October and November).

[HS] Peaks were generally less distinct for hatching success in *A. tonsa* than in *A. hudsonica*. In general, *A. tonsa* hatching success was generally higher in warmer months than cooler months, regardless of incubation temperature.

[Production] When combined, the variation in optimum temperatures and maximum values for EPR and hatching success curves yielded Production curves that were highly variable. Collections from warmer months generally had slightly higher optimum temperatures as well as higher maximum production values.

[Survivorship] Thermal survivorship curves were highly variable, with a LD50 range of  $>_{-}$  in both species. As observed in previous studies, *A. tonsa* had higher thermal tolerance than *A. hudsonica* (Sasaki & Dam, 2020). Thermal tolerance increased with collection temperature in both species.

[Thermal safety margin] Vulnerability to heatwaves was estimated as thermal safety margins (difference between thermal optimum and environmental temperature) and warming tolerance (difference between thermal tolerance and environmental temperature) (REF). TPC parameters for *A. tonsa* but not *A. hudsonica* were correlated with collection temperature. The invariant thermal optimum for *A. hudsonica* production results in seasonal variation in the effect of heatwaves, as illustrated by the decline in thermal safety margin




Figure 2: caption

for this species. Later in its season of occurrence, environmental temperatures actually exceed the thermal optimum, suggesting that the seasonal distribution of this species may be directly affected by temperature, and that heatwaves occurring in the late Spring and early Summer could have strong detrimental effects. By contrast, the seasonally variable *A. tonsa* production TPCs resulted in relatively stable thermal safety margins. Throughout its season of occurrence, *A. tonsa* maintained a buffer of at least 5 degrees C, suggesting that at all times a mild heatwave would increase production by pushing environmental temperatures closer to the thermal optimum. The one outlier was the second November collection, which was collected at 11 degrees C. This is around the threshold for resting egg production in *A. tonsa*. The extremely high estimated production optimum temperature may reflect the difference in hatching requirements between resting and subitaneous eggs.

[Warming tolerance] Thermal tolerance increased with collection temperature in both temperatures, but this trend was not significant (STATS). Despite this, warming tolerance decreased as collection temperature increased, indicating imperfect compensation via either acclimatization or adaptation. While collection temperatures never exceeded thermal tolerance for either species, the lower thermal tolerance of *A. hudsonica* translated to reduced warming tolerance relative to *A. tonsa*, again suggesting that the seasonal occurrence of this species may be delimited at least in part by direct effects of temperature.

### Effects of Simulated Heatwaves

The second component of this project examined the effects of heatwaves across generations. This necessarily began with reassessing the field collected F0. We examined the effects of short or long duration artificial heatwaves, during different collection months on EPR, HS, and production. We will focus the analysis on production, as it integrates the other traits. The ANOVA indicated no significant overall effect of heatwaves on




Figure 3: caption

production ( $p$ -value = 0.38). There was, however, a significant interaction between treatment and collection month ( $p$ -value = 0.01), indicating different effects of heatwaves across months. The different trial duration and collection months were also significantly different, with a significant interaction term as well. ANOVA results are shown in Supp. Table 1.

The effect sizes (the difference between heatwave and control trials, or between long and short duration trials) and confidence intervals were estimated using non-parametric bootstrap resampling. These estimates are shown below for the effects of treatment and the effect of duration. Short and long duration trials are shown in white and black symbols, respectively, in the first panel. The control and heatwave treatments are shown with different shapes in the second panel. Full Gardner-Altman estimation plots are shown in Supp. Fig 1 and 2. Effects of heatwaves were generally weak; only long heatwaves in June and short heatwaves in August had confidence intervals that did not (or nearly did not) overlap zero. In contrast, there were strong decreases in long duration trials relative to short trials, in both control and heatwave treatments.

### Transgenerational Effects of Simulated Heatwaves

We also examined the effects of parental exposure to heatwaves on offspring traits. Comparing between Control and Heatwave treatments now examines not the direct effects of increased temperature, but the indirect effect on offspring of parental exposure to heatwaves. In both panels, the gray boxes indicate the developmental temperature most similar to that experienced by offspring of the parental generation in the field.




Figure 4: caption

**Body Size** - Offspring body size generally decreased with developmental temperature, as expected (Supp. Fig. 4). Within individual developmental temperatures, parental exposure to heatwaves also generally resulted in small decreases in body size. Short and long duration events are shown with open and filled circles respectively.

**Fecundity** - There were no consistent patterns in the effects of parental exposure to heatwaves across developmental temperatures, heatwave duration, or time of year.

## Discussion

We have shown that there is seasonal variation in the TPC for multiple fitness-related traits in the Long Island Sound population of *A. tonsa*, but not *A. hudsonica*. The different levels of seasonal variation result in different relative vulnerabilities of the two species to heatwaves; stronger seasonal variation in *A. tonsa* prevents decreases in the thermal safety margin throughout its season of occurrence, whereas the thermal safety margin shrinks drastically in less seasonally-variable *A. hudsonica*.

[Simulated heatwave summary]

Trans-generational effects of heatwaves were highly variable. Parental exposure to heat waves generally decreased offspring body size, but had no consistent effects on F1 production rates. Effects were particularly small when looking just at the developmental temperatures the offspring would experience in the field - At these temperatures, only parental exposure to heatwaves in August resulted in a decrease in offspring production or reduced body sizes.

As a result, we'd predict heat waves should have a beneficial effect, regardless of seasonal timing, by moving the population towards its optimum temperature. However, strong heatwaves during the warmest times may have an adverse effect on survivorship, unless other mechanisms (e.g. - acclimation and phenotypic plasticity) adjust thermal tolerance.

The expectation is that production should decrease as body size decreases due to the effects on EPR. However, the changes in production were independent of changes in body size (Supp. Fig. 6). Instead, the observed effects are more likely to be the result of other mechanisms (such as maternal effects or transgenerational plasticity).



## Supplemental Information

Supp. Table 1


Supp. Table 2

Supp. Table 3


Supp. Fig. 1

These plots follow best practices for the visualization of differences between groups. The top half of each figure shows the underlying data points in a swarm plot. To the right of each each swarm is the mean and standard deviation of the group, represented using a gapped bar (gap = mean value). Below the raw data, the effect size and 95% confidence intervals are shown, which were obtained using non-parametric bootstrap resampling. Confidence intervals that do not cross the 0 effect size line indicate significant differences between groups.

### Short Heat Waves




### Long Heat Waves




## Supp. Fig. 2

Effects of parental exposure to heatwaves on F1 body size at different developmental temperatures.




## Supp. Fig. 3

Effects of parental exposure to heatwaves on F1 egg production at different developmental temperatures.



## Supp. Fig. 4

Effects of parental exposure to heatwaves on F1 hatching success at different developmental temperatures.



## Supp. Fig. 5

Effects of parental exposure to heatwaves on F1 production at different developmental temperatures.

