
A
Seasonal GLM: count of SNPs p < 0.001
Temporal Fc: excess above observed
SNPs under selection
10% 0.0 0.1 0.7 1.0 1.0 1.0 1.0 1.0
5% 0.0 0.0 0.4 0.8 1.0 1.0 1.0 1.0
2% 0.0 0.0 0.0 0.7 1.0 1.0 1.0 1.0
1% 0.0 0.0 0.0 0.2 0.9 1.0 1.0 1.0
0.5% 0.0 0.0 0.0 0.1 0.4 1.0 1.0 1.0
0.1% 0.0 0.0 0.0 0.0 0.0 0.5 1.0 1.0
0.0 0.0 0.0 1.0 1.0 1.0 1.0 1.0
0.0 0.0 0.0 0.0 1.0 1.0 1.0 1.0
0.0 0.0 0.0 0.0 0.0 1.0 1.0 1.0
0.0 0.0 0.0 0.0 0.0 0.0 0.2 1.0
0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0
0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0
X
10% 0.0 0.1 0.7 1.0 1.0 1.0 1.0 1.0
5% 0.1 0.0 0.2 1.0 0.9 1.0 1.0 1.0
2% 0.0 0.0 0.0 0.7 0.9 1.0 1.0 1.0
1% 0.0 0.0 0.0 0.3 0.8 1.0 1.0 1.0
0.5% 0.0 0.0 0.0 0.2 0.4 1.0 1.0 1.0
0.1% 0.0 0.0 0.0 0.0 0.0 0.2 0.8 1.0
0.0 0.0 0.0 1.0 1.0 1.0 1.0 1.0
0.0 0.0 0.0 0.0 1.0 1.0 1.0 1.0
0.0 0.0 0.0 0.0 0.0 0.0 1.0 1.0
0.0 0.0 0.0 0.0 0.0 0.0 0.0 1.0
0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0
0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0
F
Power
1.00
0.75
0.50
0.25
0.00
2 pp 6 pp 12 pp 18 pp 23 pp 32 pp 38 pp 45 pp
Seasonal change in allele frequency at selected SNPs (from 50%, first to last sample)
B
Temperature−associated SNPs
X
F
top 100% – 1 pp conc 3 pp conc 7 pp conc 13 pp conc 24 pp
top 20% – 3 pp conc 5 pp conc 13 pp conc 25 pp conc 40 pp
top 5% – 3 pp conc 7 pp conc 17 pp conc 30 pp conc 44 pp
top 1% – 4 pp conc 9 pp conc 20 pp conc 35 pp conc 47 pp
conc 5 pp conc 10 pp conc 23 pp r 38 pp r 48 pp
conc 5 pp conc 10 pp conc 24 pp r 39 pp – 48 pp
conc 6 pp conc 12 pp conc 27 pp – 42 pp – 49 pp
conc 7 pp conc 14 pp conc 31 pp – 45 pp – 50 pp
. = 0.05 . = 0.1 . = 0.25 . = 0.5 . = 1
not detected
detected
Seasonal response as a fraction of the spatial slope
Cells: statistic ranking first (conc: sign agreement, top 1%; r: correlation) and implied seasonal change.