
A
X
F
p < 0.01
p < 0.001
SNPs below threshold
200
150
100
50
0
15
10
5
0
time
temperature
time
temperature
Seasonal model
B
Seasonal GLM (SNPs at p < 0.001)
Temporal Fc (excess above observed)
SNPs under selection
10%
5%
2%
1%
0.5%
0.1%
2 pp
6 pp
12 pp
18 pp
23 pp
32 pp
38 pp
45 pp
Power
1.00
0.75
0.50
0.25
0.00
Seasonal change in allele frequency at selected SNPs
(percentage points from 50%, first to last sample)
C
X
F
Temperature−associated SNPs
top 100%
top 20%
top 5%
top 1%
– 1 pp
conc 3 pp
conc 7 pp
conc 13 pp
conc 24 pp
conc 5 pp
conc 10 pp
conc 23 pp
r 38 pp
r 48 pp
– 3 pp
conc 5 pp
conc 13 pp
conc 25 pp
conc 40 pp
conc 5 pp
conc 10 pp
conc 24 pp
r 39 pp
– 48 pp
– 3 pp
conc 7 pp
conc 17 pp
conc 30 pp
conc 44 pp
conc 6 pp
conc 12 pp
conc 27 pp
– 42 pp
– 49 pp
– 4 pp
conc 9 pp
conc 20 pp
conc 35 pp
conc 47 pp
conc 7 pp
conc 14 pp
conc 31 pp
– 45 pp
– 50 pp
not detected
detected
. = 0.05
. = 0.1
. = 0.25
. = 0.5
. = 1
Seasonal response as a fraction of the spatial slope