Physic formulary

School

2022.02.10

Contents

$e = 1.6021766\mathrm{C}$	last text
$\varepsilon_0 = 8.8541878\frac{\mathrm{F}}{\mathrm{m}}$	more text
$\varepsilon_{\mathtt{r}} = 1.00059$	sometext
$F = 96485.3 \frac{\text{C}}{\text{mol}}$	Faraday const

1 Test

$e = 1.6021766\mathrm{C}$	last text
$\varepsilon_0 = 8.8541878\frac{\mathrm{F}}{\mathrm{m}}$	more text
$\varepsilon_{\mathtt{r}} = 1.00059$	sometext
$F = 96485.3 \frac{C}{mol}$	Faraday const

2 TEst2

asdf

3 Constants

 $a_0 = \frac{4\pi\varepsilon_0\hbar^2}{m_{\rm e}e^2} = \frac{\varepsilon_0h^2}{\pi m_{\rm e}e^2} = \frac{\hbar}{m_{\rm e}c\alpha} = 5.291\,772\,109\,03 \times 10^{-11}\,{\rm m}$ Bohr radius

 $= 299792458 \frac{m}{s}$ Velocity of light:

 $= 1.602176634 \times 10^{-19} \,\mathrm{C}$ Elementary charge: $\varepsilon_0 = 8.8541878128 \times 10^{-12} \frac{F}{m}$ Vacuum permittivity:

Permittivity of air: = 1.00059 $\varepsilon_{\mathtt{r}}$

 $= 96485.3321233\frac{C}{mol}$ Faraday constant

 $= 9.80665 \frac{\text{m}}{\text{s}^2}$ Acceleration due to gravity:

 $= 6.67430 \times 10^{-11} \, \frac{\mathrm{m}^3}{\mathrm{kg \, s}^2}$ GGravitational constant: $= 6.626\,070\,15 \times 10^{-34}\,\tfrac{\mathrm{J}}{\mathrm{Hz}}$ Planck constant: $= 1.280649 \times 10^{-23} \frac{J}{K}$ Boltzmann constant:

 $m_e = 9.1093837015 \times 10^{-31} \,\mathrm{kg}$ Electron mass:

 $m_{\rm e} = 0.510\,998\,950\,00\,\frac{\rm MeV}{{\rm c_0}^2}$

 $m_{\mu} = 1.883531627 \times 10^{-28} \,\mathrm{kg}$

 $m_{\mu} = 105.6583755 \frac{\text{MeV}}{\text{co}^2}$ Muon mass:

 $m_{\mu} = 0.1134289259 \,\mathrm{Da}$

 $m_{\rm n} = 1.674\,927\,498\,04 \times 10^{-27}\,{\rm kg}$ Neutron mass:

 $m_{\rm n} = 939.56542052 \, \frac{\rm MeV}{{\rm c_0}^2}$

 $m_{\rm p} = 1.67262192369 \times 10^{-27} \,{\rm kg}$

Proton mass: $m_{\rm p} = 938.272\,088\,16\,{{\rm MeV}\over{{\rm c_0}^2}}$

 $\mu_{\rm 0} \ = \ 1.256\,637\,062\,12\times 10^{-6}\,{\textstyle\frac{\rm H}{\rm m}}$ Vacuum permeability:

Permeability of air: = 1.00000037 μ_{r}

4 Other physical interrelationships

Visible spectrum: $380\,\mathrm{nm}$ to $750\,\mathrm{nm}$

Speed of sound under standart conditions: $343\,\frac{\mathrm{m}}{\mathrm{s}}$

Dalton Da / unified atomic mass unit $\mathrm{u}\colon \mathrm{Da/u} =$

 $1.660\,539\,066\,60\times10^{-27}\,\mathrm{kg}$

Hydrogen mass: $m_{
m H} =$

1.00784Da to 1.00811Da

Atomic mass of helium ${}^4\mathrm{He}$ $m_{\mathrm{He}} = 4.002\,603\,254\,\mathrm{Da}$

Kilowatt-hour: $kW\,h$ = $3.6 \times 10^6\,J$

Kilowatt-hour: $eV = 1.602176634 \times 10^{-19} \,\mathrm{J}$

Pressure: $1 \, \mathrm{Pa} \qquad = 1 \, \frac{\mathrm{N}}{\mathrm{m}^2}$ Pressure: $1 \, \mathrm{bar} \qquad = 10^5 \, \mathrm{Pa}$ Absolute zero: $-273.15 \, ^{\circ}\mathrm{C} \qquad = 0 \, \mathrm{K}$

2

5 Energy

Kinetic Energy

$$E_{\mathbf{k}} = \frac{1}{2}mv^2$$

Potential Energy

$$U=mgh$$
 (gravitational) $U=rac{1}{2}\cdot k\cdot x^2$ (elastic) $U=rac{1}{2}\cdot C\cdot V^2$ (electric) $U=-mB$ (magnetic) $U=\int F(r)\,dr$ (general)

6 Motion

uniform linear motion

$$s(t) = vt \ (+s_0)$$

$$v(t) = const.$$

$$a(t) = 0$$

non-uniform linear motion

$$s(t) = \frac{1}{2}at^2 (+v_0 t + s_0)$$

$$v(t) = at \, (+v_0)$$

$$a(t) = const.$$

Circular motion

$$F_{
m Z}=rac{mv^2}{r}=m\,\omega^2 r$$
 $\omega=rac{v}{r}=rac{2\pi}{T}=rac{\Deltaarphi}{\Delta t}$ $arphi$ in rad $f=rac{1}{T}$

Momentum 7

momentum itself

$$\vec{p} = m\vec{v}$$

$$\vec{p} = \frac{\hbar}{\lambda}$$

$$\vec{p} = \sqrt{m_0^2 c_0^2 + \frac{E^2}{c_0^2}}$$

photons

general

relations to momentum

$$\sum_{i} m_{i} u_{i} = \sum_{i} m_{i} v_{i}$$

conservation of momentum

 $\Delta p = F\Delta t$

$$E_{\mathbf{k}} = \int p \, dv$$

Electricity

General

$$I = \frac{\Delta Q}{\Delta t} = \frac{\partial Q}{\partial t} = \dot{Q}$$

$$R = \frac{U}{I} = \rho \frac{l}{A}$$

 $\mathtt{in}\ A$

$$R = \frac{U}{I} = \rho \frac{l}{A}$$

in $\boldsymbol{\Omega}$

$$E = U \cdot Q$$

$$P = \frac{\Delta E}{\Delta t} = UI$$

in \boldsymbol{W} Energy flow /

"Power"

9 Fields

9.1 Newtonian gravitation

Homogeneous field

$$\begin{split} E &= mgh \\ \vec{F}_g &= m\vec{g} \\ \vec{g} &= \frac{\vec{F}_g}{m} \\ \Delta\varphi &= \frac{E}{m} = gh \end{split}$$

Radial symmetric field

$$\begin{split} U &= GMm \left(\frac{1}{r_1} - \frac{1}{r_2}\right) \\ U &= -GMm \frac{1}{r} & \text{for } r_1 \to \infty \\ \vec{F_g} &= -G\frac{Mm}{r^2} \hat{r} \\ \vec{g} &= \frac{\vec{F_g}}{m} = -G\frac{M}{r_2} \hat{r} \\ \Delta \varphi &= \frac{E}{m} = \gamma M \left(\frac{1}{r_1} - \frac{1}{r_2}\right) \end{split}$$

9.2 Electromagnetism

homogenous field

$$E = q\vec{\mathbf{E}}d$$

$$\vec{F} = q\vec{\mathbf{E}}$$

$$\vec{\mathbf{E}} = \frac{\vec{F}}{q}$$

$$\mathbf{E} = \frac{V}{d}$$

$$V = \frac{E}{q} = \vec{\mathbf{E}}d$$

Radial symetric field

$$\begin{split} U &= -\frac{1}{4\pi\varepsilon_0\varepsilon_r}Qq\left(\frac{1}{r_1} - \frac{1}{r_2}\right) \\ U &= \frac{Qq}{4\pi\varepsilon_0}\frac{1}{r} & \text{for } r_1 \to \infty \\ \vec{F} &= \frac{1}{4\pi\varepsilon_0\varepsilon_r}\frac{Qq}{r^2}\hat{r} \\ \mathbf{E} &= \frac{\vec{F}}{q} = \frac{1}{4\pi\varepsilon_0\varepsilon r}\frac{Q}{r^2}\hat{r} \\ \varepsilon_0 &= \frac{\sigma}{\varepsilon_r\mathbf{E}} & \text{const.} \end{split}$$

Index

В	F
Bohr radius1	Faraday constant1
Boltzmann constant1	•
E Elecromagnetism Electrostatics Radial symetric field	G Gravitation
Leptons	L
Muon1	Light1
Energy	
Kinetic Energy3	P
Potential Energy3	Plank constant1

Bibliography

1. <u>CODATA Internationally recommended 2018 values of the Fundamental Physical Constants</u>. May 2019. https://physics.nist.gov/cuu/Constants/index.html (2022).