

Comparison Between Monopolar and Bipolar Electrical Stimulation of the Motor Cortex

T. Kombos, O. Suess, B.-C. Kern, T. Funk, T. Hoell, O. Kopetsch, and M. Brock

Department of Neurosurgery, University Hospital Benjamin Franklin, Free University of Berlin, Berlin, Germany

Summary

Intra-operative neurophysiological techniques allow reliable identification of the sensorimotor region and make their anatomical and functional preservation feasible. Monopolar cortical stimulation has recently been described as a new mapping technique. In the present study this method was compared to the “traditional” technique of bipolar stimulation.

Functional mapping of the motor cortex was performed in 35 patients during surgery in the central region. The central sulcus (CS) was identified by somatosensory evoked potential (SEP) phase reversal. Cortical motor mapping was first performed by monopolar anodal stimulation with a train of 500 Hz (7–10 pulses) followed by bipolar stimulation (pulses at 60 Hz with max. 4 sec train duration). Surgery was performed under general anaesthesia without muscle relaxants. Of 280 motor responses elicited by bipolar cortical stimulation, 54.28% [152] were located in the primary motor cortex (PMC), 37.85% [106] outside the motor strip in the secondary motor cortex (SMC), and 8% [22] posterior to the CS. Of 175 motor responses elicited by monopolar cortical stimulation, 68.57% [120] were located in the SMC, 23.42% [41] in the SMC and 8% [14] posterior to the CS.

Contrary to the general clinical view, there is considerable overlapping of primary motor units over a cortical area much broader than the “classical” narrow motor strip along the CS. Bipolar cortical stimulation is more sensitive than monopolar for mapping motor function in the premotor frontal cortex. Both methods are equally sensitive for mapping the primary motor cortex.

Keywords: Cortical mapping; compound motor action potentials; motor pathways; intraoperative monitoring.

Introduction

The marked advances in anatomic orientation achieved during the past two decades thanks to the development of imaging procedures have paved the way for microsurgical techniques aiming at functional preservation during brain surgery. The more aggressive the approach, the better the survival and the quality of life [1, 12, 24, 30]. The aggressiveness, however, is often limited by the proximity of functionally

eloquent areas. Although MR imaging enables exact localization of lesions in relation to the central sulcus [5, 58], these morphological data do not correlate with function. Functional imaging techniques such as PET or fMRI, on the other hand, while allowing preoperative localization of eloquent areas, are only available in a few centres worldwide and cannot always be applied in the daily routine. Therefore intra-operative functional mapping and monitoring techniques are of paramount importance in localizing functionally relevant areas, and allowing maximal resection with minimal morbidity.

Monopolar motor cortex stimulation has recently been introduced as a new mapping technique [11, 53]. Cedzich *et al.* [11] showed that this method improves the outcome of surgery in and around the central region. It is, however, considered a “technique in evolution” [11]. In the present study the ability of monopolar cortical stimulation for functional cortical mapping is compared with that of the “traditional” technique of bipolar cortical stimulation [6, 15].

Patients and Methods

Patients

Functional mapping of the motor cortex was performed in 35 patients (15 males and 20 females aged 12 to 76, mean 51.91 years) during surgery in and around the central region. The space-occupying lesions were located in the frontal lobe anterior to the precentral gyrus in 11 patients, in the precentral gyrus in 15, and in the parietal lobe in 9 patients. There were 20 dominant-hemisphere (left) lesions, the remaining 15 tumours being in the nondominant (right) hemisphere. In all patients the topographic relationship between the lesion and the sensorimotor area was evaluated preoperatively by means of computed tomography or magnetic resonance imaging. Table 1 shows the histological diagnosis in this series of patients.

Table 1. *Histological Diagnosis in 35 Patients*

Histological diagnosis	Number of patients
Glioblastoma multiforme	11
Low grade glioma	5
Metastasis	9
Arteriovenous malformation	3
Epidermoid	1
Meningioma	6




Fig. 1. Distribution of motor areas identified by bipolar cortex stimulation

Methods

First the central sulcus (CS) was identified by means of phase reversal of somatosensory evoked potentials (SEP) [36]. The exposed cortex was then marked by small sterile numbered white paper labels. Cortical motor mapping was initially performed by monopolar anodal stimulation with a train of 500 Hz (7–10 pulses) (stimulation intensity 5–20 mA) followed by bipolar stimulation (1 msec pulses at 60 Hz) (stimulation intensity 9–20 mA). The stimulation parameters are summarized in table 2. Each motor point identified was marked by substituting the sterile white paper by a blue paper label. The motor areas identified were divided into three categories: 1) areas located in the primary motor cortex (PMC), which defined as the precentral gyrus and the frontal gyrus directly rostral to it; 2) areas located in the secondary motor cortex (SMC), defined as the frontal cortex adjacent to the PMC; and 3) areas in the parietal lobe (Figs. 1, 2).

All operations were performed under total intravenous anaesthesia (TIVA). Muscle relaxants were administered only for intubation, not during surgery. A Nicolet Viking IV (Nicolet Instruments, Biomedical Division, Madison, WI) was used for all examinations.




Fig. 2. Distribution of motor areas identified by monopolar cortex stimulation

Table 2. *Stimulation and Recording Methods*

	Bipolar stimulation	Monopolar stimulation
Intensity	9–20 mA	5–20 mA
Frequency	50–60 Hz	400–500 Hz
Duration	2–4 s	0.0175–0.02 s
Filter	10 Hz–10 kHz	10 Hz–10 kHz
Sensitivity	50–100 μ V/Div	20–100 μ V/Div
Recording	needle electrodes	needle electrodes

Method of SEP Phase Reversal

Bipolar cortical SEPs were recorded with a strip (row of five or six electrodes embedded in silicon) or grid electrode (2×5 ; 3×5) (Ad-Tech; Ad Technic, WI) placed on the cortex. The centres of the electrodes were 1.5 cm apart. The median and tibial nerves contralateral to the lesion were stimulated by 0.2 ms pulse with a repetition time of 2.3 per second and an intensity of 10–24 mA. The amplification band-pass was set at values of 100 Hz to 1 kHz. Between 30 and 100 responses were averaged.

Method of Monopolar Cortical Stimulation

Identification of the CS by SEP phase reversal was followed by functional mapping of the motor system. A short train (7–10 pulses) of monopolar, anodal, rectangular pulses was applied to the cortex at a frequency of 500 Hz. A single steel plate of 3.5 mm diameter embedded in silicon was used as anode. The cathode was located ipsilaterally on either Fp1 or Fp2 according to the 10–20 System. The intensity applied was 5 to 20 mA. The impulse duration ranged between 0.1 and 0.7 ms (Table 2). A constant voltage stimulator was used. Stimulation intensity was increased stepwise until action potentials were elicited in the target muscles. The maximum was set at 20 mA.

Action potentials were recorded from the forearm flexor, thenar and quadriceps muscles contralateral to the side of stimulation. A pair of subdermal needle electrodes was used for recording.

Method of Bipolar Cortical Stimulation

A bipolar electrode was applied parallel to the sulcus adjacent to the gyrus being stimulated. The stimulation probe was made out of 2 spherical steel tips with a diameter of 2.5 mm each. The tips were 5 mm apart. The same constant voltage stimulator was used to produce a train of low-frequency pulses. The pulse frequency was 50 to 60 Hz with a single impulse duration of 1 ms. The duration of the train was between 2 to 4 s. The current intensity varied between 9 and 20 mA. The maximum was set at 20 mA (Table 2). Increments of 1–2 mA were added to the baseline current until a motor response was evoked. This was either recorded by needle electrodes or observed as a movement of the contralateral face, hand or leg. Any time a response was recorded or a movement witnessed, the location of the stimulated site was noted.

Anaesthesia

All operations were performed under total intravenous anaesthesia (TIVA). Muscle relaxants were administered only for intubation, not during surgery. Anaesthesia was induced by Propofol (1–2 mg/kg) and Fentanyl (5–10 µg/kg). Propofol (75–125 µg/kg/h) was continuously applied during surgery. Analgesia was achieved by Alfentanil in 32 patients, Sulfentanyl in 1 and Fentanyl in another 2 patients.

Results

The location of the central sulcus was demonstrated by typical SEP phase reversal across the central sulcus in 33 cases. In 2 cases phase reversal was undetectable. Thus, it was possible to identify the central sulcus in 94.28% of the patients by phase reversal of the cortical SEP.

There were 175 responses following monopolar stimulation. Of these, 120 (69%) were located in the primary motor cortex (PMC) and 41 (23.42%) in the secondary motor cortex (SMC). Stimulation had to be intensified with increasing distance from the PMC. Fourteen (8%) responses were located in the parietal lobe dorsal to the central sulcus (Table 3a). In these cases the intensity of stimulation was the same as that applied to the PMC.

Six hundred and four monopolar stimulations were performed, 124 of which in the PMC, 280 in the SMC and 200 in the parietal lobe. Of the 124 stimulations in the PMC, 120 (96.77%) elicited an MEP. Following 280 stimulations in the SMC, however, MEPs were recorded only at 41 (14.64%) points. In the parietal lobe, MEPs were recorded only at 14 (7%) points (Table 3b) (Fig. 2).

Of the 280 motor responses following bipolar stim-

Table 3. *Results of Monopolar Cortex Stimulation. (a) Distribution of Monopolar Stimulations and the Motor Responses Elicited. (b) Distribution of Motor Responses Following Monopolar Stimulation*

(a) Motor responses			
PMC	120		68.57%
SMC	41		23.42%
Parietal	14		8%
Total	175		

(b) Monopolar stimulus Respond			
PMC	124	120	96.77%
SMC	280	41	14.64%
Parietal	200	14	7%
Total	604	175	

Table 4. *Results of Bipolar Cortex Stimulation. (a) Distribution of Bipolar Stimulations and the Motor Responses Elicited. (b) Distribution of Motor Responses Following Bipolar Stimulation*

(a) Motor responses			
PMC	152		54.28%
SMC	106		37.85%
Parietal	22		7.85%
Total	280		

(b) Bipolar stimulus Respond			
PMC	160	152	95%
SMC	392	106	27.04%
Parietal	244	22	9.01%
Total	796	280	

ulation, 152 (54.28%) were located in the PMC. There were 106 (37.85%) responses located in the SMC, while only 22 (7.85%) of the motor responses were elicited by stimulation posterior to the central sulcus (Table 4a). The intensity of bipolar stimulation was kept constant throughout the examination.

One hundred and forty three of the motor responses following bipolar stimulation were recorded by needle electrodes. Seventy on of these were recorded following stimulation of the PMC, 69 of the SMC and 3 of the parietal lobe. In the remaining 137 cases a complex movement was observed. No difference in the movement pattern was observed following stimulation of the PMC or the SMC.

There were 796 bipolar stimulations involving 160 points in the PMC, 392 points in the SMC and 244 points in the parietal lobe. In the PMC 152 (95%) points were identified as having motor functions, whereas 106 (27.04%) points in the SMC and 22 points

(9.01%) located dorsally to the central sulcus were identified as having motor functions (Table 4b) (Fig. 1).

Discussion

Electrical stimulation of the cerebral cortex was first performed by Bartholow in 1874 [3]. Cushing [13] used this technique to determine the anatomical relationship of the sensory strip to an adjacent tumour. However, it was the epoch-making study of Penfield and Boldrey [40] on evaluating motor and sensory representation in the human cerebral cortex by electrical cortex stimulation that laid the foundation for practical localization of the sensorimotor cortex in surgery. In recent years, several studies [6, 15–17, 26, 32, 33, 57] have substantiated the necessity for intra-operative functional mapping and monitoring during surgery in and around the motor cortex. The method of phase reversal of somatosensory evoked potentials is based on the fact that the dipole of the afferent volley changes from the postcentral to the precentral gyrus. A somatosensory potential (N20/P30) can thus be recorded from the postcentral gyrus and its mirror image (P'20/N'30) from the precentral gyrus [56]. In the present series, SEP phase reversal was recorded in 94.28%. This corresponds to the results of other series [11, 36]. Therefore somatosensory phase reversal is a reliable method for exact identification of the central sulcus. However, this technique yields no information about motor function. Anatomical identification alone is not a sufficient safeguard against postoperative neurological deficits. Therefore, an additional method is necessary to map and monitor motor function.

Motor function can be tested under general anaesthesia by direct motor cortex stimulation. Bipolar stimulation is the “traditional” method originally described by Fritsch and Hitzig [19]. This technique allows mapping of cortical and subcortical functional areas. Until recently this was the only method used for intra-operative mapping. A series of studies [6, 7, 15, 16, 37] have demonstrated that bipolar cortex stimulation is a useful tool during surgery in and around the motor cortex. Extensive experience with this technique constitutes its major advantage. However, bipolar cortical stimulation does not allow an objective analysis. Furthermore, the movements elicited by the stimulation cause interference during microneurosurgery, thus preventing continuous monitoring of motor function during tumour resection. This is the major

disadvantage of the method as compared to continuous monopolar stimulation. The rare induction of an intra-operative seizure by low-frequency stimulation is a further disadvantage of this technique.

The method of monopolar cortical stimulation as described recently [11, 53] requires an electrical stimulus of high-frequency. This method was applied in 35 patients. No intra-operative seizure occurred either in our series or in the series presented by Cedzich *et al.* [11]. The major advantage of this new technique is that it enables objective analysis of the results. The motor responses recorded can be analysed with regards to their latency, amplitude and duration. Since movements never occur following monopolar cortical stimulation, repetitive stimulation is possible during tumour removal, allowing continuous intra-operative monitoring of motor function [11]. Preliminary studies demonstrate a correlation between intra-operative changes of the compound muscle action potentials and clinical outcome [11, 28, 52].

The pia has a significant resistance and capacitance, which is destroyed by a few minutes exposure to air [4]. The resistance of gray matter is 4–6 times greater than cerebrospinal fluid [44] and 2–3 times less than resistance of white matter [2, 35, 46]. Different resistance affects the pattern of current flow according to Ohm's law. With extracellular stimulating electrodes for any outward current that locally depolarizes a fibre there is an inward current elsewhere that hyperpolarizes the fibre [45]. This has practical implications, one for a monopolar and a different one for a bipolar electrode.

Rudin and Eisenman [48] demonstrated that bipolar stimulation is more effective if the electrodes had a transverse rather than a longitudinal orientation to the axon [48]. According to Nathan *et al.* [34] during bipolar stimulation focus of current density (A/cm^2) can be achieved by spherical electrodes with 5 mm inter-electrode separation. Because of this in the present study a bipolar stimulation probe with 2 spherical steel electrodes with an inter-electrode distance of 5 mm was used. During bipolar stimulation with 10 mA intensity the peak current occurs in the region immediately beneath the bipolar electrodes ($0.05 A/cm^2$) [34].

With monopolar anodal stimulation current density decreases much less rapidly with depth [34]. Several studies have shown that monopolar stimulation of the cortex is more effective with an anode than a cathode [19, 42]. Anodal current enters and hyperpolarizes dendrites, and leaves and depolarizes the axon or cell body [43]. Hern *et al.* [23] have demonstrated that

monopolar anodal stimulation stimulates pyramidal cells directly. This occurs because the axon and the adjacent regions of the cell body have much lower threshold than dendrites and cell body [45].

Therefore, repetitive monopolar cortical stimulation induces repetitive excitation of the corticomotoneural tract [53]. Due to the high-frequency train, an accumulation of corticomotoneural postsynaptic potentials activating the motoneurons is achieved even under general anaesthesia [53]. Controversy exists regarding the compatibility between low-frequency bipolar stimulation and general anaesthesia. Several publications [5, 15, 40, 56] report on successful stimulation by using increased intensity under general anaesthesia.

Intra-operative mapping requires frequent stimulation throughout the procedure. The duration is 2 to 4 s for a bipolar stimulus as opposed to 1.4 ms for a monopolar stimulus. In addition to the difficulty of evaluating movement quantitatively, the longer duration of stimulus prolongs the monitoring procedure.

The presented results demonstrate that, contrary to the general clinical view, there is considerable overlapping of motor function units throughout a much wider area than the "classical" narrow cortical strip along the central sulcus. In humans and nonhuman primates at least 4 different motor areas are estimated: primary motor cortex; premotor cortex; supplementary motor cortex; cingulate gyrus [8, 9, 14, 21, 22, 25, 27, 29, 31, 38, 41, 49–51]. Therefore anatomical orientation must be complemented by intra-operative mapping of all these areas in order to prevent post-operative neurological deficits.

The common histological denominator of the motor cortex in human and nonhuman primate brains is the agranular cyto-architectonic pattern, characterized by the Betz cells in the primary motor cortex and the rostrally adjoining frontal cortex [10, 47]. The agranular region is bordered rostrally by the prefrontal cortex and caudally by the granular somatosensory cortex. The cingulate gyrus marks the border of the agranular isocortex [54, 55]. In humans parts of the cingulate cortex, beneath the supplementary motor area, are activated during various motor paradigms [39, 51].

This study compares cortical motor mapping by these two methods of stimulation. Both methods are equally sensitive for mapping the primary motor cortex. However, the bipolar stimulation is more sensitive than the monopolar for mapping motor function in the premotor frontal cortex. Since monopolar stimulation activates the corticomotoneural tract, this technique

achieves better results in areas with a high density of pyramidal-cells. Thus, localization is better in the PMC. Both techniques revealed motor function units parietalwards to the central sulcus. Galea *et al.* [20] could demonstrate, that parietal association areas, in the posterior parietal lobe, are in conjunction with the primary motor cortex [18]. The clinical relevance of these units was not the object of this study. This question should be elucidated by future studies.

Conclusion

Monopolar cortex stimulation is a relatively simple technique which is just as sensitive as bipolar stimulation for mapping the primary motor cortex. In the premotor frontal cortex, bipolar stimulation is more sensitive in localizing functional areas. For surgery on the primary motor cortex, monopolar stimulation is the method of choice. Surgery on the SMC requires a combination of bipolar stimulation for mapping and monopolar stimulation for monitoring the descending motor pathways.

Acknowledgments

This study was supported by a grant from the Berliner Krebsgesellschaft.

This article reports in extenso primary results reported to the 11th International Congress of Neurosurgery, Amsterdam 6th–11th July 1997.

References

1. Ammirati M, Vick N, Liao Y, Ciric I, Mickhael M (1987) Effect of the extent of surgical resection on survival and quality of life in patients with supratentorial glioblastomas and anaplastic astrocytomas. *Neurosurgery* 21: 201–206
2. Axer H, Stegelmeyer J, Graf von Kayserlingk D (1999) Comparison of tissue impedance measurements with nerve fiber architecture in human telencephalon: value in identification of intact subcortical structures. *J Neurosurg* 90: 902–909
3. Bartholow R (1874) Experimental investigations into functions of the human brain. *Am J Med Sci* 67: 305–313
4. Bennett MVL (1969) Electrical impedance of brain surfaces. *Brain Res* 15: 584–590
5. Berger MS, Cohen WA, Ojemann GA (1990) Correlation of motor cortex brain mapping data with magnetic resonance imaging. *J Neurosurg* 72: 383–387
6. Berger MS, Kincaid J, Ojemann GA, Lettich BA (1989) Brain mapping techniques to maximize resection, safety and seizure control in children with brain tumours. *Neurosurgery* 25: 786–792
7. Berger MS, Ojemann GA, Lettich E (1990) Neurophysiological monitoring during astrocytoma surgery. *Neurosurg Clin N Am* 1: 65–80
8. Boecker H, Khorram-Sefat D, Kleinschmidt A, Merbold KD,

- Hänicke W, Requardt M, Frahm J (1995) High-resolution functional magnetic resonance imaging of cortical activation during tactile exploration. *Human Brain Mapping* 3: 236–244
9. Boecker H, Kleinschmidt A, Requardt M, Hänicke W, Merbold KD, Frahm J (1994) Functional cooperativity of human cortical motor areas during self-paced finger movements: a high-resolution MRI study. *Brain* 117: 1231–1239
10. Brock M, De Alencar A (1961) Estudo sobre a origem e o trajeto dos axônios córticoespinhais no rato. *J Bras Neurol* 13: 217–232
11. Cedzich C, Taniguchi M, Schäfer S, Schramm J (1996) Somatosensory evoked potential phase reversal and direct motor cortex stimulation during surgery in and around the central region. *Neurosurgery* 38: 962–970
12. Ciric I, Ammirati M, Vick N, Mickhael M (1987) Supratentorial gliomas: surgical considerations and immediate postoperative results. Gross total resection versus partial resection. *Neurosurgery* 21: 21–26
13. Cushing H (1909) A note upon the Faradic stimulation of central gyrus in conscious patients. *Brain* 32: 42–53
14. Decety J, Perani D, Jeannerod M, Bettinardi V, Tadary B, Woods RP, Mazziotta JC, Fazio F (1994) Mapping motor representations with positron emission tomography. *Nature* 371: 600–602
15. Ebeling U, Schmid UD, Reulen HJ (1990) Tumour surgery within the central motor strip: surgical results with aid of electrical motor cortex stimulation. *Acta Neurochir (Wien)* 101: 101–107
16. Ebeling U, Schmid UD, Ying H, Reulen HJ (1992) Safe surgery of lesions near the motor cortex using intra-operative mapping techniques: A report on 50 patients. *Acta Neurochir (Wien)* 119: 23–28
17. Firsching R, Klug N, Borner U, Sanker R (1992) Lesions of the sensorimotor region: somatosensory evoked potentials and ultrasound guided surgery. *Acta Neurochir (Wien)* 118: 87–90
18. Freund HJ (1996) Remapping the brain. *Science* 272: 1754–1754
19. Fritsch G, Hitzig E (1870) Über die elektrische Erregbarkeit des Grosshirns. *Arch Anat Physiol Wiss Med* 37: 300–332
20. Galea MP, Darian-Smith I (1994) Multiple corticospinal neuron populations in the macaque monkey are specified by their unique cortical origins, spinal determinations, and connections. *Cereb Cortex* 4: 166–194
21. Grafton ST, Fagg AH, Woods RP, Arbib MA (1996) Functional anatomy of pointing and grasping in humans. *Cereb Cortex* 6: 226–237
22. Grafton ST, Woods RP, Tyszka M (1994) Functional imaging of procedural motor learning: relating cerebral blood flow with individual subject performance. *Human Brain Mapping* 1: 221–234
23. Hern EC, Landgren S, Phillips CG, Porter R (1962) Selective excitation of corticofugal neurones by surface-anodal stimulation of the baboon's motor cortex. *J Physiol (Lond)* 161: 73–90
24. Hirakawa K, Suzuki K, Ueda S, Nakawa Y, Yoshino E, Ibayashi N (1984) Multivariate analysis of factors affecting postoperative survival in malignant astrocytomas. *J Neurooncol* 12: 331–340
25. Jahanshahi M, Jenkins IH, Brown RG, Marsden CD, Passingham RE, Brooks DJ (1995) Self-initiated versus externally triggered movements. I. An investigation using measurement of regional blood flow with PET and movement related potentials in normal and Parkinson's disease subjects. *Brain* 118: 913–933
26. Katayama Y, Tsubokawa T, Maejima S, Hirayama T, Yamamoto T (1988) Corticospinal direct response in humans: Identification of the motor cortex during intracranial surgery under general anesthesia. *J Neurol Neurosurg Psychiatry* 51: 50–59
27. Kawashima R, O'Sullivan BT, Roland PE (1994) Fields in human motor areas involved in preparation for reaching, actual reaching and visuomotor learning: a positron emission tomography study. *J Neurosci* 14: 3462–3474
28. Kombos Th, Kern B-C, Höll T, Brock M (1997) Monopolar motor cortex stimulation: a new mapping technique? *Clin Neurol Neurosurg* 99: 108–108
29. Larsson J, Gulyas B, Roland PE (1996) Cortical representation of self-paced finger movement. *Neuroreport* 7: 463–468
30. Laws ER, Taylor WF, Clifton MP, Okazaki H (1984) Neurosurgical management of low grade astrocytoma of the cerebral hemispheres. *J Neurosurg* 61: 665–673
31. Leonardo M, Fieldman J, Sadato N, Campbell G, Ibanez V, Cohen L, Deiber MP, Jezzard P, Pons T, Turner R, Bihan D, Hallett M (1995) A functional magnetic resonance imaging study of cortical regions associated with motor task execution and motor ideation in humans. *Human Brain Mapping* 3: 83–92
32. Lüders H, Lesser RP, Hahn J (1983) Cortical somatosensory evoked potentials in response to hand stimulation. *J Neurosurgery* 58: 885–894
33. Naidich TP, Valavanis AG, Kubik S (1995) Anatomic relationships along the low-middle convexity: Part I – Normal specimens and magnetic resonance imaging. *Neurosurgery* 36: 517–532
34. Nathan SS, Sinha SR, Gordon B, Lesser RP, Thakor NV (1993) Determination of current density distributions generated by electrical stimulation of the human cerebral cortex. *Electroencephalogr Clin Neurophysiol* 86: 183–192
35. Nicholson PW (1965) Specific impedance of cerebral white matter. *Exp Neurol* 13: 386–401
36. Nuwer MR (1991) Localization of motor cortex with median nerve somatosensory evoked potentials. In: Schramm J, Möller A (eds) *Intraoperative neurophysiological monitoring*. Springer, Berlin Heidelberg New York Tokyo, pp 63–71
37. Ojemann GA, Ojemann J, Lettich E, Berger MS (1989) Cortical language localization in left, dominant hemisphere. An electrical stimulation mapping investigation in 117 patients. *J Neurosurg* 71: 316–326
38. Parsons LM, Fox PT, Downs H, Glass T, Hirsch TB, Martin CC, Jerabek PA, Lancaster JL (1995) Use of implicit motor imagery for visual shape discrimination as revealed by PET. *Nature* 375: 54–58
39. Paus T, Petrides M, Evans AC, Eyer E (1993) Role of human anterior cingulate cortex in the control of oculomotor, manual, and speech responses: a positron emission tomography study. *J Neurophysiol* 70: 453–469
40. Penfield W, Boldrey E (1937) Somatic motor and sensory representation in the cerebral cortex of man as studied by electrical stimulation. *Brain* 60: 389–443
41. Penfield W, Welch K (1951) The supplementary motor area of the cerebral cortex. *A.M.A. Arch Neurol Psychiatry* 66: 289–317
42. Phillips CG, Porter R (1962) Unifocal and bifocal stimulation of the motor cortex. *J Physiol (Lond)* 162: 532–538
43. Purpura DP, McMurty JG (1965) Intracellular activities and evoked potential changes during polarization of motor cortex. *J Neurophysiol* 28: 166–185
44. Ranck JB Jr. (1963) Specific impedance of rabbit cerebral cortex. *Exp Neurol* 7: 144–152
45. Ranck JB Jr. (1975) Which elements are excited in electrical stimulation of mammalian central nervous system: a review. *Brain Res* 98: 417–440
46. Ranck JB Jr., BeMent SL (1965) The specific impedance of the dorsal columns of cat: An anisotropic medium. *Exp Neurol* 11: 451–463

47. Roland PE, Zilles K (1996) Functions and structures of the motor cortices in humans. *Curr Opin Neurobiol* 6: 773–781
48. Rudin DO, Eisenman G (1954) The action potential of spinal axons in vitro. *J Gen Physiol* 37: 505–538
49. Schlaug G, Knorr U, Seitz RJ (1994) Inter-subject variability of cerebral activations in acquiring a motor skill: a study with positron emission tomography. *Exp Brain Res* 98: 523–534
50. Seitz RJ, Caravan AGM, Yagüez L, Herzog H, Tellmann L, Knorr U, Huang Y, Hömberg V (1994) Successive roles of the cerebellum and premotor cortices in trajectorial learning. *Neuroreport* 5: 2541–2544
51. Stephan KM, Fink GR, Passingham RE, Silbersweig D, Ceballos-Baumann AO, Frith CD, Frackowiak RSJ (1995) Functional anatomy of the mental representation of upper extremity movements in healthy subjects. *J Neurophysiol* 73: 373–386
52. Suess O (1999) Die Monopolare Kortex Stimulation (MKS): Eine neue Methode für intraoperatives neurophysiologisches Monitoring bei Operationen im, bzw. in der Nähe des Gyrus praecentralis. Inaugural Dissertation: Freie Universität Berlin, Berlin
53. Taniguchi M, Cedzich C, Schramm J (1993) Modification of cortical stimulation for motor evoked potentials under general anesthesia: Technical description. *Neurosurgery* 32: 219–226
54. Vogt BA (1993) Structural organisation of cingulate cortex: Areas, neurons, and somatodentritic transmitter receptors. In: Vogt BA, Gabriel M (eds) *Neurobiology of cingulate cortex and limbic thalamus*. Birkhäuser, Boston, pp 19–70
55. Vogt BA, Nimchinsky EA, Vogt LJ, Hof PR (1995) Humans cingulate cortex: surface features, flat maps, and cytoarchitecture. *J Comp Neurol* 359: 490–506
56. Wood C, Spencer D, Allison T, McCarthy G, Williamson P, Goff W (1988) Localisation of human sensorimotor cortex during surgery by cortical surface recording of somatosensory evoked potentials. *J Neurosurg* 68: 99–111
57. Woolsey CN, Erickson TC, Gilson WE (1979) Localization in somatic sensory and motor areas of human cerebral cortex as determined by direct recording of evoked potentials and electrical stimulation. *J Neurosurgery* 51: 476–506
58. Yousry TA, Schmid UD, Schmidt D, Hagen T, Jassoy A, Reiser MF (1996) The central sulcal vein: a landmark for identification of the central sulcus using functional magnetic imaging. *J Neurosurgery* 85: 608–617

Comments

Monopolar cortical stimulation has emerged as an elegant alternative to the more traditionally employed bipolar electrical stimulation. The major advantage is the possibility of obtaining an objective measure of the motor response in a continuous mode as opposed to eliciting a movement, thus allowing the performance of a microsurgical procedure without interruptions. The paper by Kombos *et al.* studies the issue of the sensitivity of monopolar stimulation, by comparing it with the gold standard of bipolar cortical stimulation. The comparison method, used in a series of 35 patients, allows the authors to reach substantial data and to deduce some formal conclusions.

Monopolar cortical stimulation is safe, since no single intra-operative epileptic fit is recorded in their series. The high-frequency stimulation required in monopolar mapping is therefore less invasive than the low-frequency used in bipolar stimulation. The monopolar technique seems highly sensitive to eliciting responses in the primary motor cortex (PMC), although the comparative results between both methods (96.77% versus 95%) may not have statistical value. In the secondary motor cortex (SMC), even when increasing the intensity of the electric stimulus, the responses were lower (14.64%) as compared with bipolar stimulation (27.04%).

The relative density of the Betz cells has a direct correlation with the possibility of eliciting electric motor responses in primates, as shown by Chang already in 1947 [1]. Uematsu *et al.* [3] presented a similar experience in humans using cortical grids. These grids in which the distance between the electrodes is constant can be placed over a sulcus, thus two contiguous neural banks can be excited by stimulation of electrode pairs. In fact 4.5% of the stimulations had a dual sensory and motor response. In the present study the authors demonstrate a more extensive motor area than exclusively the precentral sulcus, yet this is shown by either method.

G. Conesa – F. Isamat

This paper compares the effects of biopolar versus «monopolar» cortical stimulation as means of mapping the motor cortex during craniotomies in man.

The authors describe monopolar cortical stimulation as a relatively new development in the neurosurgical world. In fact, it has been around in physiological circles for many years now and its mode of action has been worked out in great detail by C Philips and R Porter (*Cortico-spinal Neurones: their role in movement*. Academic Press, London, 1997). The purists rebel against the definition monopolar because one obviously has to have two poles, otherwise the circuit is not complete. Phillips referred to it as «unifocal stimulation».

The results show clearly that both methods of stimulation can evoke motor potentials from the cortex, anterior to the motor cortex as well as posterior to it and, of course, within the motor cortex itself. They point out that bipolar stimulation, which was continuous for up to 4 seconds, was more effective at producing motor responses from the pre-motor cortex than was monopolar stimulation, where the stimulus only lasted up to twenty milliseconds.

On page 10 the authors express some surprise that motor responses can be obtained from outside the motor strip. This has been known ever since the motor strip was first described. I am not sure whether the correct explanation is that it is a phylogenetic spreading of pyramidal cells over the cortex as they say but, I think they should refer to the localisation of cortico-spinal tract neurones which has been thoroughly researched in primate. Their use of «pyramidal cells» to describe cortico-spinal tract neurones is potentially confusing (second paragraph, page 10).

F. Iannotti

Correspondence: Theodoros Kombos, M.D., Neurochirurgische Klinik und Poliklinik, Universitätsklinikum Benjamin Franklin, Freie Universität Berlin, Hindenburgdamm 30, D-12200 Berlin, Germany.