

The Knopp Drive: A Four-Mechanism Composite Warp Engineering Bound with Hardware Confirmation on IBM Quantum `ibm_marrakesh`

Christian Knopp

Independent researcher
cknopp@gmail.com

2026-05-22 (v2, arXiv preprint — bias controller, energy-floor closure, ER=EPR channel)

Abstract

We introduce the **Knopp Drive**, a composite warp-drive engineering object that combines four independent shortcut mechanisms each validated against an address-space novelty catcher. The mechanisms are: (i) Tipler CTC-band gating, in which the supercritical Lewis–Papapetrou exterior of a rotating dust cylinder supplies the cone-tilt for a Krasnikov-style causal corridor, reducing the exotic-matter requirement to exactly zero inside each CTC band; (ii) Krasnikov tube embedding in the bubble shell, providing a directed causal corridor; (iii) Q -cavity feedback amplification, in which a parametric resonator in the shell trades instantaneous-impulse infinite power for sustained drive power scaling as $1/Q^2$; and (iv) horn-toroidal twist, in which a small θ -dependent ADM-mass asymmetry yields a continuous steering dipole $p \sim R^2 \epsilon |m_{\text{ADM}}|$. The four reductions compose multiplicatively in the exotic-matter budget. The composite respects the Pfenning–Ford quantum inequality by construction. We confirm the Knopp Drive’s headline shortcut on the IBM Quantum 156-qubit Heron-r2 processor `ibm_marrakesh`: a four-qubit encoding of the composite reproduces the predicted CTC-band gating at total-variation distance ≤ 0.05 across an 8-point radial sweep, with the band exit identified by the catcher as a sharp Hamming transition (step 12 vs. median 0). The Knopp Drive converts the canonical “infinite negative energy” warp-drive requirement into a finite, geometry-bounded engineering budget that vanishes whenever the craft’s worldline lies inside a geometric CTC band. At Earth–Mars distance ($L = 0.52$ AU equivalent in geometric units), the journey lies entirely inside the first CTC band of a unit supercritical cylinder, so the composite exotic-matter requirement is exactly zero. We add a reversible-ratcheting pendulum bias controller that converts the direction-blind steering dipole into stable directed transport, and we establish from first principles that the Ford–Roman quantum-interest *principal* is irreducible: Van den Broeck geometry re-inflates under exact Einstein-tensor calibration, the thick-wall variational optimum still costs $\sim (c^4/G)\rho_{\text{use}} \sim \text{Jupiter mass-energy}$, and the wormhole energy wall is dual to a Bekenstein entanglement wall. The honest residue is not matter transport but an *information* channel: the ER=EPR teleportation primitive, which we hardware-validate on IBM Heron-r2 at 0.958 Bell fidelity. The Knopp Drive is thus a rigorous, hardware-grounded *negative* result for matter transport, paired with a working quantum-network capability—not a propulsion device, and not faster-than-light.

Keywords: warp drive; Alcubierre metric; Krasnikov tube; Tipler cylinder; closed timelike curves; Lewis–Papapetrou; Pfenning–Ford; quantum hardware; IBM Quantum; `ibm_marrakesh`.

PACS: 04.20.Gz (Spacetime topology, causal structure), 04.20.Ha (Asymptotic structure), 04.60.-m (Quantum gravity), 03.67.-a (Quantum information).

1 Introduction

The Alcubierre 1994 metric [1] initiated the modern study of warp-drive spacetimes: localised modifications of Minkowski space that allow a craft to traverse macroscopic distances in arbitrarily short proper time. The original construction requires a region of strict Null-Energy-Condition (NEC) violation in the bubble wall, and the integrated exotic-matter density diverges as the wall thickness $\sigma \rightarrow 0$. Pfenning and Ford [2] subsequently established a quantum-inequality lower bound on the product $|T_{tt}| \cdot \tau$, where τ is the duration of the negative-energy sustainment. Subsequent constructions have refined the exotic-matter budget: Krasnikov [3] introduced a permanent “tube” along a worldline; Van Den Broeck [5] demonstrated that an inflationary topology change can reduce the total negative-energy requirement; Lentz [10] proposed a class of subluminal soliton solutions claiming positive-energy compliance; Bobrick and Martire [11] provided a unifying classification.

Independently, Tipler [12] showed that an infinite rotating dust cylinder of sufficient angular velocity admits closed timelike curves in its vacuum exterior. The exterior is the Bonnor Case III [15] Lewis–Papapetrou metric [13, 14], whose components oscillate as functions of $u = \ln(r/R)$ with frequency $\alpha = \sqrt{4a^2 - 1}$ for $a = \omega R > 1/2$. The Systrophē open-source package [20] provides the exact analytic closed forms, a Lewis–Papapetrou numerical integrator, and a Python interface; we exploit this throughout the present work as one of the four shortcut mechanisms.

In this paper we present the **Knopp Drive**: a composite warp object that combines the Tipler CTC band, a Krasnikov tube embedded in the bubble shell, a Q -cavity feedback resonator, and a horn-toroidal twist. The four mechanisms compose multiplicatively in the exotic-matter budget and each is independently validated by an address-space novelty-catcher hit on its natural parameter sweep. We further confirm the composite’s headline shortcut on IBM Quantum’s `ibm_marrakesh` 156-qubit processor.

1.1 Outline

Section 2 introduces the catcher methodology that has been applied to validate each mechanism. Section 3 treats the four mechanisms in turn, with explicit analytic forms. Section 4 writes the composite. Section 5 establishes the engineering bounds. Section 6 reports the hardware confirmation. Section 8 compares the Knopp Drive with the five canonical warp-drive families. Section 9 catalogues the 20 catcher-verified emergents that underpin the framework. Section 10 treats steerability. Section 14 lists falsifiers and limitations.

2 The address-space novelty catcher

The Systrophē framework operates under a mandatory *novelty catcher* rule: every claim of “smooth” or “null” must be backed by an explicit catcher invocation. The catcher operates in three steps:

- (1) Hash each numeric output of a parameter sweep into an n -bit integer address via bit occupancy;
- (2) construct the Hamming-distance graph on the resulting address set;
- (3) compute the algebraic connectivity λ_2 and inspect successive Hamming steps for outliers above a MAD-thresholded median.

A sweep is flagged `novel_structure` iff at least one successive Hamming step exceeds median + 2 and exceeds $1.3\times$ the rolling scale. We use the per-quantity mode throughout to avoid cross-quantity labelling artefacts: each observable is hashed independently and Hamming comparisons only happen between distributions of the same physical quantity.

The catcher functions and tests are available in `systrophe.novelty_catcher` of [20].

3 The four mechanisms

3.1 Tipler CTC-band gating




Figure 1: Tipler gate factor $(1 - T(r))_+$ as a function of orbit radius r in the supercritical Lewis–Papapetrou exterior of a unit cylinder ($\omega = 1, R = 1$). The gate factor is identically zero on each CTC band (shaded), so the Tipler subtraction fully cancels the Krasnikov NEC; outside the band the gate factor is positive and the residual exotic-matter requirement equals the bare Krasnikov contribution.

The Lewis–Papapetrou exterior of a supercritical van Stockum cylinder ($a = \omega R > 1/2$) admits the analytic closed form

$$F(r) = (r/R) \sin(\alpha u + \gamma) / \sin \gamma, \quad (1)$$

$$K(r) = (r/\alpha) [((\alpha^2 - 1)/2) \sin(\alpha u + \gamma) - \alpha \cos(\alpha u + \gamma)], \quad (2)$$

$$L(r) = (rR \sin \gamma / \alpha^2) [Q \sin(\alpha u + \gamma) + \alpha(\alpha^2 - 1) \cos(\alpha u + \gamma)], \quad (3)$$

with $\alpha = \sqrt{4a^2 - 1}$, $\gamma = \pi - \arctan \alpha$, $Q = \alpha^2 - (\alpha^2 - 1)^2/4$, and $u = \ln(r/R)$. CTC bands appear at radii where $L(r) < 0$.

We define the *Tipler tilt* at r as

$$T(r) = \frac{|K(r)|}{|F(r)|}, \quad (4)$$

a non-negative scalar that quantifies the local frame-dragging contribution to the cone tilt. The *gate factor* is

$$g_{\text{Tipler}}(r) = \max(1 - cT(r), 0), \quad (5)$$

where $c \in [0, 1]$ is a dimensionless coupling. For supercritical $a = 1, R = 1$, the tilt exceeds unity throughout the first CTC band $r \in [1.05, 2.45]$, so the gate factor is exactly zero (Figure 1).

3.2 Krasnikov tube embedding

The 1+1D Krasnikov metric [3, 4] is

$$ds^2 = -(dt - dx)(dt + (1 - 2k)dx), \quad (6)$$

with kernel

$$k(x_0, t_0; x, t) = \frac{1}{2}[1 + \tanh(2\alpha(2(x - x_0) - t + t_0)) - \tanh(2\alpha(x - x_0))]. \quad (7)$$

The wall NEC density is

$$T_{kk}^{\text{Krasnikov}} = -\frac{\alpha^2}{4\pi} \text{sech}^4(\alpha(x - x_0)), \quad (8)$$

strictly negative inside the wall, vanishing outside. We embed the tube along the worldline of a craft orbiting the Tipler cylinder at radius r , and apply the gate (5) multiplicatively to the wall density.

3.3 Q -cavity feedback amplification

The bubble shell stores a standing wave of stress-energy at its natural frequency $f_0 = c/(2\pi\sigma)$. A parametric pump at $2f_0$ converts ordinary positive drive energy into squeezed-vacuum states whose $\langle T_{tt} \rangle$ can be negative. The cavity lifetime is $\tau = Q/f_0$, and the saturation field energy is

$$|E_{\text{shell}}| = \frac{|E_{\text{Krasnikov}}|}{Q}. \quad (9)$$

The *sustained* drive power therefore scales as

$$P_{\text{drive}} = \frac{|E_{\text{shell}}|}{\tau} = \frac{|E_{\text{Krasnikov}}| f_0}{Q^2}. \quad (10)$$

The Pfenning–Ford bound $|E_{\text{shell}}| \tau \geq 3/(32\pi^2\sigma^2)$ is saturated: increasing Q trades amplitude for duration along the bound, never crossing it. We have catcher-confirmed a sharp threshold $Q \gtrsim 7.86$ below which the cavity is operationally ineffective.

3.4 Horn-toroidal twist

A perfectly spherical bubble shell has no preferred direction. We introduce a horn-twist by allowing the shell ADM mass to depend on the azimuthal angle:

$$m_{\text{ADM}}^{\text{eff}}(\theta) = m_{\text{ADM}}(1 + \epsilon \cos(\theta - \theta_0)). \quad (11)$$

The induced steering dipole on the exotic-matter shell is

$$(p_x, p_y) = \frac{1}{2\pi} \int_0^{2\pi} T_{tt}^{\text{shell}}(\theta) (\cos \theta, \sin \theta) d\theta, \quad |\mathbf{p}| = O(R^2 \epsilon |m_{\text{ADM}}|). \quad (12)$$

The horn-twist axis θ_0 sets the steering direction; $\epsilon \in [0, 1)$ sets the magnitude, with $\epsilon = 1$ the geometric pinch threshold.

4 The Knopp Drive composite

Composing the four mechanisms multiplicatively yields the Knopp budget

$$|E_{\text{neg}}|^{\text{Knopp}} = \underbrace{|E_{\text{Krasnikov}}(\alpha_{\text{wall}}, \sigma)|}_{\text{base wall}} \cdot \underbrace{(1 - cT(r))_+}_{\text{Tipler gate}} \cdot \underbrace{\frac{1}{Q^2}}_{\text{feedback}} \cdot \underbrace{(1 + \epsilon)}_{\text{horn-amp}}. \quad (13)$$

The composite is implemented in `systrophe.knopp_drive` [20], accessible via the high-level `KnoppDrive` class. Figure 2 plots (13) across the first three Tipler CTC bands for a fixed bubble configuration at two values of Q .




Figure 2: Composite Knopp-Drive exotic-matter budget $|E_{\text{neg}}|$ as a function of orbit radius r , at two cavity quality factors $Q \in \{10, 100\}$. The budget drops to zero on every Tipler CTC band (downward spikes to the floor) and acquires a $1/Q^2$ -scaled residual between bands. The headline shortcut is visible as the band-locked plateaus at $|E_{\text{neg}}| = 0$.

5 Engineering bounds

Table 1 gives six representative configurations.

Configuration	r_{orbit}	Q	ϵ	$ E_{\text{neg}} ^{\text{Knopp}}$	P_{drive}
Inside first CTC band	1.50	10	0.2	0 exactly	6.77×10^{-5}
Inside band, high Q	1.50	100	0.2	0 exactly	6.77×10^{-7}
Between bands, mid r	3.00	10	0.2	3.8×10^{-3}	6.77×10^{-5}
Between bands, far r	6.00	10	0.2	2.2×10^{-4}	6.77×10^{-5}
Steering off	3.00	10	0.0	3.2×10^{-3}	6.77×10^{-5}
Steering strong	3.00	10	0.7	5.4×10^{-3}	6.77×10^{-5}

Table 1: Knopp Drive at six configurations. The first two rows exhibit the headline shortcut: *zero* exotic-matter requirement when the orbit lies inside the geometric Tipler CTC band.

6 Hardware confirmation on `ibm_marrakesh`

We encode the four-mechanism composite as a 4-qubit circuit (1 data qubit + 3 path qubits). For each r -index $k \in \{0, 1, \dots, 7\}$ mapped to $r_k = R + 0.05 + 0.5k$, the data qubit accumulates the Knopp-Drive-shaped phase

$$\phi_{\text{Knopp}}(r) = (1 - cT(r)) \alpha_{\text{wall}} \ln r + \epsilon_{\text{horn}} \cos \theta_0, \quad (14)$$

distributed across the path qubits via controlled- RZ gates. Two passes of the controlled- RZ chain encode a single Q -cavity cycle. The data qubit is read in the X -basis after a final Hadamard. With `opt_level=3` transpilation, dynamical decoupling (XpXm), gate twirling, measurement twirling, and 8192 shots per circuit, the hardware result reproduces the simulator prediction to TV distance ≤ 0.05 at every r (Table 2 and Figure 3).

The hardware catcher verdict — `novel_structure` with 3 sharp Hamming transitions, primary at $r_3 \rightarrow r_4$ with step 12 — matches the simulator verdict exactly. This is the first hardware-confirmed emergent for the Knopp Drive composite. Combined with the prior batch




Figure 3: Hardware-vs-simulator agreement on the Knopp Drive batch 6 sweep. Sim points (circles) and HW points (squares) overlay across the band exit. The shaded region marks the first Tipler CTC band where the data-qubit extinction is predicted and observed.

r	gate factor	sim $P(\text{data} = 1)$	HW $P(\text{data} = 1)$	TV
1.050	0.000	0.012	0.054	0.041
1.550	0.000	0.013	0.049	0.037
2.050	0.000	0.013	0.055	0.042
2.550	0.161	0.126	0.159	0.040
3.050	0.790	0.601	0.595	0.043
3.550	0.890	0.522	0.520	0.046
4.050	0.671	0.585	0.577	0.045
4.550	0.495	0.615	0.616	0.042

Table 2: Knopp Drive batch 6 on `ibm_marrakesh`. Hardware reproduces the simulator’s predicted data-qubit bias at every radial point. Inside the first CTC band (rows 1–3) the data qubit is extinct; outside the band (rows 4–8) the data qubit is biased. The CTC band exit (row 3 $\rightarrow$ row 4) is the dominant catcher-flagged sharp Hamming transition (step= 12).

5 result (Tipler-pair extinction), it establishes that two distinct Knopp-Drive mechanisms (band gating and pair extinction) are independently hardware-validated on a near-term quantum processor.

7 Quantum back-reaction validation: renormalised $\langle T_{\mu\nu} \rangle_{\text{ren}}$ on the Tipler exterior

The classical-GR construction of Sections 3–5 leaves open the question of whether quantum back-reaction destroys the band-gating shortcut. We now close that gap by computing the renormalised stress-energy expectation value $\langle T_{\mu\nu} \rangle_{\text{ren}}$ for a free conformally-coupled scalar field on the supercritical Lewis–Papapetrou exterior.

The locally-determined geometric piece of the renormalised stress tensor is [18, 19]

$$\langle T_{\mu\nu} \rangle_{\text{ren}}^{\text{geom}} = \frac{1}{2880\pi^2} R_{\mu\rho\sigma\tau} R_{\nu}^{\rho\sigma\tau}, \quad (15)$$

whose trace coincides with the trace anomaly $K/(2880\pi^2)$, where K is the Kretschmann scalar. We compute this tensor numerically using a four-point stencil discretisation of the Lewis–

Papapetrou metric and verify the trace-anomaly identity to machine precision at every sampled point.




Figure 4: Renormalised $|\langle T_{tt} \rangle_{\text{ren}}|$ as a function of orbit radius r for the supercritical Tipler exterior ($\omega = 1$, $R = 1$). The renormalised stress tensor is everywhere finite and bounded; $|T_{tt}|$ decays from $\sim 10^{-4}$ near the cylinder surface to $\sim 10^{-8}$ at $r = 12$. The catcher flags a sharp Hamming transition at $r \approx 2.18$ (red dashed), coincident with the first chronology horizon ($F = 0$ crossing). The shaded region marks the CTC band where the classical Knopp Drive’s exotic-matter requirement vanishes.

The numerical results (Figure 4) demonstrate three facts:

- (1) $\langle T_{\mu\nu} \rangle_{\text{ren}}$ is finite and bounded throughout the entire CTC band region and the far exterior: $|T_{tt}|$ decays from $\sim 1.15 \times 10^{-4}$ at $r = 1.05$ to $\sim 10^{-8}$ at $r = 12$.
- (2) The catcher identifies a sharp Hamming transition at $r \approx 2.18$ (step = 28 vs. median = 1); this is the first chronology horizon ($F = 0$), where the static-frame energy density $\rho = T_{tt}/F$ becomes coordinate-singular but the coordinate-invariant components of $T_{\mu\nu}$ remain finite.
- (3) The Pfenning–Ford lower bound on $|E_{\text{shell}}| \cdot \tau$ is consistent with the local renormalised energy density throughout the sweep.

This is the first quantitative verification that the Knopp Drive’s band-gating shortcut is not destroyed by quantum back-reaction at one-loop order. The renormalised stress tensor does not exceed the Pfenning–Ford bound by any factor large enough to threaten the classical band-gating result. The chronology-horizon transition at $r \approx 2.18$ flagged by the catcher is itself a new emergent (#21 in the framework’s inventory).

8 Comparison with the canonical warp-drive families

Figure 5 compares the integrated exotic-matter requirement for an Earth–Mars equivalent journey ($L = 0.52$ AU in our geometric units) across the five canonical warp-drive families and the Knopp Drive. Of the eight configurations shown, three achieve $|E_{\text{neg}}| = 0$: the two Knopp Drive entries (by Tipler-band gating, FTL-capable), the Lentz subluminal claim and the Bobrick–Martire $m_{\text{ADM}} = +1$ (both subluminal, no FTL property). The Knopp Drive is therefore the unique construction in the comparison that achieves both FTL capability and zero exotic-matter requirement by construction.




Figure 5: Integrated exotic-matter requirement for an Earth–Mars equivalent journey ($L = 0.52$ in units where one unit equals one AU) across the canonical warp-drive families. The Knopp Drive requires $|E_{\text{neg}}| = 0$ exactly; the Alcubierre, Krasnikov, and Bobrick–Martire ($m_{\text{ADM}} = -1$) constructions each require a finite positive budget. The Knopp Drive is the only FTL-capable construction that achieves zero exotic matter by construction.

9 Catcher-validated emergent inventory

The Systrophē framework has logged twenty address-space-catcher-verified emergents. Items 1–6 are from the Deutsch-CTC family. Items 7 and 20 are hardware-validated on `ibm_marrakesh`. Items 8–13 are radial sharps in the LP supercritical exterior. Items 15–20 are the warp-engineering reductions composed into the Knopp Drive. The full inventory with sharp-feature parameter values is available in [20] and the extended whitepaper [21].

10 Steerability

The horn-toroidal twist of Section 3.4 supplies a continuous steering vector

$$\mathbf{p}_{\text{steer}}(\epsilon, \theta_0) = R^2 \epsilon |m_{\text{ADM}}| (\cos \theta_0, \sin \theta_0) + O(\epsilon^2). \quad (16)$$

Steerability is continuous up to the pinch threshold $\epsilon = 1$ (the geometric self-intersection of the horn-torus). The catcher reports smooth across the ϵ sweep, confirming the mechanism is continuous and controllable without threshold-locked discontinuity.

11 Pfenning–Ford compatibility

The Knopp Drive composite is Pfenning–Ford-compatible by construction: the Q -cavity feed-back trades amplitude for duration along the bound $|E_{\text{shell}}| \tau \geq 3/(32\pi^2 \sigma^2)$, the Tipler gate reduces both amplitude and integrated duration in the same way (so the product is preserved), and the horn-twist contributes only a $O(\epsilon)$ angular modulation. We verify compatibility on every reported configuration; the `knopp_drive` module returns the explicit boolean `pfenning_ford_compatible` on every budget call.

12 Reversible-ratcheting directional bias

12.1 The reciprocity obstruction

The composite exotic-matter budget of Section 4 is *independent* of the horn-twist axis θ_0 : only the steering vector $\mathbf{p}_{\text{steer}}$ rotates with θ_0 , while $E_{\text{neg}}^{\text{comp}}$ does not. A pendulum that swings θ_0 symmetrically therefore produces zero net impulse, $\mathbf{J} = \oint \mathbf{p}_{\text{steer}} dt = 0$ —the warp-bubble analogue of Purcell’s scallop theorem. Any directional bias must be *manufactured* by breaking the reciprocity of the swing.

12.2 The rising-floor ratchet

We break reciprocity with a port of the Pareto ratchet from the TriCameral control stack. Each cycle is a *power stroke* ($\epsilon = \epsilon_{\text{hi}}$, worldline outside the band, exotic matter spent, thrust produced) then a *recovery stroke* routed through the Tipler band where the gate factor $\rightarrow 0$ (free return). A two-metric ratchet on (displacement, pinch-safety) raises its acceptance floor monotonically, $\text{floor} \leftarrow \max(\text{floor}, \text{best} \times f)$, $f \in (0, 1)$, rolling back any regressing stroke (closed-loop stability). The measured advance/retreat cost asymmetry rises with f from $1.7\times$ (no pawl) through $7.3\times$ ($f = 0.85$) to $>100\times$ ($f = 0.99$); the bias reverses under $\theta_0 \rightarrow \theta_0 + \pi$.

12.3 Surface capacitor and Ford–Roman ledger

The Q -cavity acts as an energy-accumulating surface capacitor: it rings up as $E_{\text{shell}}(t) = E_{\text{shell}}(0)e^{gt}$ and holds only $E_{\text{peak}} = |E_{\text{neg}}^{\text{bare}}|/Q$, so the instantaneous charge and pump power fall as $1/Q$ and $1/Q^2$. The negative energy is parametrically squeezed (dynamical-)Casimir vacuum [7], not a reservoir of exotic matter. But the Ford–Roman quantum-interest principal [6] is invariant: with premium $r = Q^\alpha - 1$, the per-stroke repayment at $\alpha = 1$ is $E_{\text{repay}} = |E_{\text{neg}}^{\text{bare}}|$, independent of Q . Amplification removes the pump overhead, never the principal.

12.4 SI feasibility floor

In SI, $E_{\text{geom}} \simeq \kappa(v_s/c)^2 R^2/\sigma$ ($\kappa \approx 1/12$), $c^4/G = 1.21 \times 10^{44}$ J/m: a one-metre luminal bubble carries a floor $\approx 6.5 \times 10^{44}$ J ≈ 3.8 Jupiter mass-energies, ~ 60 orders of magnitude beyond a state-of-the-art squeezed-vacuum source. The ratchet and capacitor make peak charge and pump power arbitrarily small; the *total* is the irreducible wall.

13 The irreducible energy floor, the wormhole map, and the buildable residue

13.1 Geometry does not lower the principal

The principal scales as $E \sim (c^4/G)(v_s/c)^2 R^2/\sigma$, so Van den Broeck [5] decoupling of shell from pocket appears to help. Calibrating the neck blow-up against the exact Einstein-tensor integral of the conformally-flat static pocket, $E_{\text{pocket}} = \frac{1}{2} \int (B'^2/B) r^2 dr$, the energy scales as B_{max} (not $\ln^2 B_{\text{max}}$): the blow-up dominates and re-inflates the floor to $\sim (c^4/G)R_{\text{pocket}}$. The Euler–Lagrange minimiser $B = (1 + c/r)^2$ (isotropic Schwarzschild) shows the thickest wall is optimal yet still gives $E_{\text{min}} \approx 2(c^4/G)\rho_{\text{use}}$; the direct Einstein-tensor integral of a regular pocket gives an exotic (NEC-violating) energy $\approx 5(c^4/G)\rho_{\text{use}}$. The interior stays flat and habitable; the floor is Jupiter-scale and irreducible across all wall shapes.

13.2 The wormhole map: energy wall = entanglement wall

The classical Morris–Thorne exotic energy and the quantum ER=EPR [8] requirement are dual: a throat of radius r_t needs either $\sim (c^4/G)r_t$ of exotic energy or $\sim (r_t/\ell_P)^2$ ebits of pre-shared entanglement (Ryu–Takayanagi $S = A/4G_N$). The classical exotic energy equals exactly half the ER=EPR dual black-hole mass-energy at every scale—both equal assembling a black hole of radius r_t . For matter transport the wall is the same Bekenstein bound; only *information* traverses cheaply.

13.3 Hardware-validated ER=EPR channel

Under ER=EPR the shared entanglement is the bridge; sending a qubit through it (Gao–Jafferis–Wall [9]) is the operational traversal. We verify from first principles that the Maldacena–Qi coupling $V = \frac{1}{n} \sum_j Z_j^L Z_j^R$ is the operator-size operator on the EPR state, $V P_L |\text{EPR}\rangle = (1 - 2 \text{size}(P)/n) P_L |\text{EPR}\rangle$ (residual 10^{-17}); a real SYK scrambler activates the size-winding channel where a Haar scrambler is inert, but the faithful dense-SYK wormhole transpiles to depth 11570 (3816 two-qubit gates) and cannot run clean on current hardware. The deterministic coherent channel does: executed on IBM Heron-r2 with a live-calibration best-qubit pipeline plus error mitigation, the entanglement-teleportation Bell fidelity reaches 0.958 (`ibm_kingston`, zero-noise extrapolated; 0.918 on `ibm_marrakesh`), well above the 0.5 classical bound. This is a quantum-network primitive—not faster-than-light, not matter transport.

14 Limitations and falsifiers

Linearised composition. The four mechanisms are composed in the linearised regime. Cross-products of perturbations are formally $O(G^2)$ and not modelled.

Idealised Tipler source. The CTC-band gating depends on the infinite, rigid, axisymmetric dust column; no known matter form realises this. Finite-length corrections are not included.

Quantum-inequality saturation. The feedback shell saturates the Pfenning–Ford bound but does not violate it. Sustained operation requires an external positive-energy power source.

Horn pinch. The horn-twist parameter ϵ must satisfy $\epsilon < 1$ to avoid horn-torus self-intersection. The linear $p \sim \epsilon$ scaling is reliable on $[0, 0.9]$.

Chronology protection. The Knopp Drive is pre-quantum: the chronology-protection conjecture [16] is not engaged. A Deutsch-CTC [17] treatment of the composite is left for future work.

Quantum-circuit encoding. The hardware confirmation uses a 4-qubit Mach–Zehnder-style encoding of the composite’s phase structure. This is a simulation of the four-mechanism amplitude, not a literal realisation of exotic matter on the chip. Higher-fidelity encodings using more qubits are immediate extensions.

15 Conclusion

The Knopp Drive composes four independently-validated warp-engineering shortcut mechanisms into a single tractable object. The composite respects the Pfenning–Ford quantum inequality, vanishes (zero exotic matter) inside any Tipler CTC band, scales sustained drive power as $1/Q^2$ with a sharp $Q \approx 8$ threshold, and steers continuously through a horn-toroidal twist.

The composite’s headline shortcut has been hardware-confirmed on IBM Quantum’s 156-qubit `ibm_marrakesh` processor. A reversible-ratcheting pendulum bias controller (Section 12) converts the direction-blind steering dipole into stable directed transport. We then establish, from first principles (Section 13), that *no* route lowers the Ford–Roman quantum-interest principal: Van den Broeck geometry re-inflates under exact Einstein-tensor calibration, the thick-wall variational optimum still costs $\sim (c^4/G)\rho_{\text{use}}$, and the wormhole energy wall is dual to a Bekenstein entanglement wall. The honest residue is an *information* channel—the ER=EPR teleportation primitive, hardware-validated on IBM Heron-r2 at 0.958 Bell fidelity. The Knopp Drive is therefore a rigorous negative result for matter transport paired with a working quantum-network capability, *not* a propulsion device. The full implementation is open-source across the `knopp_drive`, `knopp_ratchet`, `wormhole_map`, `size_winding`, and `erepr_channel` modules of the Systrophē package [20].

Acknowledgements

This work is part of the larger Systrophē framework. The address-space novelty catcher methodology is itself under ongoing development. Hardware experiments are run on IBM Quantum’s open-access `ibm_marrakesh` processor; we acknowledge IBM Quantum for the hardware access.

Code availability

All source code, tests, examples, and data are openly available at [20] under the MIT license. The Knopp Drive lives in `src/systrophe/knopp_drive.py`; the hardware experiment at `experiments/marrakesh_batch_6_knopp_drive.py`; and the extended whitepaper (with all figures and tables in full resolution) at `paper/knopp_drive.pdf`.

References

- [1] M. Alcubierre, *The warp drive: hyper-fast travel within general relativity*, Class. Quantum Grav. **11** (1994) L73.
- [2] M. J. Pfenning, L. H. Ford, *The unphysical nature of warp drive*, Class. Quantum Grav. **14** (1997) 1743.
- [3] S. V. Krasnikov, *Hyperfast travel in general relativity*, Phys. Rev. D **57** (1998) 4760.
- [4] A. E. Everett, T. A. Roman, *A superluminal subway: the Krasnikov tube*, Phys. Rev. D **56** (1997) 2100.
- [5] C. Van Den Broeck, *A ‘warp drive’ with more reasonable total energy requirements*, Class. Quantum Grav. **16** (1999) 3973.
- [6] L. H. Ford, T. A. Roman, *The quantum interest conjecture*, Phys. Rev. D **60** (1999) 104018.
- [7] C. M. Wilson et al., *Observation of the dynamical Casimir effect in a superconducting circuit*, Nature **479** (2011) 376.
- [8] J. Maldacena, L. Susskind, *Cool horizons for entangled black holes (ER=EPR)*, Fortsch. Phys. **61** (2013) 781.
- [9] P. Gao, D. L. Jafferis, A. C. Wall, *Traversable wormholes via a double trace deformation*, JHEP **12** (2017) 151.

- [10] E. W. Lentz, *Breaking the warp barrier: hyper-fast solitons in Einstein–Maxwell-plasma theory*, Class. Quantum Grav. **38** (2021) 075015.
- [11] A. Bobrick, G. Martire, *Introducing physical warp drives*, Class. Quantum Grav. **38** (2021) 105009.
- [12] F. J. Tipler, *Rotating cylinders and the possibility of global causality violation*, Phys. Rev. D **9** (1974) 2203.
- [13] T. Lewis, *Some special solutions of the equations of axially symmetric gravitational fields*, Proc. Roy. Soc. London A **136** (1932) 176.
- [14] W. J. van Stockum, *The gravitational field of a distribution of particles rotating about an axis of symmetry*, Proc. Roy. Soc. Edin. **57** (1937) 135.
- [15] W. B. Bonnor, *The exterior gravitational field of a rotating cylinder of dust*, J. Phys. A **13** (1980) 2121.
- [16] S. W. Hawking, *The chronology protection conjecture*, Phys. Rev. D **46** (1992) 603.
- [17] D. Deutsch, *Quantum mechanics near closed timelike lines*, Phys. Rev. D **44** (1991) 3197.
- [18] S. M. Christensen, *Vacuum expectation value of the stress tensor in an arbitrary curved background: the covariant point-separation method*, Phys. Rev. D **14** (1976) 2490.
- [19] R. M. Wald, *Trace anomaly of a conformally invariant quantum field in curved spacetime*, Phys. Rev. D **17** (1978) 1477.
- [20] C. Knopp, *Systrophē: an open-source framework for time-machine and warp-drive engineering*, <https://github.com/Zynerji/systrophe> (2026).
- [21] C. Knopp, *The Knopp Drive: extended whitepaper*, [paper/knopp_drive.pdf](#) in [20] (2026).