

## LGS GROUP OF COLLEGES A PROJECT OF LAHORE GRAMMAR SCHOOL

Sheet #\_

ame: Kida Fatima	Class: 1-4ea	Roll No	
bject: Statistic	Test No. <u><b>WT-5</b></u>		
<b>8860 88</b>		0000	Marks Obtained
$     \begin{array}{ccccccccccccccccccccccccccccccccc$			
	II-MOIT)	10	
	SECTION-1		
	12 18.13.63.13.0.133.20	2 = 11 2 2002 5/0	Q3. 8 SALL
BANGE LEWINGS OF	( s) (i) (s s) (s s) (i)	00,000,0	W
SET:	1.70X 23Y 27 23 44 27 65	202120421	2.4
A set	is a well defined c	allection of	distinct
- 1. 1 1. 1 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1.	objects making up a		
H set is usua	ally denoted by a co	apital letter	ie, AB, Cetc.
(A. N.H. 1997)	-, 3,4,5} is a set.		
e.g A = \$1,2	-, 3,4,5} is a set.		eits
유민들 아름답다 중심하는 얼마나 없는 것이 되었다.		$\begin{array}{cccccccccccccccccccccccccccccccccccc$	
(A. N.H. 1997)	-, 3,4,5} is a set.	(1) 2 (1) 2 (1) (1) (1) (1) (1) (1) (1) (1)	
2 000/100PFH )=(3,1)		· A real consumer control of	A)9-
2,000/100/FP )=(31)	\$ 0 (ii) 15 =	(4)	A)9- 1 (jii)
2 000/100PFH )=(3.1)	\$ 0 (ii) 15 =	(12) (12) (2) (3)	A)9- 1 (jii)
(1,8) (3,5) (3,1) (1,8)=(479001600 (3,1) (3,6) (8,1) (8,1)	\$ 0 (ii) 15 =	(12) (12) (2) (3)	A)9
(1,8) (3,5) (3,1) (1,8)=(479001600 (3,1) (3,6) (8,1) (8,1)	\$ 0 (ii) 15 =	(12) (12) (2) (3)	4.9
(1.8) (3,1) (3,1) (1.8) (3,1)	\$ 0 (ii) 15 =	(1) y = (1)	4.9
1,8) (3,5) (3,1) (1,8)=(479001600 5 (1,8)=(319000 5 (1,8)=(319000 5 (1,8)=(319000 5 (1,8)=(319000 5 (1,8)=(319000 5 (1,8)=(319000 5 (1,8)=(319000 5 (1,8)=(319000 5 (1,8)=(319000 5 (1,8)=(319000 5 (1,8)=(319000 5 (1,8)=(319000 5 (1,8)=(319000	2 6 (ii) 3 6 (iii)	(P) = (A) $(D) = (A)$ $(D) = (A)$ $(D) = (A)$ $(D) = (A)$	4.9
18) (S.S.)  (18)=(479001600)  3628800)  = 132 Ans	20 (E12 (1:1) (2:2) 2 (1:1) (E12 (1:1) (1:2) 2 (1:1) (E12 (1:1) (1:2) 3 (1:1) (E12 (1:1) (1:2) 3 (1:1) (E12 (1:1) (1:2) 4 (1:1) (E12 (1:1) (1:2) 4 (1:1) (E12 (1:1) (1:2) 4 (1:1) (E12 (1:1) (1:2) (1:2) (1:2) 4 (1:1) (E12 (1:1) (1:2) (1:2) (1:2) 4 (1:1) (E12 (1:1) (1:2) (1:2) (1:2) (1:2) 4 (1:1) (E12 (1:1) (1:2) (1:2) (1:2) (1:2) (1:2) 4 (1:1) (E12 (1:1) (1:2)	(12) (12) (13) (13) (13) (13) (13) (13) (13) (13	
18) (S.S.)  (18)=(479001600)  3628800)  = 132 Ans  Sample space  A=A+ le	2 0 (E) 2 (1.10) = 2 2 (1.10) (E) 2 (1.10) = 2	(12) (12) (13) (13) (13) (13) (13) (13) (13) (13	
18) (S.S.)  479001600  3628800) (S.S.)  132 Ans  Sample space  A=A+ le  P(A)= v	2 0 (E) 2 (1.10) = 2 2 (1.10) (E) 2 (1.10) = 2	(12) (12) (13) (13) (13) (13) (13) (13) (13) (13	
18) (S.S.)  479001600  3628800) (S.S.)  132 Ans  Sample space  A=A+ le  P(A)= v	2 12 (21) (25) (2.12) (	(12) (12) (13) (13) (13) (13) (13) (13) (13) (13	
18) (S.S.)  479001600  3628800) (S.S.)  132 Ans  Sample space  A=A+ le  P(A)= v	Cins (SHS) (SES)  Constant one head, A=	(12) (12) (13) (13) (13) (13) (13) (13) (13) (13	
18) (S.S.)  479001600  3628800) (S.S.)  132 Ans  Sample space  A=A+ le  P(A)= v	20 (20 0.0) =  20 (20 0.0) =  20 (20 0.0) =  20 (20 0.0) =  20 (20 0.0) =  20 (20 0.0) =  20 (20 0.0) =  20 (20 0.0) =  21 (20 0.0) =  22 (20 0.0) =  23 (20 0.0) =  24 (20 0.0) =  25 (20 0.0) =  26 (20 0.0) =  27 (20 0.0) =  27 (20 0.0) =  28 (20 0.0) =  28 (20 0.0) =  29 (20 0.0) =  20 (20 0.0) =  20 (20 0.0) =  20 (20 0.0) =  20 (20 0.0) =  21 (20 0.0) =  22 (20 0.0) =  23 (20 0.0) =  24 (20 0.0) =  26 (20 0.0) =  27 (20 0.0) =  27 (20 0.0) =  28 (20 0.0) =	(12) (12) (13) (13) (13) (13) (13) (13) (13) (13	



ciis

## PROJECT OF AN AGRECORA VOLVERS

Rida Fatima

Sample space = S= {1, 2, 3	1.	5.61

Livi

A = even number = { 2, 4, 6}

P(A) = N(A) =, 3 =, 1 =, 0.5=P(A)

## SECTION-II

A = sum is 6 = { (1,5), (2,4), (3,3), (4,2), (5,1)}

P(A) = n(A) = 5 n(s) 36

A = sum is 4 or less = { <1,1)(1,2), <1,3), (2,1), <2,2), (3,1)}

 $\frac{P(A) = v(A)}{v(1)} = \frac{6}{36}$ 

(3,3) (3,5), (4,2), (4,4), (4,6), (5,1), (5,3), (5,5), (6,2), (6,4), (6,6)}

 $\frac{P(A) = \gamma(A)}{\gamma(1)} = \frac{18}{31}$ 

D(B) = 1

Scanned with

CS CamScanner