



## Operations Research

Publication details, including instructions for authors and subscription information:

<http://pubsonline.informs.org>

### Technical Note—A Note on Zero-One Programming

Manfred W. Padberg,

To cite this article:

Manfred W. Padberg, (1975) Technical Note—A Note on Zero-One Programming. Operations Research 23(4):833-837. <http://dx.doi.org/10.1287/opre.23.4.833>

Full terms and conditions of use: <http://pubsonline.informs.org/page/terms-and-conditions>

This article may be used only for the purposes of research, teaching, and/or private study. Commercial use or systematic downloading (by robots or other automatic processes) is prohibited without explicit Publisher approval. For more information, contact [permissions@informs.org](mailto:permissions@informs.org).

The Publisher does not warrant or guarantee the article's accuracy, completeness, merchantability, fitness for a particular purpose, or non-infringement. Descriptions of, or references to, products or publications, or inclusion of an advertisement in this article, neither constitutes nor implies a guarantee, endorsement, or support of claims made of that product, publication, or service.

© 1975 INFORMS

Please scroll down for article—it is on subsequent pages



INFORMS is the largest professional society in the world for professionals in the fields of operations research, management science, and analytics.

For more information on INFORMS, its publications, membership, or meetings visit

<http://www.informs.org>

## REFERENCES

1. J. W. COHEN, *The Single Server Queue*, American Elsevier, New York, 1969.
2. D. R. COX, *Renewal Theory*, Wiley, New York, 1962.
3. D. R. COX AND W. L. SMITH, *Queues*, Wiley, New York, 1961.
4. W. FELLER, *An Introduction to Probability Theory and its Applications*, Vol. II, Wiley, New York, 1966.
5. J. KEILSON, "The General Bulk Queue as a Hilbert Problem," *J. Roy. Statist. Soc. Ser. B*, **24**, 344-358 (1962).
6. L. KOSTEN, *Stochastic Theory of Service Systems*, Pergamon Press, Oxford, 1973.
7. P. LE GALL, *Les Systèmes Avec ou Sans Attente*, Dunod, Paris, 1962.

## A Note on Zero-One Programming

MANFRED W. PADBERG

*New York University, New York, New York*

(Received original May 15, 1973; final, March 3, 1975)

We generalize a method for constructing facets for the convex hull of integer solutions to set packing problems to arbitrary zero-one problems having nonnegative constraint-matrices. A particular class of facets is obtained explicitly and illustrated by a numerical example.

WE CONSIDER the convex polytope in  $R^n$  defined by

$$P = \{x \in R^n \mid \sum_{j=1}^{j=n} a_j x_j \leq a_0, 0 \leq x_j \leq 1, j \in N = \{1, \dots, n\}\}, \quad (1)$$

where  $a_j \in R_+^m$ ,  $j=0, 1, \dots, n$ , are nonnegative vectors with  $m$  integral components. Denote by  $P_I$  the convex hull of the integer vertices of  $P$ , i.e.

$$P_I = \text{conv}\{x \in P \mid x_j = 0 \text{ or } 1, \forall j \in N\}. \quad (2)$$

In this note we describe a general method that can be used to obtain 'facets' of  $P_I$ . An inequality  $\pi x \leq \pi_0$  is called a *facet* of  $P_I$  if (i)  $x \in P_I$  implies  $\pi x \leq \pi_0$  and (ii) there exist exactly  $d$  affinely independent vertices  $x^i$  of  $P_I$  satisfying  $\pi x^i = \pi_0$ ,  $i=1, \dots, d$ , if  $\dim P_I = d$ . Any inequality  $\pi x \leq \pi_0$  for which (i) holds is called a *valid inequality* for  $P_I$ . We note that 'cutting planes' used in integer programming generally constitute *valid* inequalities,<sup>[1]</sup> whereas *facets* of  $P_I$  constitute 'deepest' cutting planes. Facets are not only deepest in the sense that they cannot be pushed further into feasible set  $P$  without cutting off at least one feasible integer point of  $P$ ,

but they also belong to the class of inequalities that uniquely determines the polytope  $P_I$ . Whenever the dimension of  $P_I$  coincides with the number of variables in (1), i.e., if  $\dim P_I = n$ , then there is a uniquely defined class of facets of  $P_I$ .

We observe first that the inequalities  $x_j \geq 0$  are facets of  $P_I$ , provided that  $\mathbf{a}_j \leq \mathbf{a}_0$  for all  $j = 1, \dots, n$ . We shall call the inequalities  $x_j \geq 0$ ,  $j = 1, \dots, n$ , 'trivial' facets of  $P_I$ . Observe that for any nontrivial facet of  $P_I$  we have  $\pi_j \geq 0$ ,  $j = 1, \dots, n$  and  $\pi_0 > 0$ . Consequently, requirement (ii) in the definition of a facet states that there must exist  $d$  linearly independent vertices of  $P_I$  satisfying the condition in (ii). We now assume explicitly that  $\mathbf{a}_j \leq \mathbf{a}_0$  for all  $j \in N$ . Hence,  $\dim P_I = n$ .

Let  $T$  be a nonempty proper subset of  $N$ . Denote by  $P^T$  the polytope obtained from  $P$  by setting the variables  $x_j$ ,  $j \in T$ , equal to zero, i.e.,

$$P^T = P \cap \bigcap_{j \in T} \{x \in R^n \mid x_j = 0\} \quad (3)$$

and define  $P_I^T$  to be the convex hull of the zero-one points of  $P^T$ .

Let  $T = \{j_1, \dots, j_t\}$ , where  $t = |T|$  and the elements of  $T$  are arbitrarily ordered. For  $q = 1, \dots, t$  define  $T_q$  to be

$$T_q = T_{q-1} \cup \{j_q\}, \quad (4)$$

with the convention that  $T_0 = \emptyset$ . Similar to  $P^T$  and  $P_I^T$ , we denote by  $P^{T-T_q}$  the polytope obtained from  $P$  by setting the variables  $x_j$ ,  $j \in T - T_q$ , equal to zero and define  $P_I^{T-T_q}$  to be the convex hull of the zero-one points of  $P^{T-T_q}$ . Note that with the above definitions  $P^{T-T_0} = P^T$  and  $P^{T-T_t} = P$ . Furthermore, by the above assumptions we have that  $\dim P^{T-T_q} = \dim P_I^{T-T_q} = (n - t + q)$ .

Let  $\pi x \leq \pi_0$  be any valid inequality for  $P_I$  that is a (nontrivial) facet for the  $(n - t)$ -dimensional polytope  $P_I^T$  and consider the zero-one problem

$$z = \max \pi x, \quad x \in P_I^{T-T_1} \cap \{x \in R^n \mid x_{j_1} = 1\}, \quad (5)$$

where we have set the variables  $x_j$ ,  $j \in T - T_1$ , equal to zero and the variable  $x_{j_1}$  equal to +1. (Remember  $T_1 = \{j_1\}$ .)

Define the vector  $\pi^1$  as follows:  $\pi_j^1 = \pi_j$ ,  $\forall j \in N - T$ ,  $\pi_{j_1}^1 = \pi_0 - \bar{z}$ ,  $\pi_j^1 = 0$  otherwise, where  $\bar{z}$  is the optimal objective function value of (5). It follows easily that  $\pi_{j_1}^1 \geq 0$  and that the inequality  $\pi^1 x \leq \pi_0$  is a valid inequality for  $P_I$ , which is a (nontrivial) facet for the  $(n - t + 1)$ -dimensional polytope  $P_I^{T-T_1}$ . Continuing the above process with  $j_2$ , etc., until  $T$  is exhausted, we obtain a (nontrivial) facet for the  $n$ -dimensional polytope  $P_I$ . To be more specific, suppose that  $\pi_j$ ,  $j \in N - T$ , and  $\pi_0 > 0$  are given. Define a sequence of maximization problems  $(H_q)$  as

$$\begin{aligned} z_q = \max \quad & \sum_{j \in N - T} \pi_j x_j + \sum_{j \in T_{q-1}} \pi_j x_j \\ & \sum_{j \in N - T} \mathbf{a}_j x_j + \sum_{j \in T_{q-1}} \mathbf{a}_j x_j \leq \mathbf{a}_0 - \mathbf{a}_{j_q} \\ & x_j = 0 \text{ or } 1, \quad \forall j \in (N - T) \cup T_{q-1}, \end{aligned} \quad (H_q)$$

where the  $\pi_j, j \in T_{q-1}$ , are defined recursively by

$$\pi_{j_q} = \pi_0 - z_q \quad (6)$$

and  $z_q$  is the optimal value of the objective function of problem  $(H_q)$ ,  $q = 1, \dots, t$ . The following theorem generalizes Theorem 3.3 of reference 3. (See also reference 4, Theorem 3.11.)

**THEOREM.** Let  $T = \{j_1, \dots, j_t\}$ , where  $1 \leq t = |T| \leq n-1$ , be an arbitrarily ordered subset of  $N$  and let  $\pi x \leq \pi_0$  be a nontrivial facet of  $P_I^T$  as defined in (3). Let  $\pi'$  be defined by  $\pi'_j = \pi_j$  for all  $j \in N - T$ ,  $\pi'_{j_q} = \pi_0 - z_q$  for  $q = 1, \dots, t$ , where  $z_q$  are obtained by solving the problems  $(H_q)$  for  $q = 1, \dots, t$ . Then  $\pi' x \leq \pi_0$  is a (nontrivial) facet of  $P_I$ .

The proof of the theorem closely follows the argument used in the proof of Theorem 3.3 in reference 3. It also follows from Theorem 4.2 of reference 13 since if we take  $S = N$  and  $\mathcal{g} = \{I | I \subseteq N \text{ and } \sum_{j \in I} a_j \leq a_0\}$ , the pair  $(S, \mathcal{g})$  is easily verified to define an *independence system*, i.e.,  $I_1 \subseteq I_2 \in \mathcal{g}$  implies  $I_1 \in \mathcal{g}$ .

In the following we show how the above result can be used to construct a special class of facets of  $P_I$ . Let  $N'$  be any subset of  $N$  satisfying

$$\sum_{j \in N'} a_j \not\leq a_0, \quad (7)$$

where the symbol ' $\not\leq$ ' is meant to say that for *at least one component* of the vectors involved in (7) the inequality  $\leq$  is violated.

**Proposition.** Let  $N'$  be any subset of  $N$  satisfying (7) and suppose

$$\sum_{j \in Q} a_j \leq a_0 \quad (8)$$

holds for all  $Q \subset N'$  such that  $|Q| = \pi_0$ , but for no  $Q \subset N'$  such that  $|Q| > \pi_0$ . Then

$$\sum_{j \in N'} x_j \leq \pi_0 \quad (9)$$

is a facet of  $P_I^{N'}$ , where  $N^* = N - N'$ .

**Proof.** From the assumption that (7) holds, we have that for every zero-one vector  $x$ ,  $\sum_{j \in N'} x_j > \pi_0$  implies  $x \notin P_I^{N'}$ ; hence (9) is satisfied by all  $x \in P_I^{N'}$ . By this assumption, furthermore,  $n' = |N'| \geq \pi_0 + 1$ . Consequently, letting  $k \in N' - Q$  for some  $Q \subseteq N'$  with  $|Q| = \pi_0$ , we find  $\pi_0 + 1$  linearly independent zero-one vectors  $x^i$  satisfying (9) with equality and with the property that  $j \in N - (Q \cup \{k\})$  implies  $x_j^i = 0$  for  $i = 1, \dots, \pi_0 + 1$ . The remaining  $n' - (\pi_0 + 1)$  vectors  $x^i$  for  $i = \pi_0 + 2, \dots, n'$  are constructed so as to ensure a lower triangular structure in the rows and columns that correspond to the index set  $\{\pi_0 + 2, \dots, n'\}$ . The latter construction is possible, since (8) holds for all subsets of  $N'$  with cardinality  $\pi_0$ . The proposition also follows from Corollary 4.4 of reference 13.

**Example.** Suppose that  $m = 1$  in (1) and consider the inequality in zero-one variables

$$x_2 + 66x_3 + 66x_4 + 66x_5 + 65x_6 + 38x_7 \leq 205. \quad (10)$$

Taking  $N'$  to be the index set  $N' = \{2, 3, 4, 5, 6\}$ , we easily check that  $N'$  satisfies the requirements of the proposition. Consequently, the inequality

$$x_2 + x_3 + x_4 + x_5 + x_6 \leq 3 \quad (11)$$

provides a facet for  $P_I^{(1,7)}$ . In order to 'extend' (11) to a facet of  $P_I$ , we set  $T = \{7, 1\}$  and solve  $(H_q)$ ,  $q = 1, 2$ . Hence

$$z_1 = \max x_2 + x_3 + x_4 + x_5 + x_6,$$

$$71x_2 + 66x_3 + 66x_4 + 66x_5 + 65x_6 \leq 167 = 205 - 38,$$

$$x_i = 0 \text{ or } 1, i = 2, \dots, 6,$$

yields  $z_1 = 2$ . Consequently, a new inequality is given by

$$x_2 + x_3 + x_4 + x_5 + x_6 + x_7 \leq 3$$

and this inequality is a facet for  $P_I^{(1)}$ . Solving  $(H_2)$ , we obtain finally

$$2x_1 + x_2 + x_3 + x_4 + x_5 + x_6 + x_7 \leq 3. \quad (12)$$

The inequality provided by (12) is a facet for the convex hull of zero-one solutions to (10). (In fact, we have chosen an example where (10) and (12) have identical zero-one solutions.) The other (and in this case the only other) facets of  $P_I$  are the trivial facets  $x_j \geq 0$  and the nontrivial facets  $x_j \leq 1$ ,  $j = 1, \dots, 7$ . In our particular example, the order of  $T$  did not affect the resulting facet. This is, however, not true in general. Finally, we would like to point out that the upper bound on the number of facets needed to describe  $P_I$  for an arbitrary underlying polytope  $P$  is given by  $2^{n-1}$  (see reference 2).

## NOTES

1. Since this paper was originally written (March 1973), a number of papers have appeared that deal with topics very closely related to the subject of this note. A (possibly incomplete) list of papers that the interested reader should consult is given in the "Additional References" below.
2. The results given here were independently and simultaneously obtained by Professor Padberg and G. L. Nemhauser and L. E. Trotter, Jr.<sup>[13]</sup> The presentation in reference 13 is somewhat more general, but also more abstract. To avoid duplication, this note is an abbreviated version of the original paper. The Editor.

## ACKNOWLEDGMENT

I am indebted to Egon Balas for helpful discussions and remarks on the topic of this paper. I did the research for this paper while visiting the University of Waterloo in March-April, 1973. Financial support by NRC

## REFERENCES

1. R. GARFINKEL AND G. L. NEMHAUSER, *Integer Programming*, Wiley, New York, 1972.
2. R. G. JEROSLOW, "On Defining Sets of Vertices of the Hypercube by Linear Inequalities," Working paper, Carnegie-Mellon University, January 1973.
3. M. W. PADBERG, "On the Facial Structure of Set Packing Polyhedra," *Math. Programming* **5**, 199-215 (1973).
4. L. E. TROTTER, "Solution Characteristics and Algorithms for the Vertex Packing Problem," Ph.D. Thesis, Cornell University, January 1973.

## ADDITIONAL REFERENCES

5. E. BALAS, "Facets of the Knapsack Polytope," *Math. Programming* **8**, 146-164 (1975).
6. E. BALAS, "On the Facial Structure of the Knapsack Polytope," paper presented at the NATO Advanced Study Institute on Combinatorial Programming, Versailles, September 2-13, 1974.
7. E. BALAS AND E. ZEMEL, "Facets of the Knapsack Polytope from Minimal Covers," Carnegie-Mellon University, Management Sciences Research Report No. 352, December 1974.
8. F. GLOVER, "Unit-Coefficient Inequalities for Zero-one Programming," University of Colorado, Management Sciences Report Series, No. 73-7, July 1973.
9. P. L. HAMMER, E. L. JOHNSON, AND U. N. PELED, "Regular 0-1 Programs," University of Waterloo, Combinatorics and Optimization Research Report, CORR No. 73-18, September 1973.
10. P. L. HAMMER, E. L. JOHNSON, AND U. N. PELED, "Facets of Regular 0-1 Polytopes," *Math. Programming* **8**, 179-206 (1975).
11. P. L. HAMMER, E. L. JOHNSON, AND U. N. PELED, "The Role of Master Polytopes in the Unit Cube," University of Waterloo, Combinatorics and Optimization Research Report, CORR No. 74-25, October 1974.
12. E. L. JOHNSON, "A Class of Facets of the Master 0-1 Knapsack Polytope," Thomas J. Watson Research Center Report RC-5106, IBM Research, October 30, 1974.
13. G. L. NEMHAUSER AND L. E. TROTTER, "Properties of Vertex Packing and Independence System Polyhedra," *Math. Programming* **6**, 48-61 (1974).
14. L. A. WOLSEY, "Faces for a Linear Inequality in 0-1 Variables," *Math. Programming* **8**, 165-178 (1975).
15. L. A. WOLSEY, "Facets and Strong Valid Inequalities for Integer Programming," Université Catholique de Louvain, CORE Discussion paper, April 1974.

Copyright 1975, by INFORMS, all rights reserved. Copyright of Operations Research is the property of INFORMS: Institute for Operations Research and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.