

DM GALPROP

Andrey Egorov

E-mail: ael14@yandex.ru

Contents

1	Introduction	1
2	DM source	1
3	Installation and usage	2

1 Introduction

In general, the GALPROP code solves the transport equation for various species of cosmic rays in the Galaxy and computes various types of emission from them. A full description of the code and respective updates can be found at [1]. Also a detailed general manual is included in the “doc” folder of the code distribution (i.e. the whole tarball for download). This brief manual describes the DM part only AND supersedes the respective part of the general manual. The current DM version is based on GALPROP v54r2766. Comments and questions on DM GALPROP can be addressed to the author’s e-mail above.

2 DM source

WIMP annihilation everywhere in the Galaxy is expected to produce various energetic final products: electrons, protons, gammas, neutrinos etc. and their respective antiparticles. Then they propagate and give a rise to various emissions from radio to gamma through synchrotron, ICS, bremsstrahlung etc. Gammas and neutrinos are obviously free-streaming products. The injection of all these species is described by the following source term $q(r, p)$, which would enter the transport equation:

$$q(r, p) = \frac{1}{2} \langle \sigma v \rangle \left(\frac{\rho_{\text{DM}}(r)}{m_\chi} \right)^2 \xi_{\text{DM}}(r) \frac{dN}{dp}(p), \quad (2.1)$$

where $\langle \sigma v \rangle$ denotes the WIMP annihilation cross section, $\rho_{\text{DM}}(r)$ is the DM density distribution, m_χ is the WIMP mass, $\xi_{\text{DM}}(r)$ is the DM annihilation rate boost factor due to substructures and $\frac{dN}{dp}(p)$ represents the spectral yield of any specie of interest. The latter depends on both WIMP mass and annihilation channel, and comes from particle physics data. Below some details are provided on every key component of the source term.

- **DM density distribution.** Currently GALPROP supports four different profiles: generalized NFW, Isothermal, Einasto and Burkert. Their exact definitions are:

$$\begin{aligned} \text{gNFW : } \rho_{\text{gNFW}}(r) &= \frac{\rho_s}{(r/r_s)^\gamma (1 + r/r_s)^{3-\gamma}} \\ \text{Einasto : } \rho_{\text{Ein}}(r) &= \rho_s \exp \left\{ -\frac{2}{\alpha} \left[\left(\frac{r}{r_s} \right)^\alpha - 1 \right] \right\} \\ \text{Isothermal : } \rho_{\text{Iso}}(r) &= \frac{\rho_s}{1 + (r/r_s)^2} \\ \text{Burkert : } \rho_{\text{Bur}}(r) &= \frac{\rho_s}{(1 + r/r_s)(1 + (r/r_s)^2)} \end{aligned} \quad (2.2)$$

GALPROP does averaging of the DM density squared (which enters eq. (2.1)) inside every spatial grid cell in order to take into account possible steep variations of the density. Also, considering that the gNFW profile in eq. (2.2) is singular in the center, there is an option (see below) to leave the density flat below some radius for that profile. This prevents formation of an artificial spike in the center, where the real DM distribution is known quite poorly in fact.

- **DM annihilation rate boost factor.** Substructures boost the DM density and, hence, the annihilation rate. However, in a small vicinity of the Galactic center ~ 1 kpc they almost don't survive due to tidal disruption. Hence, the boost factor can be negligible there. However, it can reach significant values up to ~ 10 far from the center. This local boost factor is assumed to be a function of the radial distance only and implemented based on the results of [2] (which were presumably derived assuming the NFW profile for the smooth component). If one would like to stay conservative in computation of the emission intensity due to DM, there is an option to turn off the boost at all.
- **DM annihilation yields.** This comes from particle physics and defines how many stable products (like electrons and others) are being produced in one annihilation depending on their energy. Such a spectra begin at kinematic threshold $m_\chi c^2$ and typically rise with energy decrease. The yields are implemented as numerical tables for every final product and annihilation channel based on results of [3–5]. Specifically, the latest version of the yields is used, which includes electroweak corrections. Yields are taken as non-polarized (if applicable), meaning average of the Left/Right or Longitudinal/Transverse. Currently, 7 annihilation channels are implemented: $\chi\chi \rightarrow b\bar{b}, \tau^+\tau^-, W^+W^-, \mu^+\mu^-, q\bar{q}, ZZ, hh$ (with $q \equiv u, d, s$).

3 Installation and usage

A typical installation procedure is described in the “README” file of the distribution folder. A user also has to download FITS+DM data folder and pass its path to the `./configure` parameter “`–with-fitsdata`”.

In general, it's recommended to run the 3D version of the code rather than 2D. The former one is more precise and better tested. Currently, DM GALPROP is capable to compute propagation of DM electrons, positrons and antiprotons. Regarding computation of an emission, the synchrotron from electrons and positrons is currently well developed and tested (see e.g. our work [6] based on the current version). In principle, user can compute gamma rays due to DM too. However, the gamma ray part is not finalized yet. In order to get the prompt gammas only, one has to turn on the following switches in `galdef` file: `DM_gammas`, `gamma_rays` AND any specie like e.g. `DM_electrons`. In order to get secondary gammas due to DM, which come from ICS and bremsstrahlung of DM electrons, one has to turn on respectively `IC_` and `bremss`.

Notes on 3D version: it's extremely recommended to choose the spatial grid steps dx, dy, dz in a way that one of the grid points would fall exactly at the Galactic center $x=0, y=0, z=0$. This is because the DM source is very concentrated around the center and, hence, the central grid point has a very important role among all others. And it has to be “resolved” in computation through the best way possible, especially for the case of cuspy profiles like NFW. Realistic runs typically require $dx \approx dy \approx dz \approx 0.2$ kpc.

DM parameters in galdef file are described below. As was already mentioned, all other parameters are described in the general GALPROP manual. There is also an example of a typical galdef file for the DM run in the “GALDEF” folder of the main distribution - “galdef_54_testDM”.

DM_positrons	= 0	1=compute DM positrons
DM_electrons	= 0	1=compute DM electrons
DM_antiprotons	= 0	1=compute DM antiprotons
DM_gammas	= 0	1=compute DM gammas
DM_int0	= 1	DM density profile $\rho_{\text{DM}}(r)$: 0=generalized NFW; 1=Isothermal; 2=Einasto; 3=Burkert; 9=DarkSUSY
DM_int1	= 1	9=DarkSUSY DM source
DM_int2	= 1	DM annihilation channel: 1= $b\bar{b}$, 2 = $\tau^+\tau^-$, 3 = W^+W^- , 4 = $\mu^+\mu^-$, 5 = $q\bar{q}$, 6 = ZZ , 7 = hh
DM_int3	= 1	Substructure on/off ($\xi_{\text{DM}}(r)$): 1=based on Kamionkowski (which is likely for NFW only)
DM_int4	= 1	doubles lepton source function (0=no) in order to save comp. time by propagating only one specie; if turned on, requires either one of DM_positrons or DM_electrons to be 0; should be a good approximation; doesn't work in case of the presence of positron annihilation
DM_int5	= 1	not used
DM_int6	= 1	not used
DM_int7	= 1	not used
DM_int8	= 1	not used
DM_int9	= 1	0=fixed increment synchrotron frequency grid (set by nu_synch_min, nu_synch_max, nu_synch_factor), >0=number of arbitrary desired frequencies
DM_double0	= 28.44	scale radius r_s of the DM density profile, kpc
DM_double1	= 0.033	scale density ρ_s of the DM density profile, GeV cm^{-3}
DM_double2	= 19.1	DM particle mass m_χ , GeV; must be in the range [5..100000] AND greater than the rest mass of the primary products (set by DM_int2); inaccuracies are possible close to the thresholds
DM_double3	= 15.0e6,30.0e6,45.0e6	list of coma-separated arbitrary desired synchrotron frequencies (Hz), if DM_int9>0
DM_double4	= not used	
DM_double5	= not used	
DM_double6	= 1.1	inner slope parameter γ for the gNFW DM density profile (1.0 gives the canonical NFW)
DM_double7	= 0.17	α parameter for the Einasto profile
DM_double8	= 4.2e-10	truncation radius for the gNFW profile, below which the DM density stays constant, kpc (must be greater than 0!)
DM_double9	= 3.0e-26	thermally averaged annihilation cross section $\langle\sigma v\rangle$, cm^3/s

Acknowledgments

The author is grateful to Andrew Strong, Elena Pierpaoli and Jennifer Gaskins for a valuable help with preparation of this version.

References

- [1] <http://galprop.stanford.edu/>, <https://sourceforge.net/projects/galprop>.
- [2] M. Kamionkowski, S. M. Koushiappas, and M. Kuhlen, *Galactic Substructure and Dark Matter Annihilation in the Milky Way Halo*, *Phys.Rev.* **D81** (2010) 043532, [[arXiv:1001.3144](#)].
- [3] <http://www.marcocirelli.net/PPPC4DMID.html>.
- [4] M. Cirelli, G. Corcella, A. Hektor, G. Hutsi, M. Kadastik, et al., *PPPC 4 DM ID: A Poor Particle Physicist Cookbook for Dark Matter Indirect Detection*, *JCAP* **1103** (2011) 051, [[arXiv:1012.4515](#)].
- [5] P. Ciafaloni, D. Comelli, A. Riotto, F. Sala, A. Strumia, et al., *Weak Corrections are Relevant for Dark Matter Indirect Detection*, *JCAP* **1103** (2011) 019, [[arXiv:1009.0224](#)].
- [6] A. E. Egorov, J. M. Gaskins, E. Pierpaoli, and D. Pietrobon, *Dark matter implications of the WMAP-Planck Haze*, *JCAP* **3** (Mar., 2016) 060, [[arXiv:1509.05135](#)].