

## Exercise 1

Recall that we have already explained in lecture that all rational function of a single variable can be integrated in finite terms. Starting from there, we introduce integrals of the form, known as the **binomial integrals**:

$$J_{p,q} = \int (a + bz)^p z^q dz, a, b \in \mathbb{R}, p, q \in \mathbb{Q}. \quad (1)$$

This exercise aims at studying the rationalization of the binomial integral, as well as some of its applications.

## Question 1

Assume that  $p \in \mathbb{Z}$ , rationalise the integrand. ?

Since  $q \in \mathbb{Q}$ , this implies there exists  $m \in \mathbb{Z}, n \in \mathbb{N}$  such that  $q = \frac{m}{n}$ . Then,  $J_{p,q}$  can be written as

$$J_{p,q} = \int (a + bz)^p z^{\frac{m}{n}} dz.$$

Let  $z = x^n$ , then  $dz = n \cdot x^{n-1} dx$ . Replace  $z$  with  $x$  we have

$$J_{p,q} = \int (a + b \cdot x^n)^p \cdot (x^n)^{\frac{m}{n}} \cdot n \cdot x^{n-1} dx.$$

Simplify it and we get

$$J_{p,q} = n \int (a + b \cdot x^n)^p \cdot x^{m+n-1} dx.$$

Now, by binomial theorem we can see the integrand is rationalised, as needed. ☺

**Question 2**

Assume that  $q \in \mathbb{Z}$ , rationalise the integrand.

?

Since  $p \in \mathbb{Q}$ , this implies there exists  $m \in \mathbb{Z}, n \in \mathbb{N}$  such that  $p = \frac{m}{n}$ . Then,  $J_{p,q}$  can be written as

$$J_{p,q} = \int (a + bz)^{\frac{m}{n}} z^q dz.$$

If  $b = 0$  then the integrand is trivially rational, thus we consider the case when  $b \neq 0$ . Let  $x^n = a + bz$ , then  $z = \frac{x^n - a}{b}$  and  $n \cdot x^{n-1} dx = b dz$ . Replace  $z$  with  $x$  we have

$$J_{p,q} = \int (x^n)^{\frac{m}{n}} \cdot \left( \frac{x^n - a}{b} \right)^q \cdot \frac{n \cdot x^{n-1}}{b} dx.$$

Simplify it and we get

$$J_{p,q} = \frac{n}{b^{q+1}} \int x^{m+n-1} \cdot (x^n - a)^q dx.$$

Now, by binomial theorem we can see the integrand is rationalised, as needed.

☺

## Question 3

Assume that  $p + q \in \mathbb{Z}$ , rationalise the integrand.

?

**Hint.** Write the integrand as

$$\int \left( \frac{a + bz}{z} \right)^p z^{p+q} dz.$$



If  $p \in \mathbb{Z}$  then this case is rationalised by Question 1. Thus we focus on the case when  $p \in \mathbb{Q} \setminus \mathbb{Z}$ , that is,  $p = \frac{m}{n}$  for some  $m \in \mathbb{Z}, n \in \mathbb{N}$ . First, we change the equation to

$$J_{p,q} = \int \left( \frac{a + bz}{z} \right)^p z^{p+q} dz.$$

Now, let  $x^n = \frac{a+bz}{z} = \frac{a}{z} + b$ , then  $z = \frac{a}{x^n - b}$  and  $n \cdot x^{n-1} dx = -\frac{a}{z^2} dz$ . Now replace  $z$  with  $x$  we have

$$\begin{aligned} J_{p,q} &= \int (x^n)^{\frac{m}{n}} \cdot \left( \frac{a}{x^n - b} \right)^{p+q} \cdot \left( \frac{n \cdot x^{n-1} \cdot z^2}{-a} \right) dx \\ &= \int x^m \cdot \frac{a^{p+q}}{(x^n - b)^{p+q}} \cdot \left( \frac{n \cdot x^{n-1} \cdot \frac{a^2}{(x^n - b)^2}}{-a} \right) dx \\ &= n \int x^m \cdot \frac{a^{p+q}}{(x^n - b)^{p+q}} \cdot \left( -\frac{ax^{n-1}}{(x^n - b)^2} \right) dx \\ &= -n \cdot a^{p+q+1} \int x^m \cdot \frac{1}{(x^n - b)^{p+q}} \cdot \left( x^{n-1} \cdot \frac{1}{(x^n - b)^2} \right) dx \\ &= -n \cdot a^{p+q+1} \int x^{m+n-1} \cdot \frac{1}{(x^n - b)^{p+q+2}} dx \\ &= -n \cdot a^{p+q+1} \int x^{m+n-1} \cdot (x^n - b)^{-2-p-q} dx \end{aligned}$$

Now, we can see that the integrand is rationalised, as needed.



**Remark 1.** So far we have shown an interesting conclusion: if either  $p$ , or  $q$ , or  $p + q$  is an integer, then the function can be integrated in finite terms.



## Question 4

Prove that

- If  $p \neq -1$ , then

$$J_{p,q} = -\frac{(a+bz)^{p+1}z^{q+1}}{a(p+1)} + \frac{p+q+2}{a(p+1)}J_{p+1,q}$$

- If  $q \neq -1$ , then

$$J_{p,q} = \frac{(a+bz)^{p+1}z^{q+1}}{a(q+1)} - b\frac{p+q+2}{a(q+1)}J_{p,q+1}$$

?

*Proof.*

- 1) When  $p \neq -1$ , using integration by parts we have

$$\begin{aligned} J_{p,q} &= \int (a+bz)^p \cdot z^q \, dz \\ &= \int \left( \frac{a+bz}{z} \right)^p \cdot z^{p+q} \, dz \\ &= \int \left( \frac{a}{z} + b \right)^p \cdot z^{p+q} \, dz \\ &= \int z^{p+q} \cdot \left( -\frac{z^2}{a} \right) \, d \left( \frac{\left( \frac{a}{z} + b \right)^{p+1}}{p+1} \right) \\ &= -\frac{1}{a(p+1)} \int z^{p+q+2} \cdot d \left( \frac{a}{z} + b \right)^{p+1} \\ &= -\frac{\frac{(a+bz)^{p+1}}{z^{p+1}} \cdot z^{p+q+2}}{a(p+1)} + \frac{1}{a(p+1)} \int \left( \frac{a}{z} + b \right)^{p+1} d(z^{p+q+2}) \\ &= -\frac{(a+bz)^{p+1} \cdot z^{q+1}}{a(p+1)} + \frac{p+q+2}{a(p+1)} \int \frac{(a+bz)^{p+1}}{z^{p+1}} \cdot z^{p+q+1} \, dz \\ &= -\frac{(a+bz)^{p+1} \cdot z^{q+1}}{a(p+1)} + \frac{p+q+2}{a(p+1)} \int (a+bz)^{p+1} \cdot z^q \, dz \\ &= -\frac{(a+bz)^{p+1} \cdot z^{q+1}}{a(p+1)} + \frac{p+q+2}{a(p+1)} J_{p+1,q} \end{aligned} \tag{2}$$

- 2) When  $q \neq -1$ , we first consider the integral form of  $J_{p+1,q}$  in (1), then substitute our result into (2), and the ending result is the desired formula.  
To this end, consider  $J_{p+1,q} = \int (a + bz)^{p+1} \cdot z^q dz$ , using integration by parts we have

$$\begin{aligned}
 J_{p+1,q} &= \int (a + bz)^{p+1} \cdot z^q dz \\
 &= \int (a + bz)^{p+1} d\left(\frac{z^{q+1}}{q+1}\right) \\
 &= \frac{1}{q+1} (a + bz)^{p+1} \cdot z^{q+1} - \frac{1}{q+1} \int z^{q+1} d(a + bz)^{p+1} \\
 &= \frac{1}{q+1} (a + bz)^{p+1} \cdot z^{q+1} - \frac{b(p+1)}{q+1} \int (a + bz)^p \cdot z^{q+1} dz \\
 &= \frac{1}{q+1} (a + bz)^{p+1} \cdot z^{q+1} - \frac{b(p+1)}{q+1} J_{p,q+1}
 \end{aligned} \tag{3}$$

Now, we substitute (3) back to (2), then the equation becomes

$$\begin{aligned}
 J_{p,q} &= -\frac{(a + bz)^{p+1} \cdot z^{q+1}}{a(p+1)} + \frac{p+q+2}{a(p+1)} J_{p+1,q} \\
 &= -\frac{(a + bz)^{p+1} \cdot z^{q+1}}{a(p+1)} + \frac{p+q+2}{a(p+1)} \left( \frac{1}{q+1} (a + bz)^{p+1} \cdot z^{q+1} - \frac{b(p+1)}{q+1} J_{p,q+1} \right) \\
 &= -\frac{(a + bz)^{p+1} \cdot z^{q+1}}{a(p+1)} + \frac{1}{a} \left[ 1 + \frac{q+1}{p+1} \right] \cdot \left( \frac{1}{q+1} (a + bz)^{p+1} \cdot z^{q+1} - \frac{b(p+1)}{q+1} J_{p,q+1} \right) \\
 &= -\frac{(a + bz)^{p+1} \cdot z^{q+1}}{a(p+1)} + \frac{1}{a} \left[ \frac{z^{q+1}}{q+1} \cdot (a + bz)^{p+1} - \frac{b(p+1)}{q+1} J_{p,q+1} \right] \\
 &\quad + \frac{q+1}{a(p+1)} \left[ \frac{z^{q+1}}{q+1} (a + bz)^{p+1} - \frac{b(p+1)}{q+1} J_{p,q+1} \right] \\
 &= -\frac{(a + bz)^{p+1} z^{q+1}}{a(p+1)} + \frac{1}{a(q+1)} (a + bz)^{p+1} z^{q+1} - \frac{b(p+1)}{a(q+1)} J_{p,q+1} \\
 &\quad + \frac{(a + bz)^{p+1} z^{q+1}}{a(p+1)} - \frac{b}{a} J_{p,q+1} \\
 &= \frac{(a + bz)^{p+1} z^{q+1}}{a(q+1)} - \frac{b(p+1+q+1)}{a(q+1)} J_{p,q+1} \\
 &= \frac{(a + bz)^{p+1} z^{q+1}}{a(q+1)} - b \frac{p+q+2}{a(q+1)} J_{p,q+1}
 \end{aligned}$$

which is precisely the equation required, as needed.

QUOD  
ERAT  
DEMONSTRATUM ■

## Question 5

Based on the previous question, prove that if  $p + q \neq -1$ , then

$$J_{p,q} = \frac{(a+bz)^p z^{q+1}}{p+q+1} + \frac{ap}{p+q+1} J_{p-1,q}$$

$$J_{p,q} = \frac{(a+bz)^{p+1} z^q}{b(p+q+1)} - \frac{aq}{b(p+q+1)} J_{p,q-1}$$

?

*Proof.* We first show the first equation. Consider (1), we will use integration by parts to show the desired equation. That is,

$$\begin{aligned} J_{p,q} &= \int (a+bz)^p z^q dz \\ &= \int \left( \frac{a+bz}{z} \right)^p z^{p+q} dz \\ &= \int \left( \frac{a+bz}{z} \right)^p d \left( \frac{z^{p+q+1}}{p+q+1} \right) \\ &= \frac{(a+bz)^p}{z^p} \cdot \frac{z^{p+q+1}}{p+q+1} - \frac{1}{p+q+1} \int z^{p+q+1} d \left( \frac{a+bz}{z} \right)^p \\ &= \frac{(a+bz)^p z^{q+1}}{p+q+1} - \frac{1}{p+q+1} \int z^{p+q+1} d \left( \frac{a}{z} + b \right)^p \\ &= \frac{(a+bz)^p z^{q+1}}{p+q+1} - \frac{1}{p+q+1} \int z^{p+q+1} p \left( \frac{a}{z} + b \right)^{p-1} \left( -\frac{a}{z^2} \right) dz \\ &= \frac{(a+bz)^p z^{q+1}}{p+q+1} + \frac{ap}{p+q+1} \int z^{p+q+1} \left( \frac{a+bz}{z} \right)^{p-1} z^{-2} dz \\ &= \frac{(a+bz)^p z^{q+1}}{p+q+1} + \frac{ap}{p+q+1} \int (a+bz)^{p-1} z^{p+q+1-p+1-2} dz \\ &= \frac{(a+bz)^p z^{q+1}}{p+q+1} + \frac{ap}{p+q+1} \int (a+bz)^{p-1} z^q dz \\ &= \frac{(a+bz)^p z^{q+1}}{p+q+1} + \frac{ap}{p+q+1} J_{p-1,q} \end{aligned} \tag{4}$$

With (4) we can show the second equation. Similar to Question 4, we first consider the integral form of  $J_{p-1,q}$  in (1), then substitute our result into (4), and the ending result is the desired formula.

$$\begin{aligned}
 J_{p-1,q} &= \int (a + bz)^{p-1} z^q dz \\
 &= \frac{1}{pb} \int z^q d(a + bz)^p \\
 &= \frac{1}{pb} (a + bz)^p z^q - \frac{q}{pb} \int (a + bz)^p z^{q-1} dz \\
 &= \frac{1}{pb} (a + bz)^p z^q - \frac{q}{pb} J_{p,q-1}
 \end{aligned} \tag{5}$$

Now, we substitute (5) back to (4), then the equation becomes

$$\begin{aligned}
 J_{p,q} &= \frac{(a + bz)^p z^{q+1}}{p + q + 1} + \frac{ap}{p + q + 1} J_{p-1,q} \\
 &= \frac{(a + bz)^p z^{q+1}}{p + q + 1} + \frac{ap}{p + q + 1} \left( \frac{1}{pb} (a + bz)^p z^q - \frac{q}{pb} J_{p,q-1} \right) \\
 &= \frac{(a + bz)^p z^{q+1}}{p + q + 1} + \frac{a}{p + q + 1} \left( \frac{1}{b} (a + bz)^p z^q - \frac{q}{b} J_{p,q-1} \right) \\
 &= \frac{(a + bz)^p z^{q+1}}{p + q + 1} + \frac{a}{p + q + 1} \frac{(a + bz)^p z^q}{b} - \frac{aq}{b(p + q + 1)} J_{p,q-1} \\
 &= \frac{bz(a + bz)^p z^q}{b(p + q + 1)} + \frac{a(a + bz)^p z^q}{b(p + q + 1)} - \frac{aq}{b(p + q + 1)} J_{p,q-1} \\
 &= \frac{(a + bz)^{p+1} z^q}{b(p + q + 1)} - \frac{aq}{b(p + q + 1)} J_{p,q-1}
 \end{aligned}$$

which is precisely the equation required, as needed.

QUOD  
ERAT  
DEMONSTRATUM ■

Now with all the information obtained above, let's study an example to get some feelings. Define for  $m \in \mathbb{Z}$  the integral

$$H_m = \int \frac{x^m}{\sqrt{1-x^2}} dx, \quad x > 0.$$

In the sequel, we will first transform  $H_m$  into a binomial integral. Since the binomial integral can be computed by iteration, it in turn provides a recursive relation for  $H_m, m \in \mathbb{Z}$ . To conclude, we aim at completely (at least formally) solve the problem of integrate  $H_m$  in finite terms for any  $m \in \mathbb{Z}$ .

**Question 6**

By introducing the substitution  $z = x^2$ , show that

$$H_m = J_{-\frac{1}{2}, \frac{m-1}{2}}.$$

?

*Proof.* Let  $z = x^2$ , then  $dz = 2x dx$ . Replace  $x$  with  $z$  we have

$$H_m = \int \frac{z^{\frac{m}{2}}}{\sqrt{1-z}} \cdot \frac{dz}{2\sqrt{z}} = \frac{1}{2} \int \frac{z^{\frac{m-1}{2}}}{\sqrt{1-z}} dz = \int \frac{z^{\frac{m-1}{2}}}{\sqrt{4-4z}} dz = \int (4-4z)^{-\frac{1}{2}} \cdot z^{\frac{m-1}{2}} dz.$$

Let  $a = 4, b = -4, p = -\frac{1}{2}, q = \frac{m-1}{2}$ , then we have

$$H_m = \int (4-4z)^{-\frac{1}{2}} \cdot z^{\frac{m-1}{2}} dz = J_{-\frac{1}{2}, \frac{m-1}{2}}.$$

QUOD  
ERAT  
DEMONSTRATUM ■



## Question 7

Prove that  $\forall m > 1$ ,

$$H_m = -\frac{1}{m}x^{m-1}\sqrt{1-x^2} + \frac{m-1}{m}H_{m-2}$$

?

**Hint.** Perhaps use consequence in question 5.



*Proof.* Assume  $m > 1$ . Let  $a = 4, b = -4, p = -\frac{1}{2}, q = \frac{m-1}{2}$  (note that here  $p + q \neq -1$ ),  $z = x^2$ , then we have

$$\begin{aligned} H_m = J_{-\frac{1}{2}, \frac{m-1}{2}} &= \frac{(a+bz)^{p+1}z^q}{b(p+q+1)} - \frac{aq}{b(p+q+1)} J_{-\frac{1}{2}, \frac{(m-2)-1}{2}} \\ &= \frac{(4-4z)^{\frac{1}{2}}z^{\frac{m-1}{2}}}{-4(\frac{m-1}{2} - \frac{1}{2} + 1)} - \frac{4(\frac{m-1}{2})}{-4(\frac{m-1}{2} - \frac{1}{2} + 1)} \int (4-4z)^{-\frac{1}{2}} z^{\frac{(m-2)-1}{2}} dz \\ &= \frac{2(1-x^2)^{\frac{1}{2}}x^{m-1}}{-2(m-1-1+2)} - \frac{2(m-1)}{-2(m-1-1+2)} \int \frac{1}{2}(1-x^2)^{-\frac{1}{2}}x^{(m-2)-1}2x dx \\ &= -\frac{(1-x^2)^{\frac{1}{2}}x^{m-1}}{m} - \frac{m-1}{-m} \int (1-x^2)^{-\frac{1}{2}}x^{m-2} dx \\ &= -\frac{1}{m}x^{m-1}\sqrt{1-x^2} + \frac{m-1}{m} \int \frac{x^{m-2}}{\sqrt{1-x^2}} dx \\ &= -\frac{1}{m}x^{m-1}\sqrt{1-x^2} + \frac{m-1}{m}H_{m-2} \end{aligned}$$

since we have chosen an arbitrary  $m > 1$ , thus we have shown the equation holds for all  $m > 1$ , as required.

QUOD  
ERAT  
DEMONSTRATUM ■

**Question 8**Compute  $H_0, H_1$  explicitly.

?

$$\begin{aligned} H_0 &= \int \frac{1}{\sqrt{1-x^2}} dx \\ &= \arcsin x + C \end{aligned} \tag{6}$$

Note that the equality follows by formula, which can be verified by taking the derivative of (6). ☺

$$\begin{aligned} H_1 &= \int \frac{x}{\sqrt{1-x^2}} dx \\ &= \frac{1}{2} \int (1-x^2)^{-\frac{1}{2}} d(x^2) \\ &= -(1-x^2)^{\frac{1}{2}} + C \end{aligned}$$

☺

**Question 9**

Based on the previous sub-question(s), conclude that  $\forall m \in \mathbb{Z}_{\geq 0}$ ,  $H_m$  can be integrated in finite terms. ?

*Proof.* We show by strong induction. From Question 8, we have shown  $H_0, H_1$  can be integrated in finite terms. Let  $H_0, H_1$  be the base cases. Assume  $H_n$  holds when  $n \in \{0, 1, \dots, m\}$  for some  $m \in \mathbb{Z}_{\geq 0}$ . Then, we want to show  $H_{m+1}$  can also be integrated in finite terms. To this end, consider 2 cases.

When  $m + 1 \in \{0, 1\}$ , this is immediately covered by our base cases.

When  $m + 1 \geq 2$ , by Question 7, we have

$$H_{m+1} = -\frac{1}{m+1}x^m\sqrt{1-x^2} + \frac{m}{m+1}H_{m-1},$$

since  $m > m - 1 \geq 0$ , this means by our inductive hypothesis  $H_{m-1}$  can be integrated in finite terms, this gives  $H_{m+1}$  can also be integrated in finite terms by our equation.

Therefore, by strong induction, we have shown  $H_m$  can be integrated in finite terms for all  $m \in \mathbb{Z}_{\geq 0}$ , as required.

QUOD  
ERAT  
DEMONSTRATUM ■

## Question 10

Prove that  $\forall m < -1$ ,

$$H_m = \frac{x^{m+1}\sqrt{1-x^2}}{m+1} + \frac{2+m}{1+m}H_{m+2}$$

?

**Hint.** Perhaps use consequence in question 4.



*Proof.* Choose any  $m < -1$ , fix  $a = 4, b = -4, p = -\frac{1}{2}, q = \frac{m-1}{2}$ , let  $z = x^2$  where  $dz = 2x dx$ , then, by formula from Question 4 and Question 6 we have

$$\begin{aligned} H_m = J_{-\frac{1}{2}, \frac{m-1}{2}} &= \frac{(a+bz)^{\frac{1}{2}} z^{\frac{m+1}{2}}}{a(q+1)} - b \frac{p+q+2}{a(q+1)} J_{-\frac{1}{2}, \frac{m+1}{2}} \\ &= \frac{(4-4z)^{\frac{1}{2}} z^{\frac{m+1}{2}}}{4(\frac{m-1}{2}+1)} - (-4) \frac{-\frac{1}{2} + \frac{m-1}{2} + 2}{4(\frac{m-1}{2}+1)} \int (4-4z)^{-\frac{1}{2}} z^{\frac{m+1}{2}} dz \\ &= \frac{2(1-x^2)^{\frac{1}{2}} x^{m+1}}{2(m+1)} + 2 \frac{-1+m-1+4}{2m-2+4} \int \frac{1}{2} (1-x^2)^{-\frac{1}{2}} x^{m+1} 2x dx \\ &= \frac{x^{m+1}\sqrt{1-x^2}}{m+1} + \frac{2+m}{1+m} \int \frac{x^{m+2}}{\sqrt{1-x^2}} dx \\ &= \frac{x^{m+1}\sqrt{1-x^2}}{m+1} + \frac{2+m}{1+m} H_{m+2} \end{aligned}$$

which is precisely the equation required, as needed.

QUOD  
ERAT  
DEM ■

**Question 11**Compute  $H_{-1}, H_{-2}$  explicitly.

?

**Claim 1**

$$\int \frac{1}{\sin x} dx = \ln |\csc x - \cot x| + C$$

◇

*Proof of Claim 1.*

$$\begin{aligned} \int \frac{1}{\sin x} dx &= \int \frac{-\cos x + 1}{\sin x(-\cos x + 1)} dx \\ &= \int \frac{-\frac{\cos x}{\sin^2 x} + \frac{1}{\sin^2 x}}{\frac{1}{\sin x} - \frac{\cos x}{\sin x}} dx \\ &= \int \frac{-\csc x \cot x + \csc^2 x}{\csc x - \cot x} dx \\ &= \int \frac{1}{\csc x - \cot x} d(\csc x - \cot x) \\ &= \ln |\csc x - \cot x| + C \end{aligned}$$

QUOD  
ERAT  
DEMONSTRATUM ■

## Claim 2

$$\int \frac{1}{\sqrt{x^2 - 1}} dx = \ln |x + \sqrt{x^2 - 1}| + C$$

◇

*Proof of Claim 2.* Let  $x = \sec u$ , then  $dx = \sec u \tan u du$ , thus we have

$$\begin{aligned} \int \frac{1}{\sqrt{x^2 - 1}} dx &= \int \frac{1}{\sqrt{\sec^2 u - 1}} \sec u \tan u du \\ &= \int \sec u du \\ &= \int \frac{1}{\cos u} du \\ &= - \int \frac{1}{\sin\left(\frac{\pi}{2} - u\right)} d\left(\frac{\pi}{2} - u\right) \\ &= - \ln \left| \csc\left(\frac{\pi}{2} - u\right) - \cot\left(\frac{\pi}{2} - u\right) \right| + C \\ &= \ln \left| \frac{1}{x - \sqrt{x^2 - 1}} \right| + C \quad \text{By Pythagorean Theorem} \\ &= \ln \left| \frac{1}{x - \sqrt{x^2 - 1}} \cdot \frac{x + \sqrt{x^2 - 1}}{x + \sqrt{x^2 - 1}} \right| + C \\ &= \ln |x + \sqrt{x^2 - 1}| + C \end{aligned} \tag{7}$$

If we assume  $x \in (-\infty, -1) \cup (1, \infty)$ , then  $u = \operatorname{arcsec} x \in (0, \pi) \setminus \{\frac{\pi}{2}\}$ . Hence, consider 2 cases.

When  $u \in (0, \pi/2)$ , we have  $\tan u > 0$ , thus our step (7) is valid. When  $u \in (\pi/2, \pi)$ , despite  $\tan u < 0$ , in this case  $x \in (-\infty, -1)$ , because the function  $\frac{1}{\sqrt{x^2 - 1}}$  is even, the integral of this case is same as the first case, hence we are allowed to change our  $\frac{\tan u}{|\tan u|}$  into 1.

QUOD  
ERAT  
DEMONSTRATUM ■

$$\begin{aligned} H_{-1} &= \int \frac{x^{-1}}{\sqrt{1-x^2}} dx \\ &= \int \frac{1}{x\sqrt{1-x^2}} dx \\ &= \int \frac{1}{x^2\sqrt{\frac{1}{x^2}-1}} dx \\ &= -\int \frac{1}{\sqrt{\left(\frac{1}{x}\right)^2-1}} d\left(\frac{1}{x}\right) \\ &= -\ln \left| \frac{1}{x} + \sqrt{\left(\frac{1}{x}\right)^2-1} \right| + C \end{aligned}$$



Using the substitution  $x = \sin u$ , and the pothagorean theorem, we have

$$\begin{aligned} H_{-2} &= \int \frac{x^{-2}}{\sqrt{1-x^2}} dx \\ &= \int \frac{1}{\sin^2 u \cos u} \cos u du \\ &= \int \csc^2 u du \\ &= -\cot u + C \\ &= -\frac{\sqrt{1-x^2}}{x} + C \end{aligned}$$



**Question 12**

Based on the previous sub-question(s), conclude that  $\forall m \in \mathbb{Z}_{<0}$ ,  $H_m$  can be integrated in finite terms. ?

*Proof.* We perform a variation of strong induction on  $\mathbb{Z}_{<0}$ . That is, first let  $H_{-1}, H_{-2}$  be the base cases which are proven in Question 11 that they can be integrated in finite terms. Now assume the statement holds for all  $H_n$  where  $n \in \{m, m+1, \dots, -2, -1\}$  for some  $m \in \mathbb{Z}_{<0}$ . Then, we want to show  $H_{m-1}$  can also be integrated in finite terms. To this end, consider 2 cases.

When  $m-1 \in \{-2, -1\}$ , this is immediately covered by our base cases.

When  $m-1 \leq -3$ , by Question 10, we have

$$H_{m-1} = \frac{x^m \sqrt{1-x^2}}{m} + \frac{1+m}{m} H_{m+1},$$

since  $m < m+1 \leq -1$ , this means by our inductive hypothesis  $H_{m+1}$  can be integrated in finite terms, this gives  $H_{m-1}$  can also be integrated in finite terms by our equation.

Therefore, by strong induction, we have shown  $H_m$  can be integrated in finite terms for all  $m \in \mathbb{Z}_{<0}$ , as required.

QUOD  
ERAT  
DEMONSTRATUM ■

**Conclusion:** We have shown that for any  $m \in \mathbb{Z}$ ,  $H_m$  can be integrated in finite terms.