

Sepsis Treatment Guidelines According To WHO guidelines 2022

➤ FIRST CHOICE (7 days) Of unknown origin

DRUG	FREQUENCY	DOSE	ROUTE OF ADMINISTRATION
CEFOTAXIME	Q8H	2000 mg	IV
OR			
CEFTRIAZONE	Q24H	2000 mg	IV

➤ SECOND CHOICE (7 days)

DRUG	FREQUENCY	DOSE	ROUTE OF ADMINISTRATION
AMIKACIN	Q24H	15 mg/kg	IV
OR			
GENTAMICIN	Q24H	5 mg/kg	IV

Sepsis Treatment Guidelines According To COOK guidelines 2022

(Adult of unknown origin)

➤ FIRST CHOICE

DRUG	FREQUENCY	DOSE	ROUTE OF ADMINISTRATION
CEFAZOLIN + GENTAMICIN	According to patient condition and clinicians experience		
IF POSSIBLE ABDOMINAL OR PELVIC INFECTION, ADD			
METRONIDAZOLE	According to patient condition and clinicians experience		
IF FAILING			
MEROPENEM +/- VANCOMYCIN	According to patient condition and clinicians experience		

➤ SECOND CHOICE

DRUG	FREQUENCY	DOSE	ROUTE OF ADMINISTRATION
AMOXICILLIN + CLAVULANATE + GENTAMICIN	According to patient condition and clinicians experience		
OR			
CHLORAMPHENICOL + GENTAMICIN	According to patient condition and clinicians experience		
OR			
VANCOMYCIN + GENTAMICIN	According to patient condition and clinicians experience		

(Children of unknown origin)

➤ FIRST CHOICE

DRUG	FREQUENCY	DOSE	ROUTE OF ADMINISTRATION
FLUCLOXACILLIN + CEFTRIAXONE	According to patient condition and clinicians experience		
IF FAILING			
FLUCLOXACILLIN +CEFTRIAXONE + GENTAMICIN	According to patient condition and clinicians experience		
IF FAILING &/OR SUSPECT MRSA			
VANCOMYCIN + GENTAMICIN	According to patient condition and clinicians experience		

➤ SECOND CHOICE

DRUG	FREQUENCY	DOSE	ROUTE OF ADMINISTRATION
CIPROFLOXACIN + VANCOMYCIN	According to patient condition and clinicians experience		
OR			
CHLORAMPHENICOL + CEFAZOLIN	According to patient condition and clinicians experience		
OR			
CHLORAMPHENICOL + CEFTRIAXONE	According to patient condition and clinicians experience		
OR			
CHLORAMPHENICOL + GENTAMICIN	According to patient condition and clinicians experience		

