Recurrence

Lines dividing the plane

Towers of Hanoi

Fibonacci staircase

Recall that we used induction to prove statements like

$$0+1+2+3+\ldots+n=\frac{n(n+1)}{2}$$
 (Arithmetic series)

$$1 + a + a^2 + a^3 + \dots + a^n = \frac{1 - a^{n+1}}{1 - a}$$
 (Geometric series)

Recurrence

Lines dividing the plane

Towers of Hanoi

Fibonacci staircase

Let's denote

$$Sum(n) = 1 + 2 + 3 + ... + n.$$

Question 1:

If we already know that Sum(99) = 4950, compute Sum(100).

Question 2:

If we know Sum(n-1), how to compute Sum(n)?

These observations suggest that this summation can be defined:

$$Sum(0) = 0$$

$$Sum(n) = Sum(n-1) + n \qquad (\forall n > 0)$$

This is a so called *recurrent* definition.

Recurrence

Lines dividing the plane

Towers of Hanoi

Fibonacci staircase

Exponentiation (to integer power n):

$$E(a,n)=a^n$$

Recurrently:

$$E(a,0) = 1$$

$$E(a,n) = E(a,n-1) \cdot a \qquad (\forall n > 0)$$

Recurrence

Lines dividing the plane

Towers of Hanoi

Fibonacci staircase

Factorial:

$$n! = 1 \cdot 2 \cdot \ldots \cdot n$$

Recurrently:

$$0! = 1$$

 $n! = (n-1)! \cdot n \quad (\forall n > 0)$

Translating a word problem into a recurrence.

A farmer had 10 cows when he started in January, 2000. The farm is working great, and each year the number of cows doubles.

Unfortunately, on every December 31, one of the cows gets abducted by a UFO.



Question: Find a recurrent expression for the number of cows in the beginning of each year.

$$(9 \times 2^{\Delta Y} + 1)$$

Recurrence


Lines dividing the plane

Towers of Hanoi

Lines dividing the plane into regions

Let's consider lines dividing the plane into regions:

- 1 line divides it into 2 regions.
- 2 lines into 4 regions.
- 3 lines into 7 regions...



(It is assumed that each pair of lines intersects, and no three lines intersect in the same point.)


When there are n-1 lines and we add one more line. How does it change the number of regions?

Recurrence

Lines dividing the plane

Towers of Hanoi

Lines dividing the plane into regions



When there are n-1 lines and we add one more line. How does it change the number of regions?

$$R(n) = R(n-1) + n$$

Full recurrence:

$$R(0) = 1$$

 $R(n) = R(n-1) + n$ (when $n > 0$)

Okay, nice, but can we find a *closed-form* expression for it?

Recurrence

Lines dividing the plane

Towers of Hanoi

Lines dividing the plane into regions

Recurrence

Lines dividing the plane

Towers of Hanoi

Fibonacci staircase

Recurrence:

$$R(0) = 1$$

 $R(n) = R(n-1) + n$ (when $n > 0$)

Answer for the closed-form expression:

$$R(n) = \frac{n(n+1)}{2} + 1.$$

It is almost identical to the Sum(n) recurrence we considered earlier, differing only in the way it is initialized.



http://www.mathsisfun.com/games/towerofhanoi.html


Recurrence

Lines dividing the plane

Towers of Hanoi

Our recursive algorithm to move a tower of height n from #1 to #3:

- 1. Move an (n-1)-tower from #1 to #2.
- 2. Move an 1-tower from #1 to #3.
- 3. Move an (n-1)-tower from #2 to #3.



Recurrence


Lines dividing the plane

Towers of Hanoi

Our recursive algorithm to move a tower of height n from #1 to #3:

- 1. Move an (n-1)-tower from #1 to #2.
- 2. Move an 1-tower from #1 to #3.
- 3. Move an (n-1)-tower from #2 to #3.

There is a way to find a recurrent formula for T_n , the total number of steps to move the tower from the peg 1 to the peg 3.




Recurrence

Lines dividing the plane

Towers of Hanoi

 T_n , the time to move a tower of height n:

$$\begin{split} T_1 &= 1 \\ T_n &= T_{n-1} + 1 + T_{n-1} \qquad (\forall n > 1) \end{split}$$



Recurrence

Lines dividing the plane


Towers of Hanoi

 T_n , the time to move a tower of height n:

$$T_1 = 1$$

 $T_n = T_{n-1} + 1 + T_{n-1}$ $(\forall n > 1)$

There is a proof by induction that this time is optimal for any algorithm.



Recurrence

Lines dividing the plane

Towers of Hanoi

Recurrence

Lines dividing the plane

Towers of Hanoi

Fibonacci staircase

$$T_1 = 1$$

 $T_n = 2T_{n-1} + 1$ $(\forall n > 1)$

Our goal is to find a closed form expression for T_n as a function of n, without any recurrence.

Before we get a closed form formula for T_n , what are the numbers?

$$T_1 = 1$$

 $T_n = 2T_{n-1} + 1$ $(\forall n > 1)$

We can compute a list like this:

$$T_1 = 1$$

 $T_2 = 3$
 $T_3 = 7$
 $T_4 = 15$
 $T_5 = 31$
 $T_6 = 63$
...

Recurrence

Lines dividing the plane

Towers of Hanoi

$$T_1 = 1$$

 $T_n = 2T_{n-1} + 1$ $(\forall n > 1)$

$$T_1 = 1$$

 $T_2 = 3$
 $T_3 = 7$
 $T_4 = 15$
 $T_5 = 31$
 $T_6 = 63$
...

Guess and verify method... Let's try $T_n = 2^n - 1$?

Recurrence

Lines dividing the plane

Towers of Hanoi

Recurrence

Lines dividing the plane

Towers of Hanoi

Fibonacci staircase

$$T_1 = 1$$

 $T_n = 2T_{n-1} + 1$ $(\forall n > 1)$

Guess and verify method... Let's try $T_n = 2^n - 1$? We can show by induction that this formula is correct.

The base case, n = 1:

$$T_1 = 2^1 - 1 = 1.$$

Ok, the base case is true.

$$T_1 = 1$$

 $T_n = 2T_{n-1} + 1$ $(\forall n > 1)$

We want to prove the closed form formula $T_n = 2^n - 1$.

The inductive step, n > 1:

Assume that $T_n = 2^n - 1$, and show that then $T_{n+1} = 2^{n+1} - 1$.

Proof. From the recurrence:

$$T_{n+1} = 2T_n + 1$$

By the inductive hypothesis:

$$2T_n + 1 = 2(2^n - 1) + 1 = 2^{n+1} - 2 + 1 = 2^{n+1} - 1$$

Recurrence

Lines dividing the plane

Towers of Hanoi

If it is difficult to guess the closed form expression for the recurrence, there is another technique:

$$T_1 = 1$$
$$T_n = 2T_{n-1} + 1$$

Recurrence

Lines dividing the plane

Towers of Hanoi

If it is difficult to guess the closed form expression for the recurrence, there is another technique:

$$T_1 = 1$$

 $T_n = 2T_{n-1} + 1$
 $= 2(2T_{n-2} + 1) + 1$

Recurrence

Lines dividing the plane

Towers of Hanoi

If it is difficult to guess the closed form expression for the recurrence, there is another technique:

$$T_1 = 1$$

$$T_n = 2T_{n-1} + 1$$

$$= 2(2T_{n-2} + 1) + 1$$

$$= 2^2T_{n-2} + 2 + 1$$

Recurrence

Lines dividing the plane

Towers of Hanoi

If it is difficult to guess the closed form expression for the recurrence, there is another technique:

$$T_{1} = 1$$

$$T_{n} = 2T_{n-1} + 1$$

$$= 2(2T_{n-2} + 1) + 1$$

$$= 2^{2}T_{n-2} + 2 + 1$$

$$= 2^{2}(2T_{n-3} + 1) + 2 + 1$$

$$= 2^{3}T_{n-3} + 2^{2} + 2 + 1$$
...

Recurrence

Lines dividing the plane

Towers of Hanoi

If it is difficult to guess the closed form expression for the recurrence, there is another technique:

```
\begin{split} T_1 &= 1 \\ T_n &= 2T_{n-1} + 1 \\ &= 2(2T_{n-2} + 1) + 1 \\ &= 2^2T_{n-2} + 2 + 1 \\ &= 2^2(2T_{n-3} + 1) + 2 + 1 \\ &= 2^3T_{n-3} + 2^2 + 2 + 1 = \dots = 2^kT(n-k) + 2^{k-1} + \dots + 2 + 1 \\ &\dots \quad \text{(can expand until } k = n-1 \text{ and } T(1) = 1) \end{split}
```

Recurrence

Lines dividing the plane

Towers of Hanoi

If it is difficult to guess the closed form expression for the recurrence, there is another technique:

$$\begin{split} T_1 &= 1 \\ T_n &= 2T_{n-1} + 1 \\ &= 2(2T_{n-2} + 1) + 1 \\ &= 2^2T_{n-2} + 2 + 1 \\ &= 2^2(2T_{n-3} + 1) + 2 + 1 \\ &= 2^3T_{n-3} + 2^2 + 2 + 1 = \dots = 2^kT(n-k) + 2^{k-1} + \dots + 2 + 1 \\ \dots & (\text{can expand until } k = n - 1 \text{ and } T(1) = 1) \\ &= 2^{n-1}\underbrace{T(n - (n-1))}_{=1} + \dots + 4 + 2 + 1 \end{split}$$

Recurrence

Lines dividing the plane

Towers of Hanoi

If it is difficult to guess the closed form expression for the recurrence, there is another technique:

$$T_{1} = 1$$

$$T_{n} = 2T_{n-1} + 1$$

$$= 2(2T_{n-2} + 1) + 1$$

$$= 2^{2}T_{n-2} + 2 + 1$$

$$= 2^{2}(2T_{n-3} + 1) + 2 + 1$$

$$= 2^{3}T_{n-3} + 2^{2} + 2 + 1 = \dots = 2^{k}T(n-k) + 2^{k-1} + \dots + 2 + 1$$

$$\dots \quad (\text{can expand until } k = n - 1 \text{ and } T(1) = 1)$$

$$= 2^{n-1}\underbrace{T(n - (n-1))}_{=1} + \dots + 4 + 2 + 1$$

$$= 2^{n-1} + \dots + 4 + 2 + 1 = \sum_{k=0}^{n-1} 2^{k} = 1$$

Recurrence

Lines dividing the plane

Towers of Hanoi

If it is difficult to guess the closed form expression for the recurrence, there is another technique:

$$T_{1} = 1$$

$$T_{n} = 2T_{n-1} + 1$$

$$= 2(2T_{n-2} + 1) + 1$$

$$= 2^{2}T_{n-2} + 2 + 1$$

$$= 2^{2}(2T_{n-3} + 1) + 2 + 1$$

$$= 2^{3}T_{n-3} + 2^{2} + 2 + 1 = \dots = 2^{k}T(n-k) + 2^{k-1} + \dots + 2 + 1$$

$$\dots \quad (\text{can expand until } k = n - 1 \text{ and } T(1) = 1)$$

$$= 2^{n-1}\underbrace{T(n - (n-1))}_{=1} + \dots + 4 + 2 + 1$$

$$= 2^{n-1} + \dots + 4 + 2 + 1 = \sum_{k=0}^{n-1} 2^{k} = \frac{1 - 2^{n}}{1 - 2} = 2^{n} - 1.$$

Recurrence

Lines dividing the plane

Towers of Hanoi

Recurrence

Lines dividing the plane

Towers of Hanoi

Fibonacci staircase

Why is it useful to know that the recurrence

$$T_1 = 1$$

 $T_n = 2T_{n-1} + 1$ $(\forall n > 1)$

is equivalent to the closed form formula $T_n = 2^n - 1$?

The 7-disk puzzle will require $T_7 = 2^7 - 1 = 127$ moves to complete.

And the 100-disk puzzle will require

$$T_{100} = 2^{100} - 1 = 1267650600228229401496703205375$$
 moves.

Fibonacci staircase



plane

Towers of Hanoi

Fibonacci staircase



A staircase has *n* steps.

You walk up by taking 1 or 2 steps at a time.

Question: In how many different ways can you walk up the stairs?