

Infinity. Cardinality.
Pairing function. Diagonalization.

Infinity?

We know that the set of natural numbers, \mathbb{N} , is infinite, so, definitely, there are sets with infinitely many elements.

How is it possible to construct such sets?

Let's define an operation

$$A^+ = A \cup \{A\}$$

We start with \emptyset and apply this operation:

$$\emptyset = \emptyset$$

$$\emptyset^+ = \{\emptyset\}$$

$$(\emptyset^+)^+ = \{\emptyset, \{\emptyset\}\}$$

$$((\emptyset^+)^+)^+ = \{\emptyset, \{\emptyset\}, \{\emptyset, \{\emptyset\}\}\}$$

...

Infinity

Infinite sets

Countable sets

Hilbert's Hotel

Ordered pairs

Power set.
Diagonalization.

Schröder-Bernstein
Theorem

Infinity?

We know that the set of natural numbers, \mathbb{N} , is infinite, so, definitely, there are sets with infinitely many elements.

How is it possible to construct such sets?

Let's define an operation

$$A^+ = A \cup \{A\}$$

We start with \emptyset and apply this operation:

$$'0' = \emptyset = \emptyset$$

$$'1' = \emptyset^+ = \{\emptyset\} = \{'0'\}$$

$$'2' = (\emptyset^+)^+ = \{\emptyset, \{\emptyset\}\} = \{'0', '1'\}$$

$$'3' = ((\emptyset^+)^+)^+ = \{\emptyset, \{\emptyset\}, \{\emptyset, \{\emptyset\}\}\} = \{'0', '1', '2'\}$$

...

This is von Neumann's construction of natural numbers.

Infinity

Infinite sets

Countable sets

Hilbert's Hotel

Ordered pairs

Power set.
Diagonalization.

Schröder-Bernstein
Theorem

Infinity

Suppose that you have infinitely many one dollar bills (numbered 1, 3, 5, ...) and you come upon the Devil, who is willing to pay two dollars for each of your one-dollar bills.



The Devil is very particular, however, about the order in which the bills are exchanged. The contract stipulates that in each sub-transaction he buys from you your lowest-numbered bill and pays you with higher-numbered bills.

First sub-transaction takes $1/2$ hour, then $1/4$ hour, $1/8$, and so on, so that after one hour the entire exchange will be complete.

Infinity

Infinite sets

Countable sets

Hilbert's Hotel

Ordered pairs

Power set.
Diagonalization.

Schröder-Bernstein
Theorem

Infinite sets

Infinity

Infinite sets

Countable sets

Hilbert's Hotel

Ordered pairs

Power set.
Diagonalization.

Schröder-Bernstein
Theorem

Consider four sets:

$$\mathbb{N} = \{0, 1, 2, 3, 4, \dots\}$$

$$Even_N = \{0, 2, 4, 6, 8, \dots\}$$

$$Odd_N = \{1, 3, 5, 7, 9, \dots\}$$

$$\mathbb{Z}^- = \{-1, -2, -3, -4, \dots\}$$

Can we compare their cardinalities?

Infinite sets

Infinity

Infinite sets

Countable sets

Hilbert's Hotel

Ordered pairs

Power set.
Diagonalization.

Schröder-Bernstein
Theorem

Consider four sets:

$$\mathbb{N} = \{0, 1, 2, 3, 4, \dots\}$$

$$Even_N = \{0, 2, 4, 6, 8, \dots\}$$

$$Odd_N = \{1, 3, 5, 7, 9, \dots\}$$

$$\mathbb{Z}^- = \{-1, -2, -3, -4, \dots\}$$

Can we compare their cardinalities?

We need a definition for the cardinality of an infinite set.

Cardinality of an infinite set

Def. The sets A and B have the same cardinality if and only if there is a bijection from A to B .

When A and B have the same cardinality, we write $|A| = |B|$.

$$\mathbb{N} = \{0, 1, 2, 3, 4, \dots\}$$

$$Even_N = \{0, 2, 4, 6, 8, \dots\}$$

$$Odd_N = \{1, 3, 5, 7, 9, \dots\}$$

$$\mathbb{Z}^- = \{-1, -2, -3, -4, \dots\}$$

Infinity

Infinite sets

Countable sets

Hilbert's Hotel

Ordered pairs

Power set.
Diagonalization.

Schröder-Bernstein
Theorem

Cardinality of an infinite set

Infinity

Infinite sets

Countable sets

Hilbert's Hotel

Ordered pairs

Power set.
Diagonalization.

Schröder-Bernstein
Theorem

$$\mathbb{N} = \{0, 1, 2, 3, 4, \dots\}$$
$$Even_N = \{0, 2, 4, 6, 8, \dots\}$$

Find a bijection

$$f : \mathbb{N} \rightarrow Even_N$$

0● 1● 2● 3● ...

●0 ●2 ●4 ●6 ...

Cardinality of an infinite set

Infinity

Infinite sets

Countable sets

Hilbert's Hotel

Ordered pairs


Power set.
Diagonalization.

Schröder-Bernstein
Theorem

$$\mathbb{N} = \{0, 1, 2, 3, 4, \dots\}$$
$$Even_N = \{0, 2, 4, 6, 8, \dots\}$$

Find a bijection

$$f : \mathbb{N} \rightarrow Even_N$$



$$f(x) = 2x$$

Cardinality of an infinite set

Infinity

Infinite sets

Countable sets

Hilbert's Hotel

Ordered pairs


Power set.
Diagonalization.

Schröder-Bernstein
Theorem

$$\mathbb{N} = \{0, 1, 2, 3, 4, \dots\}$$
$$Even_N = \{0, 2, 4, 6, 8, \dots\}$$

Alternatively

$$f : \mathbb{N} \rightarrow Even_N$$



Cardinality of an infinite set

Infinity

Infinite sets

Countable sets

Hilbert's Hotel

Ordered pairs

Power set.
Diagonalization.

Schröder-Bernstein
Theorem

$$\mathbb{N} = \{0, 1, 2, 3, 4, \dots\}$$
$$Odd_N = \{1, 3, 5, 7, 9, \dots\}$$

Find a bijection

$$f : \mathbb{N} \rightarrow Odd_N$$

0● 1● 2● 3● ...

●1 ●3 ●5 ●7 ...

Cardinality of an infinite set

Infinity

Infinite sets

Countable sets

Hilbert's Hotel

Ordered pairs


Power set.
Diagonalization.

Schröder-Bernstein
Theorem

$$\mathbb{N} = \{0, 1, 2, 3, 4, \dots\}$$
$$Odd_N = \{1, 3, 5, 7, 9, \dots\}$$

Find a bijection

$$f : \mathbb{N} \rightarrow Odd_N$$



$$f(x) = 2x + 1$$

Cardinality of an infinite set

Infinity

Infinite sets

Countable sets

Hilbert's Hotel

Ordered pairs

Power set.
Diagonalization.

Schröder-Bernstein
Theorem

$$\mathbb{N} = \{0, 1, 2, 3, 4, \dots\}$$

$$\mathbb{Z}^- = \{-1, -2, -3, -4, -5, \dots\}$$

Find a bijection

$$f : \mathbb{N} \rightarrow \mathbb{Z}^-$$

0 ● 1 ● 2 ● 3 ● ...

●-1 ●-2 ●-3 ●-4 ...

Cardinality of an infinite set

Infinity

Infinite sets

Countable sets

Hilbert's Hotel

Ordered pairs


Power set.
Diagonalization.

Schröder-Bernstein
Theorem

$$\mathbb{N} = \{0, 1, 2, 3, 4, \dots\}$$
$$\mathbb{Z}^- = \{-1, -2, -3, -4, -5, \dots\}$$

Find a bijection

$$f : \mathbb{N} \rightarrow \mathbb{Z}^-$$



$$f(x) = -x - 1$$

Cardinality of an infinite set

Infinity

Infinite sets

Countable sets

Hilbert's Hotel

Ordered pairs

Power set.
Diagonalization.

Schröder-Bernstein
Theorem

Therefore, all these sets have the same cardinality

$$|\mathbb{N}| = |Even_N| = |Odd_N| = |\mathbb{Z}^-|$$

Countable sets

- Infinity
- Infinite sets
- Countable sets
- Hilbert's Hotel
- Ordered pairs
- Power set.
Diagonalization.
- Schröder-Bernstein
Theorem

Therefore, all these sets have the same cardinality

$$|\mathbb{N}| = |Even_N| = |Odd_N| = |\mathbb{Z}^-|$$

Def. A set S is called *countable* if $|S| = |\mathbb{N}|$ or if S is a finite set.

Countable sets

- Infinity
- Infinite sets
- Countable sets
- Hilbert's Hotel
- Ordered pairs
- Power set.
Diagonalization.
- Schröder-Bernstein
Theorem

Since \mathbb{N} is an infinite set, the cardinality $|\mathbb{N}|$ is greater than any natural number. We need a way to denote the cardinality of this set.


The following symbol is used

$$|\mathbb{N}| = \aleph_0$$

It reads as “aleph naught”, “aleph null”, “aleph zero”.

All infinite countable sets have the same cardinality \aleph_0 .

Hilbert's Hotel




Imagine a hotel with a countably infinite number of rooms.

Each room is occupied by a guest.

Question: Can it accomodate one more guest?

- Infinity
- Infinite sets
- Countable sets
- Hilbert's Hotel
- Ordered pairs
- Power set.
- Diagonalization.
- Schröder-Bernstein Theorem

Hilbert's Hotel




- Infinity
- Infinite sets
- Countable sets
- Hilbert's Hotel
- Ordered pairs
- Power set.
- Diagonalization.
- Schröder-Bernstein Theorem

There is a bijection between $\{x\} \cup \mathbb{N}$ (guests) and \mathbb{N} (rooms)

$x \bullet \quad 0 \bullet \quad 1 \bullet \quad 2 \bullet \quad 3 \bullet \quad \dots$

$\bullet 0 \quad \bullet 1 \quad \bullet 2 \quad \bullet 3 \quad \dots$

Hilbert's Hotel



Infinity

Infinite sets

Countable sets

Hilbert's Hotel

Ordered pairs

Power set.

Diagonalization.

Schröder-Bernstein
Theorem

There is a bijection between $\{x\} \cup \mathbb{N}$ (guests) and \mathbb{N} (rooms)



More complex cases

- Infinity
- Infinite sets
- Countable sets
- Hilbert's Hotel
- Ordered pairs
- Power set.
- Diagonalization.
- Schröder-Bernstein Theorem

We want to prove that $B = \mathbb{N} \times \{T, F\}$ is countable.

Can we find a bijection between \mathbb{N} and $B = \mathbb{N} \times \{T, F\}$?

$$\mathbb{N} = \{0, 1, 2, 3, 4, 5, \dots\}$$


$$B = \{(0, T), (1, T), (2, T), \dots (0, F), (1, F), (2, F), \dots\}$$

More complex cases

Can we find a bijection between \mathbb{N} and $B = \mathbb{N} \times \{T, F\}$?

$$\mathbb{N} = \{0, 1, 2, 3, 4, 5, \dots\}$$

$$B = \{(0, T), (1, T), (2, T), \dots (0, F), (1, F), (2, F), \dots\}$$



$$(0, T), (0, F), (1, T), (1, F), (2, T), (2, F), \dots$$

Infinity
Infinite sets
Countable sets
Hilbert's Hotel
Ordered pairs
Power set.
Diagonalization.
Schröder-Bernstein
Theorem

More complex cases

Infinity
Infinite sets
Countable sets
Hilbert's Hotel
Ordered pairs
Power set.
Diagonalization.
Schröder-Bernstein
Theorem

Similarly, there is a bijection between \mathbb{N} and \mathbb{Z}

$$\mathbb{N} = \{0, 1, 2, 3, \dots\}$$

$$\mathbb{Z} = \{\dots - 3, -2, -1, 0, 1, 2, 3, \dots\}$$

We just rearrange the order of integers:

$$0, 1, -1, 2, -2, 3, -3, \dots$$

In general, if there is a way *to list* the elements of a given set in linear order, then it is *countable* (i.e. there is a bijection between this set and \mathbb{N}).

More complex cases

Find a bijection $h : A \rightarrow B$, where

$$A = \mathbb{N} \times \{T, F\}$$

$$B = \mathbb{Z}$$

Infinity

Infinite sets

Countable sets

Hilbert's Hotel

Ordered pairs

Power set.

Diagonalization.

Schröder-Bernstein
Theorem

More complex cases

Find a bijection $h : A \rightarrow B$, where

$$A = \mathbb{N} \times \{T, F\}$$

$$B = \mathbb{Z}$$

A and B are countable, and we know how to construct the following two bijections

$$f : \mathbb{N} \rightarrow A$$

$$g : \mathbb{N} \rightarrow B$$

Since f is a bijection, there exist an inverse function $f^{-1} : A \rightarrow \mathbb{N}$, which is a bijection too, and we can find it, so

$$h(x) = g(f^{-1}(x))$$

Infinity

Infinite sets

Countable sets

Hilbert's Hotel

Ordered pairs

Power set.

Diagonalization.

Schröder-Bernstein
Theorem

$$A \times \mathbb{N}$$

Infinity
Infinite sets
Countable sets
Hilbert's Hotel
Ordered pairs
Power set.
Diagonalization.
Schröder-Bernstein
Theorem

We have shown that \mathbb{Z} is countable, $\mathbb{N} \times \{T, F\}$ is countable.


Similarly, it's not hard to show that for any *finite* set A , its Cartesian products

$A \times \mathbb{N}$ and $\mathbb{N} \times A$ are countable.

$\mathbb{N} \times A$ and $A \times \mathbb{N}$ when A is finite

Similarly, it's not hard to show that for any *finite* set A , its Cartesian products

$A \times \mathbb{N}$ and $\mathbb{N} \times A$ are countable.



- Infinity
- Infinite sets
- Countable sets
- Hilbert's Hotel
- Ordered pairs
- Power set.
- Diagonalization.
- Schröder-Bernstein Theorem

Is the set $\mathbb{N} \times \mathbb{N}$ countable?

Can we find a bijection $\mathbb{N} \rightarrow \mathbb{N} \times \mathbb{N}$? If yes, then the set of ordered pairs of natural numbers, $\mathbb{N} \times \mathbb{N}$, is a countable set.

| | | | | | |
|----------|----------|----------|----------|----------|---------|
| \vdots | \vdots | \vdots | \vdots | \vdots | |
| (0, 3) | (1, 3) | (2, 3) | (3, 3) | (4, 3) | \dots |
| (0, 2) | (1, 2) | (2, 2) | (3, 2) | (4, 2) | \dots |
| (0, 1) | (1, 1) | (2, 1) | (3, 1) | (4, 1) | \dots |
| (0, 0) | (1, 0) | (2, 0) | (3, 0) | (4, 0) | \dots |

Infinity

Infinite sets

Countable sets

Hilbert's Hotel

Ordered pairs


Power set.

Diagonalization.

Schröder-Bernstein
Theorem

Is the set $\mathbb{N} \times \mathbb{N}$ countable?


Can we find a bijection $\mathbb{N} \rightarrow \mathbb{N} \times \mathbb{N}$? If yes, then the set of ordered pairs of natural numbers, $\mathbb{N} \times \mathbb{N}$, is a countable set.



Infinity
Infinite sets
Countable sets
Hilbert's Hotel
Ordered pairs
Power set.
Diagonalization.
Schröder-Bernstein
Theorem

Pairing function $\mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$

$$P(x, y) = \frac{1}{2}(x + y)(x + y + 1) + y$$



Infinity
Infinite sets
Countable sets
Hilbert's Hotel
Ordered pairs
Power set.
Diagonalization.
Schröder-Bernstein
Theorem

The set of rational numbers, \mathbb{Q}

- Infinity
- Infinite sets
- Countable sets
- Hilbert's Hotel
- Ordered pairs
- Power set.
Diagonalization.
- Schröder-Bernstein
Theorem

We can define the set of rational numbers as the set of all quotients p/q such that $p \in \mathbb{Z}$ and $q \in \mathbb{Z}^+$:

$$\mathbb{Q} = \left\{ \frac{p}{q} \mid p \in \mathbb{Z} \wedge q \in \mathbb{Z}^+ \right\}$$

We can prove that \mathbb{Q} is countable. The argument is similar to the proof for $\mathbb{N} \times \mathbb{N}$.

Power set

Infinity
Infinite sets
Countable sets
Hilbert's Hotel
Ordered pairs
Power set.
Diagonalization.
Schröder-Bernstein
Theorem

Is the power set $\mathcal{P}(\mathbb{N})$ countable?

Power set

Infinity

Infinite sets

Countable sets

Hilbert's Hotel

Ordered pairs

Power set.

Diagonalization.

Schröder-Bernstein
Theorem

Theorem. The power set $\mathcal{P}(\mathbb{N})$ is not countable.

Power set

Proof. (by contradiction)

Assume that $\mathcal{P}(\mathbb{N})$ is countable, so all subsets of \mathbb{N} can be listed:

$$A_0, A_1, A_2, \dots$$

We know that subsets can be encoded by strings of 1s and 0s.

| Subset | 0 | 1 | 2 | 3 | 4 | 5 | ... |
|--------|---|---|---|---|---|---|-----|
| A_0 | 0 | 0 | 0 | 1 | 0 | 0 | ... |
| A_1 | 1 | 1 | 1 | 0 | 0 | 1 | ... |
| A_2 | 1 | 1 | 1 | 1 | 1 | 1 | ... |
| A_3 | 0 | 0 | 0 | 0 | 0 | 1 | ... |
| A_4 | 1 | 0 | 0 | 0 | 0 | 1 | ... |
| A_5 | 1 | 1 | 0 | 0 | 1 | 1 | ... |

Now, we want to construct a counter-example subset $C \subseteq \mathbb{N}$ that is different from each A_i .

Infinity
Infinite sets
Countable sets
Hilbert's Hotel
Ordered pairs
Power set.
Diagonalization.
Schröder-Bernstein
Theorem

Power set

| Subset | 0 | 1 | 2 | 3 | 4 | 5 | ... |
|--------|---|---|---|---|---|---|-----|
| A_0 | 0 | 0 | 0 | 1 | 0 | 0 | ... |
| A_1 | 1 | 1 | 1 | 0 | 0 | 1 | ... |
| A_2 | 1 | 1 | 0 | 1 | 1 | 1 | ... |
| A_3 | 0 | 0 | 0 | 1 | 0 | 1 | ... |
| A_4 | 1 | 0 | 0 | 0 | 1 | 1 | ... |
| A_5 | 1 | 1 | 0 | 0 | 1 | 0 | ... |
| ... | | | | | | | |
| C | 1 | 0 | 0 | 1 | 1 | 0 | ... |

We construct a counter-example set C that is different from each subset A_i . How can we do it?

For all $i = 0, 1, 2, 3, \dots$: Whenever $i \in A_i$, we choose $i \notin C$, and vice versa, when $i \notin A_i$, we choose $i \in C$. Thus, by construction, C is different from each A_i . Effectively, the set C inverts the diagonal.

Infinity

Infinite sets

Countable sets

Hilbert's Hotel

Ordered pairs

Power set.
Diagonalization.

Schröder-Bernstein
Theorem

Power set

Since $C \neq A_i$ for all i , and C is obviously a subset of \mathbb{N} by construction, the list of subsets A_i does not contain all subsets of \mathbb{N} (it does not contain C , for example), therefore, our assumption was incorrect: the subsets of \mathbb{N} are not countable.

That is, *the power set $\mathcal{P}(\mathbb{N})$ is uncountable.* □

This proof strategy is called diagonalization.

Similarly, we can show that the *unit interval* $0 \leq x \leq 1$ of real numbers is uncountable. (Also, see Rosen's book for the proof). And because you can make a bijection between this interval, $[0, 1]$, and \mathbb{R} , the set of all real number is uncountable.

- Infinity
- Infinite sets
- Countable sets
- Hilbert's Hotel
- Ordered pairs
- Power set.
- Diagonalization.
- Schröder-Bernstein Theorem

More results about cardinality

Theorem. If A and B are countable sets, then their union $A \cup B$ is also countable.

Proof. Without loss of generality, we can assume that A and B are disjoint. (If they are not, we continue the proof with A and $B \setminus A$)

If at least one of the sets is finite, we first list this set, then the other set.

Otherwise, if both are infinite countable sets, we list both sets by alternating elements:

$$a_0, b_0, a_1, b_1, a_2, b_2, \dots$$

where $a_i \in A$ and $b_i \in B$.



- Infinity
- Infinite sets
- Countable sets
- Hilbert's Hotel
- Ordered pairs
- Power set.
- Diagonalization.
- Schröder-Bernstein Theorem

Cardinality, one-to-one and onto

Infinity
Infinite sets
Countable sets
Hilbert's Hotel
Ordered pairs
Power set.
Diagonalization.
Schröder-Bernstein
Theorem

Mapping rules

If there is a *one-to-one* function $f : A \rightarrow B$ then

$$|A| \leq |B|.$$

If there is an *onto* function $g : A \rightarrow B$ then

$$|A| \geq |B|.$$

If there is a *bijection* $h : A \rightarrow B$ then

$$|A| = |B|.$$

Schröder-Bernstein Theorem

Infinity

Infinite sets

Countable sets

Hilbert's Hotel

Ordered pairs

Power set.

Diagonalization.

Schröder-Bernstein
Theorem

Theorem (Schröder-Bernstein). Given two sets A and B , if there exist one-to-one functions $f : A \rightarrow B$ and $g : B \rightarrow A$, then there is a bijection between A and B .

In other words, to prove existence of a bijection, it's enough to prove existence of two one-to-one functions:

Once you have found a one-to-one function $f : A \rightarrow B$, instead of proving that f is onto, you can prove that there exists another one-to-one function that maps B to A .