

Modular Arithmetic

Previously, we defined

GCD is a linear combination

Relative primes

Fundamental theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's Algorithm

Def (Divisibility). We say that a *divides* b if there is an integer k such that

$$b = a \cdot k.$$

We write $a \mid b$ if a divides b . Otherwise, we write $a \nmid b$.

Theorem (The Division Algorithm). Let a be an integer and d a positive integer. Then there are *unique* integers q and r , such that $0 \leq r < d$ and

$$a = dq + r.$$

GCD is a linear combination

GCD is a linear
combination

Relative primes

Fundamental
theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's
Algorithm

Def (GCD).

Theorem (Bezout's Theorem). If a and b are positive integers, then there exist integers s and t such that

$$\gcd(a, b) = sa + tb.$$

Exmaple: $\gcd(52, 44) = 4$

$$6 \cdot 52 + (-7) \cdot 44 = 4$$

So called Extended Euclid's algorithm constructs such s and t , and so proves the theorem. The algorithm is described in the last section of this lecture.

Relative primes (co-primes)

GCD is a linear combination

Relative primes

Fundamental theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's Algorithm

Def (Prime numbers).

Def (Relative primes). a and b are *relative primes* (or co-primes) if

$$\gcd(a, b) = 1.$$

By Bezout's theorem, a and b are co-primes if and only if there exist s and t such that

$$sa + tb = 1$$

Factorization of positive integers

GCD is a linear combination

Relative primes

Fundamental theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's Algorithm

Theorem (Fundamental theorem of arithmetic). Every positive integer n can be written in a unique way as a product of primes

$$n = p_1 \cdot p_2 \cdot \dots \cdot p_j \quad (p_1 \leq p_2 \leq \dots \leq p_j)$$

This product is called prime factorization.

See Lehman and Leighton (p. 67) for the proof.

Modular arithmetic

... -2 -1 0 1 2 3 4 5 6 7 8...

What if instead of integers, we deal with
a finite set of periodically repeating integers?

... 5 6 → 0 1 2 3 4 5 6 → 0 1...

GCD is a linear
combination

Relative primes

Fundamental
theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's
Algorithm

Modular arithmetic

GCD is a linear combination

Relative primes

Fundamental theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's Algorithm

$\dots -2 \quad -1 \quad \quad 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \quad 6 \quad \quad 7 \quad 8 \dots$

What if instead of integers, we deal with
a finite set of periodically repeating integers?

$\dots 5 \quad 6 \rightarrow 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \quad 6 \rightarrow 0 \quad 1 \dots$

For example, the days of the week behave in this way.

Mon, Tue, Wed, Thr, Fri, Sat, Sun,

are followed again by Mon, Tue, and so on.

Modular arithmetic

...5 6 \rightarrow 0 1 2 3 4 5 6 \rightarrow 0 1...

We want to add, subtract, multiply, and, hopefully, divide such special “integers” ...

$$4 + 4 \text{ is } 1$$

$$2 - 3 \text{ is } 6$$

$$14 \cdot 5 \text{ is } 0$$

$$-7 \text{ is } 0 \text{ is } 7 \text{ is } 14 \text{ is } 21 \dots$$

First, we need to rigorously define, which integers can be called “equal” in such modular arithmetic. We will call them congruent.

GCD is a linear combination

Relative primes

Fundamental theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's Algorithm

Congruence

Def. For a positive integer n , a is *congruent* to b modulo n if

$$n \mid (a - b).$$

This is denoted

$$a \equiv b \pmod{n}.$$

Example:

$$8 \equiv 1 \pmod{7}$$

$$15 \equiv 1 \pmod{7}$$

$$8 \equiv 15 \pmod{7}$$

because

$$7 \mid \underbrace{(8-1)}_{=7}, \quad 7 \mid \underbrace{(15-1)}_{=14}, \quad 7 \mid \underbrace{(15-8)}_{=7}.$$

GCD is a linear combination

Relative primes

Fundamental theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's Algorithm

Congruence

GCD is a linear combination

Relative primes

Fundamental theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's Algorithm

Lemma. If $a \equiv b \pmod{n}$, then exists $k \in \mathbb{Z}$ s.t. $a = b + kn$.

Lemma. Two numbers are congruent modulo n if and only if they have the same remainder when divided by n .

$$a \equiv b \pmod{n} \quad \text{if and only if} \quad a \bmod n = b \bmod n.$$

Proof: By the division algorithm,

$$a = q_1n + r_1, \quad b = q_2n + r_2.$$

$$a - b = (q_1 - q_2)n + (r_1 - r_2)$$

“ \Rightarrow ”: If $a \equiv b \pmod{n}$ then $n \mid (a - b)$. So $r_1 - r_2 = 0$, the remainders are equal.

“ \Leftarrow ”: If $r_1 = r_2$, then $n \mid (a - b)$, so $a \equiv b \pmod{n}$. □

Congruence

-3 0 3 6
 -2 1 4 7
 -1 2 5 8

$x \bmod 3$ 0 1 2 0 1 2 0 1 2 0 1 2

Integers are divided into 3 congruence classes:

..., -3, 0, 3, 6, 9, 12, ... are congruent modulo 3.

..., -2, 1, 4, 7, 10, 13, ... are congruent modulo 3.

..., -1, 2, 5, 8, 11, 14, ... are congruent modulo 3.

By the way, all Mondays, all Tuesdays, all Wednesdays, etc. are congruence classes too.

GCD is a linear combination
 Relative primes
 Fundamental theorem of arithmetic
 Modular arithmetic
 Congruence
 Modular arithmetic
 Multiplicative inverse
 Extended Euclid's Algorithm

Modular arithmetic

Addition, subtraction, and multiplication preserve congruence.

Theorem. if $a \equiv b \pmod{n}$ and $c \equiv d \pmod{n}$, then

$$a + c \equiv b + d \pmod{n}.$$

Theorem. if $a \equiv b \pmod{n}$ and $c \equiv d \pmod{n}$, then

$$ac \equiv bd \pmod{n}.$$

Proof.

Exist $x, y \in \mathbb{Z}$ such that $a - b = xn$ and $c - d = yn$.

$$ac - bd = (b + xn)(d + yn) - bd = n(xd + by + xny)$$

Thus $ac \equiv bd \pmod{n}$.



GCD is a linear combination

Relative primes

Fundamental theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's Algorithm

Modular arithmetic

Addition, subtraction, and multiplication preserve congruence.
What does it mean practically?

If we have to find x such that

$$12^2 \cdot (-11) + 80 \equiv x \pmod{5}$$

We know that

$$\begin{aligned} 12 &\equiv 2 \pmod{5}, \\ -11 &\equiv -1 \pmod{5}, \\ 80 &\equiv 0 \pmod{5} \end{aligned}$$

Therefore, we are free to substitute 12 with 2, -11 with -1 , and 80 with 0:

$$12^2 \cdot (-11) + 80 \equiv 2^2 \cdot (-1) + 0 \equiv 2 \cdot 2 \cdot (-1) \equiv -4 \equiv 1 \pmod{5}.$$

GCD is a linear combination

Relative primes

Fundamental theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's Algorithm

Multiplicative inverse

What about division?

Theorem. if a and n are relative primes, i.e. $\gcd(a, n) = 1$, then exists integer a^{-1} called *multiplicative inverse*, such that

$$aa^{-1} \equiv 1 \pmod{n}$$

Proof.

Exist s and t , such that $sa + tn = 1$. Therefore,

$$sa - 1 = tn$$

$$sa \equiv 1 \pmod{n}$$

Therefore, $a^{-1} = s$.



GCD is a linear combination

Relative primes

Fundamental theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's Algorithm

Multiplicative inverse

GCD is a linear combination

Relative primes

Fundamental theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's Algorithm

Corollary. If a and n are relative primes, then there exists a *unique* multiplicative inverse $a^{-1} \in \{1, 2, \dots, n-1\}$ such that

$$aa^{-1} \equiv 1 \pmod{n}.$$

Ok, uniqueness is great, but we need a procedure for finding multiplicative inverses.

Multiplicative inverse

Find inverse of 101 modulo 4620, that is x such that

$$101 \cdot x \equiv 1 \pmod{4620}$$

GCD is a linear combination

Relative primes

Fundamental theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's Algorithm

Multiplicative inverse

Find inverse of 101 modulo 4620, that is x such that

$$101 \cdot x \equiv 1 \pmod{4620}$$

If 101 and 4620 are relative primes:

$$\gcd(101, 4620) = 1,$$

by Bezout's theorem: Exist s and t such that

$$101 \cdot s + 4620 \cdot t = \gcd(101, 4620) = 1$$

$$101 \cdot s \equiv 1 \pmod{4620}$$

We have to find Bezout coefficients s and t . Then s is the inverse.

GCD is a linear combination

Relative primes

Fundamental theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's Algorithm

Recall Euclid's Algorithm

$$a_0 = 4620 = 45 \cdot 101 + 75$$

$$a_1 = 101 = 1 \cdot 75 + 26$$

$$a_2 = 75 = 2 \cdot 26 + 23$$

$$a_3 = 26 = 1 \cdot 23 + 3$$

$$a_4 = 23 = 7 \cdot 3 + 2$$

$$a_5 = 3 = 1 \cdot 2 + 1$$

$$a_6 = 2 = 2 \cdot 1$$

$$a_7 = 1$$

Let's adapt this algorithm for finding Bezout coefficients s and t :

$$101 \cdot s + 4620 \cdot t = 1$$

GCD is a linear combination

Relative primes

Fundamental theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's Algorithm

Extended Euclid's Algorithm

$$101 \cdot s + 4620 \cdot t = 1$$

Run Euclid's algorithm:

$$a_0 = 4620 = 45 \cdot 101 + 75$$

$$a_1 = 101 = 1 \cdot 75 + 26$$

$$a_2 = 75 = 2 \cdot 26 + 23$$

$$a_3 = 26 = 1 \cdot 23 + 3$$

$$a_4 = 23 = 7 \cdot 3 + 2$$

$$a_5 = 3 = 1 \cdot 2 + 1$$

$$a_6 = 2 = 2 \cdot 1$$

$$a_7 = 1$$

Work backwards, to express GCD in terms of $a_1 = 101$ and $a_0 = 4620$:

$$1 = 3 - 1 \cdot 2$$

GCD is a linear combination

Relative primes

Fundamental theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's Algorithm

Extended Euclid's Algorithm

$$101 \cdot s + 4620 \cdot t = 1$$

Run Euclid's algorithm:

$$a_0 = 4620 = 45 \cdot 101 + 75$$

$$a_1 = 101 = 1 \cdot 75 + 26$$

$$a_2 = 75 = 2 \cdot 26 + 23$$

$$a_3 = 26 = 1 \cdot 23 + 3$$

$$a_4 = 23 = 7 \cdot 3 + 2$$

$$a_5 = 3 = 1 \cdot 2 + 1$$

$$a_6 = 2 = 2 \cdot 1$$

$$a_7 = 1$$

Work backwards, to express GCD in terms of $a_1 = 101$ and $a_0 = 4620$:

$$1 = 3 - 1 \cdot 2$$

$$= 3 - 1 \cdot (23 - 7 \cdot 3) = -1 \cdot 23 + 8 \cdot 3$$

GCD is a linear combination

Relative primes

Fundamental theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's Algorithm

Extended Euclid's Algorithm

GCD is a linear combination

Relative primes

Fundamental theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's Algorithm

$$101 \cdot s + 4620 \cdot t = 1$$

Run Euclid's algorithm:

$$a_0 = 4620 = 45 \cdot 101 + 75$$

$$a_1 = 101 = 1 \cdot 75 + 26$$

$$a_2 = 75 = 2 \cdot 26 + 23$$

$$a_3 = 26 = 1 \cdot 23 + 3$$

$$a_4 = 23 = 7 \cdot 3 + 2$$

$$a_5 = 3 = 1 \cdot 2 + 1$$

$$a_6 = 2 = 2 \cdot 1$$

$$a_7 = 1$$

Work backwards, to express GCD in terms of $a_1 = 101$ and $a_0 = 4620$:

$$1 = 3 - 1 \cdot 2$$

$$= 3 - 1 \cdot (23 - 7 \cdot 3) = -1 \cdot 23 + 8 \cdot 3$$

$$= -1 \cdot 23 + 8 \cdot (26 - 1 \cdot 23) = 8 \cdot 26 - 9 \cdot 23$$

Extended Euclid's Algorithm

GCD is a linear combination

Relative primes

Fundamental theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's Algorithm

$$101 \cdot s + 4620 \cdot t = 1$$

Run Euclid's algorithm:

$$a_0 = 4620 = 45 \cdot 101 + 75$$

$$a_1 = 101 = 1 \cdot 75 + 26$$

$$a_2 = 75 = 2 \cdot 26 + 23$$

$$a_3 = 26 = 1 \cdot 23 + 3$$

$$a_4 = 23 = 7 \cdot 3 + 2$$

$$a_5 = 3 = 1 \cdot 2 + 1$$

$$a_6 = 2 = 2 \cdot 1$$

$$a_7 = 1$$

Work backwards, to express GCD in terms of $a_1 = 101$ and $a_0 = 4620$:

$$1 = 3 - 1 \cdot 2$$

$$= 3 - 1 \cdot (23 - 7 \cdot 3) = -1 \cdot 23 + 8 \cdot 3$$

$$= -1 \cdot 23 + 8 \cdot (26 - 1 \cdot 23) = 8 \cdot 26 - 9 \cdot 23$$

$$= 8 \cdot 26 - 9(75 - 2 \cdot 26) = -9 \cdot 75 + 26 \cdot 26$$

Extended Euclid's Algorithm

GCD is a linear combination

Relative primes

Fundamental theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's Algorithm

$$101 \cdot s + 4620 \cdot t = 1$$

Run Euclid's algorithm:

$$a_0 = 4620 = 45 \cdot 101 + 75$$

$$a_1 = 101 = 1 \cdot 75 + 26$$

$$a_2 = 75 = 2 \cdot 26 + 23$$

$$a_3 = 26 = 1 \cdot 23 + 3$$

$$a_4 = 23 = 7 \cdot 3 + 2$$

$$a_5 = 3 = 1 \cdot 2 + 1$$

$$a_6 = 2 = 2 \cdot 1$$

$$a_7 = 1$$

Work backwards, to express GCD in terms of $a_1 = 101$ and $a_0 = 4620$:

$$1 = 3 - 1 \cdot 2$$

$$= 3 - 1 \cdot (23 - 7 \cdot 3) = -1 \cdot 23 + 8 \cdot 3$$

$$= -1 \cdot 23 + 8 \cdot (26 - 1 \cdot 23) = 8 \cdot 26 - 9 \cdot 23$$

$$= 8 \cdot 26 - 9(75 - 2 \cdot 26) = -9 \cdot 75 + 26 \cdot 26$$

$$= -9 \cdot 75 + 26 \cdot (101 - 1 \cdot 75) = 26 \cdot 101 - 35 \cdot 75$$

Extended Euclid's Algorithm

GCD is a linear combination

Relative primes

Fundamental theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's Algorithm

$$101 \cdot s + 4620 \cdot t = 1$$

Run Euclid's algorithm:

$$a_0 = 4620 = 45 \cdot 101 + 75$$

$$a_1 = 101 = 1 \cdot 75 + 26$$

$$a_2 = 75 = 2 \cdot 26 + 23$$

$$a_3 = 26 = 1 \cdot 23 + 3$$

$$a_4 = 23 = 7 \cdot 3 + 2$$

$$a_5 = 3 = 1 \cdot 2 + 1$$

$$a_6 = 2 = 2 \cdot 1$$

$$a_7 = 1$$

Work backwards, to express GCD in terms of $a_1 = 101$ and $a_0 = 4620$:

$$1 = 3 - 1 \cdot 2$$

$$= 3 - 1 \cdot (23 - 7 \cdot 3) = -1 \cdot 23 + 8 \cdot 3$$

$$= -1 \cdot 23 + 8 \cdot (26 - 1 \cdot 23) = 8 \cdot 26 - 9 \cdot 23$$

$$= 8 \cdot 26 - 9(75 - 2 \cdot 26) = -9 \cdot 75 + 26 \cdot 26$$

$$= -9 \cdot 75 + 26 \cdot (101 - 1 \cdot 75) = 26 \cdot 101 - 35 \cdot 75$$

$$= 26 \cdot 101 - 35 \cdot (4620 - 45 \cdot 101)$$

$$= -35 \cdot 4620 + 1601 \cdot 101$$

Extended Euclid's Algorithm

GCD is a linear combination

Relative primes

Fundamental theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's Algorithm

$$-35 \cdot 4620 + 1601 \cdot 101 = 1$$

Bezout coefficients are $s = 1601$ and $t = -35$.

Therefore, $s = 1601$ is the multiplicative inverse:

$$101 \cdot 1601 \equiv 1 \pmod{4620}$$

It works, but it's easy to make a mistake using this method. Let's describe the extended Euclid's algorithm more systematically.

Extended Euclid's Algorithm (II)

The task:

Given two numbers $a_0 \geq a_1$, run Euclid's algorithm, computing

$$a_2 = \dots$$

$$a_3 = \dots$$

\dots

$$a_k = \gcd(a_0, a_1)$$

In addition, find the coefficients x_k and y_k such that

$$a_k = x_k a_0 + y_k a_1$$

We find a recurrent solution for x_k and y_k .

GCD is a linear combination

Relative primes

Fundamental theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's Algorithm

Extended Euclid's Algorithm (II)

Need to find the coefficients x_k and y_k such that

$$a_k = \gcd(a_0, a_1) = x_k a_0 + y_k a_1$$

But we compute more than that. We want to represent all a_i as a linear combination of a_0 and a_1

$$a_0 = x_0 a_0 + y_0 a_1$$

$$a_1 = x_1 a_0 + y_1 a_1$$

$$a_2 = x_2 a_0 + y_2 a_1$$

$$a_3 = x_3 a_0 + y_3 a_1$$

...

$$a_k = \gcd(a_0, a_1) = x_k a_0 + y_k a_1$$

GCD is a linear combination

Relative primes

Fundamental theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's Algorithm

Extended Euclid's Algorithm (II)

$$a_0 = x_0 a_0 + y_0 a_1$$

$$a_1 = x_1 a_0 + y_1 a_1$$

$$a_2 = x_2 a_0 + y_2 a_1$$

$$a_3 = x_3 a_0 + y_3 a_1$$

...

$$a_k = x_k a_0 + y_k a_1$$

GCD is a linear combination

Relative primes

Fundamental theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's Algorithm

Extended Euclid's Algorithm (II)

$$a_0 = x_0 a_0 + y_0 a_1 \quad a_0 = 1a_0 + 0a_1, \quad x_0 = 1, \quad y_0 = 0,$$

$$a_1 = x_1 a_0 + y_1 a_1 \quad a_1 = 0a_0 + 1a_1, \quad x_1 = 0, \quad y_1 = 1,$$

$$a_2 = x_2 a_0 + y_2 a_1$$

$$a_3 = x_3 a_0 + y_3 a_1$$

...

$$a_k = x_k a_0 + y_k a_1$$

GCD is a linear combination

Relative primes

Fundamental theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's Algorithm

Extended Euclid's Algorithm (II)

$$\begin{aligned}a_0 &= x_0 a_0 + y_0 a_1 & a_0 &= 1 a_0 + 0 a_1, & x_0 &= 1, & y_0 &= 0, \\a_1 &= x_1 a_0 + y_1 a_1 & a_1 &= 0 a_0 + 1 a_1, & x_1 &= 0, & y_1 &= 1, \\a_2 &= x_2 a_0 + y_2 a_1 \\a_3 &= x_3 a_0 + y_3 a_1 \\&\dots \\a_k &= x_k a_0 + y_k a_1\end{aligned}$$

The other x_i and y_i can be derived using the relations between a_i 's:

$$a_i = a_{i-2} - q_{i-1} \cdot a_{i-1}$$

GCD is a linear combination

Relative primes

Fundamental theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's Algorithm

Extended Euclid's Algorithm (II)

Euclid's algorithm computes the next remainder, a_i , this way:

$$a_i = a_{i-2} - q_{i-1} \cdot a_{i-1}$$

Two previous remainders are

$$a_{i-2} = x_{i-2}a_0 + y_{i-2}a_1 \quad \text{and} \quad a_{i-1} = x_{i-1}a_0 + y_{i-1}a_1$$

$$a_i = a_{i-2} - q_{i-1} \cdot a_{i-1}$$

$$= x_{i-2} \cdot a_0 + y_{i-2} \cdot a_1 - q_{i-1}(x_{i-1} \cdot a_0 + y_{i-1} \cdot a_1)$$

$$= (x_{i-2} - q_{i-1}x_{i-1}) \cdot a_0 + (y_{i-2} - q_{i-1}y_{i-1}) \cdot a_1$$

$$= \underbrace{\left(x_{i-2} - \left(\frac{a_{i-2} - a_i}{a_{i-1}} \right) x_{i-1} \right)}_{=x_i} \cdot a_0 + \underbrace{\left(y_{i-2} - \left(\frac{a_{i-2} - a_i}{a_{i-1}} \right) y_{i-1} \right)}_{=y_i} \cdot a_1$$

GCD is a linear combination

Relative primes

Fundamental theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's Algorithm

Extended Euclid's Algorithm (II)

$$a_i = a_{i-2} - q_{i-1} \cdot a_{i-1}$$

This is how we compute all x_i and y_i up to x_k and y_k :

$$x_0 = 1$$

$$x_1 = 0$$

...

$$x_i = x_{i-2} - \underbrace{\left(\frac{a_{i-2} - a_i}{a_{i-1}} \right)}_{=q_{i-1}} x_{i-1}$$

...

$$y_0 = 0$$

$$y_1 = 1$$

...

$$y_i = y_{i-2} - \underbrace{\left(\frac{a_{i-2} - a_i}{a_{i-1}} \right)}_{=q_{i-1}} y_{i-1}$$

...

In the end, we get two numbers x_k and y_k , so we can express the GCD as a linear combination of a_0 and a_1 :

$$\gcd(a_0, a_1) = a_k = x_k \cdot a_0 + y_k \cdot a_1$$

GCD is a linear combination

Relative primes

Fundamental theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's Algorithm

Extended Euclid's Algorithm (II)

$$x_i = x_{i-2} - \underbrace{\left(\frac{a_{i-2} - a_i}{a_{i-1}} \right)}_{=q_{i-1}} x_{i-1}$$

$$y_i = y_{i-2} - \underbrace{\left(\frac{a_{i-2} - a_i}{a_{i-1}} \right)}_{=q_{i-1}} y_{i-1}$$

i	a_i	q	x_i	y_i
0	$a_0 = 4620$	-	1	0
1	$a_1 = 101$	-	0	1
2	$4620 = 45 \cdot 101 + 75$ $a_2 = 75$	45	$1 - 45 \cdot 0 =$ 1	$0 - 45 \cdot 1 =$ -45

GCD is a linear combination

Relative primes

Fundamental theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's Algorithm

Extended Euclid's Algorithm (II)

$$x_i = x_{i-2} - \underbrace{\left(\frac{a_{i-2} - a_i}{a_{i-1}} \right)}_{=q_{i-1}} x_{i-1}$$

$$y_i = y_{i-2} - \underbrace{\left(\frac{a_{i-2} - a_i}{a_{i-1}} \right)}_{=q_{i-1}} y_{i-1}$$

i	a_i	q	x_i	y_i
0	$a_0 = 4620$	-	1	0
1	$a_1 = 101$	-	0	1
2	$4620 = 45 \cdot 101 + 75$ $a_2 = 75$	45	$1 - 45 \cdot 0 =$ 1	$0 - 45 \cdot 1 =$ -45
3	$101 = 1 \cdot 75 + 26$ $a_3 = 26$	1	$0 - 1 \cdot 1 =$ -1	$1 - 1 \cdot (-45) =$ 46
4	$75 = 2 \cdot 26 + 23$ $a_4 = 23$	2	$1 - 2 \cdot (-1) =$ 3	$-45 - 2 \cdot 46 =$ -137

GCD is a linear combination

Relative primes

Fundamental theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's Algorithm

Extended Euclid's Algorithm (II)

i	a_i	q	x_i	y_i
0	$a_0 = 4620$	-	1	0
1	$a_1 = 101$	-	0	1
2	$a_2 = 75$	45	1	-45
3	$a_3 = 26$	1	-1	46
4	$a_4 = 23$	2	3	-137
5	$26 = 1 \cdot 23 + 3$ $a_5 = 3$	1	$-1 - 1 \cdot 3 = -4$	$46 - 1 \cdot (-137) = 183$
6	$23 = 7 \cdot 3 + 2$ $a_6 = 2$	7	$3 - 7 \cdot (-4) = 31$	$-137 - 7 \cdot 183 = -1418$
7	$3 = 1 \cdot 2 + 1$ $a_7 = 1$	1	$-4 - 1 \cdot 31 = -35$	$183 - 1 \cdot 1418 = -1235$

GCD is a linear combination

Relative primes

Fundamental theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's Algorithm

Extended Euclid's Algorithm (II)

While computing the sequence of a_i 's with Euclid's algorithm, we eventually produced coefficients

$$x_7 = -35, \quad y_7 = 1601$$

By construction, they satisfy the equation

$$a_7 = x_7 \cdot a_0 + y_7 \cdot a_1$$

$$1 = \underbrace{-35}_{=x_7} \cdot \underbrace{4620}_{=a_0} + \underbrace{1601}_{=y_7} \cdot \underbrace{101}_{=a_1}$$

But from the last equation we can find the inverse of 101 modulo 4620, and the inverse of 4620 modulo 101.

GCD is a linear combination

Relative primes

Fundamental theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's Algorithm

Finding a multiplicative inverse

Take this equation and find the multiplicative inverse of $a_1 = 101$ modulo $a_0 = 4620$.

$$1 = \underbrace{-35}_{=x_7} \cdot \underbrace{4620}_{=a_0} + \underbrace{1601}_{=y_7} \cdot \underbrace{101}_{=a_1}$$

$$1601 \cdot 101 - 1 = 35 \cdot 4620$$

Therefore, by definition of congruence,

$$101 \cdot 1601 \equiv 1 \pmod{4620}.$$

So, 1601 is a multiplicative inverse of 101 modulo 4620.

We were able to find the inverse, because 101 and 4620 are relative primes, that is, their GCD is equal to 1.

GCD is a linear combination

Relative primes

Fundamental theorem of arithmetic

Modular arithmetic

Congruence

Modular arithmetic

Multiplicative inverse

Extended Euclid's Algorithm