

Induction



The idea


Idea

Principle

Examples

Summations

Inequalities



Consider a logical problem

Idea

Principle

Examples

Summations

Inequalities

$$\begin{array}{c} p_0 \\ p_0 \rightarrow p_1 \\ p_1 \rightarrow p_2 \\ \dots \\ p_n \rightarrow p_{n+1} \\ \dots \\ \hline p_k \text{ for all } k \geq 0 \end{array}$$

The same, but using predicate P

Idea

Principle

Examples

Summations

Inequalities

We can represent all propositions with a predicate P :

$$P(n) = p_n$$

$$P(0)$$

$$P(n) \rightarrow P(n+1) \text{ for all } n \geq 0$$

$$P(k) \text{ for all } k \geq 0$$

Principle of induction


Idea

Principle

Examples

Summations

Inequalities



Principle of induction

Idea

Principle

Examples

Summations

Inequalities



Principle of induction


Idea

Principle

Examples

Summations

Inequalities



Principle of induction

Idea

Principle

Examples

Summations

Inequalities

If

- $P(0)$ is true
(the base case), and
- for all $n \geq 0$, $P(n)$ implies $P(n + 1)$
(the inductive step),

then $P(k)$ is true for every $k \in \mathbb{N}$.

Example proof

Idea

Principle

Examples

Summations

Inequalities

Let's use induction to prove the formula of the sum of natural numbers from 0 to n :

$$0 + 1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2}$$

(We have to show that both, the *base case* and the *inductive step* are correct)

Example proof

Idea
Principle
Examples
Summations
Inequalities

Prove that

$$0 + 1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2}$$

Part 1. *The base case.*

For $n = 0$:

$$0 = \frac{0 \cdot (0 + 1)}{2}$$

It is true.

Example proof

Part 2. *The inductive step.*

Assume that the formula is true for an arbitrary $n \geq 0$:

$$0 + 1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2}$$

We have to prove that

$$0 + 1 + 2 + 3 + \dots + (n+1) = \frac{(n+1)((n+1)+1)}{2}$$

Idea

Principle

Examples

Summations

Inequalities

Example proof

Idea

Principle

Examples

Summations

Inequalities

Part 2. *The inductive step.*

Assume that the formula is true for an arbitrary $n \geq 0$:

$$0 + 1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2}$$

We have to prove that

$$0 + 1 + 2 + 3 + \dots + (n+1) = \frac{(n+1)((n+1)+1)}{2}$$

Consider the left-hand-side:

$$\begin{aligned} \underbrace{1 + 2 + 3 + \dots + n}_{= \frac{n(n+1)}{2}} + (n+1) &= \frac{n(n+1)}{2} + n + 1 = \frac{n(n+1) + 2(n+1)}{n+1} \\ &= \frac{(n+1)(n+2)}{2} = \frac{(n+1)((n+1)+1)}{2} \end{aligned}$$

Example proof

- Idea
- Principle
- Examples
- Summations
- Inequalities

The base case and the inductive step are true.

Therefore, *by induction*, the formula is correct for every natural number n .


$$0 + 1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2}$$

This is a very useful formula, by the way.

Rewrite it using the sigma-notation

$$\sum_{k=1}^n k = 1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2}$$

Tiling $2^n \times 2^n$ with 1 square removed



For all $n > 0$, a checkerboard $2^n \times 2^n$ with one square removed


can be tiled by L-shaped tiles



- Idea
- Principle
- Examples
- Summations
- Inequalities

Tiling $2^n \times 2^n$ with 1 square removed

Idea
Principle
Examples
Summations
Inequalities



For all $n > 0$, a checkerboard $2^n \times 2^n$ with one square removed

can be tiled by L-shaped tiles




The base case for $n=1$



Inductive step.

Assuming that we can tile $2^n \times 2^n$ with one removed, prove that it's possible to tile $2^{n+1} \times 2^{n+1}$ with one removed



Another example proof

Idea
Principle
Examples
Summations
Inequalities

Let $P(n)$ be the predicate, "*I can lift n grains of sand.*"

- I can lift one grain of sand, so $P(1)$ is true. This is my base case.

Another example proof

Idea
Principle
Examples
Summations
Inequalities

Let $P(n)$ be the predicate, “I can lift n grains of sand.”

- I can lift one grain of sand, so $P(1)$ is true. This is my base case.
- Then, surely, if I can lift n grains, then I can lift $n + 1$, it does not make any difference!

$$P(n) \rightarrow P(n + 1)$$

This is my inductive step.

Another example proof

Idea

Principle

Examples

Summations

Inequalities

Let $P(n)$ be the predicate, “I can lift n grains of sand.”

- I can lift one grain of sand, so $P(1)$ is true. This is my base case.
- Then, surely, if I can lift n grains, then I can lift $n + 1$, it does not make any difference!

$$P(n) \rightarrow P(n + 1)$$

This is my inductive step.

So, by induction, *I can lift any amount of sand*. Right?

Where is a mistake?

Idea

Principle

Examples

Summations

Inequalities

Of course, the proof is wrong. But should we blame induction for that?

Well, we made an error in the proof of $P(n) \rightarrow P(n + 1)$.

It is hard to say for exactly which n it is false, but certainly there is some value!

Sum of b^k

Idea

Principle

Examples

Summations

Inequalities

Prove by induction that

$$b^0 + b^1 + b^2 + \dots + b^n = \frac{b^{n+1} - 1}{b - 1}$$

First we can do the base case, then the inductive step.

Sum of b^k

Idea

Principle

Examples

Summations

Inequalities

Prove that

$$b^0 + b^1 + b^2 + \dots + b^n = \frac{b^{n+1} - 1}{b - 1}$$

Base case ($n = 0$):

$$b^0 = 1, \quad \text{and} \quad \frac{b^1 - 1}{b - 1} = 1$$

.

Sum of b^k

Idea

Principle

Examples

Summations

Inequalities

Prove that

$$b^0 + b^1 + b^2 + \dots + b^n = \frac{b^{n+1} - 1}{b - 1}$$

Inductive step:

As always, we make a hypothesis that

$$b^0 + b^1 + b^2 + \dots + b^n = \frac{b^{n+1} - 1}{b - 1} \quad \text{is true for } n \geq 0$$

And we have to prove that the formula is correct for $n + 1$:

$$b^0 + b^1 + b^2 + \dots + b^{n+1} = \frac{b^{n+2} - 1}{b - 1}$$

Sum of b^k

Idea

Principle

Examples

Summations

Inequalities

Inductive step:

We have to prove that the formula is correct for $n + 1$:

$$b^0 + b^1 + b^2 + \dots + b^{n+1} = \frac{b^{n+2} - 1}{b - 1}$$

$$\begin{aligned} \underbrace{b^0 + b^1 + b^2 + \dots + b^n}_{= \frac{b^{n+1} - 1}{b - 1} \text{ by the hypothesis}} + b^{n+1} &= \frac{b^{n+1} - 1}{b - 1} + b^{n+1} \\ &= \frac{b^{n+1} - 1 + b^{n+2} - b^{n+1}}{b - 1} = \frac{b^{n+2} - 1}{b - 1}. \end{aligned}$$

Sum of b^k

Idea

Principle

Examples

Summations

Inequalities

So, this formula for the sum is correct

$$b^0 + b^1 + b^2 + \dots + b^n = \frac{b^{n+1} - 1}{b - 1}$$

In the sigma-notation:

$$\sum_{k=0}^n b^k = b^0 + b^1 + b^2 + \dots + b^n = \frac{b^{n+1} - 1}{b - 1}$$

Sum of b^k

Idea

Principle

Examples

Summations

Inequalities

We can multiply both sides by a constant a :

$$\sum_{k=0}^n ab^k = ab^0 + ab^1 + ab^2 + \dots + ab^n = \frac{a(b^{n+1} - 1)}{b - 1},$$

The sequence of numbers

$$a, ab, ab^2, ab^3, \dots, ab^n, \dots$$

is called a *Geometric progression*.

So, we proved the formula for the partial sum of a geometric progression.

Sum of kb^{k-1}

Idea

Principle

Examples

Summations

Inequalities

The partial sum of the geometric progression:

$$\sum_{k=0}^n ab^k = ab^0 + ab^1 + ab^2 + \dots + ab^n = \frac{a(b^{n+1} - 1)}{b - 1},$$

Can we compute

$$\begin{aligned} \sum_{k=0}^n kb^{k-1} \\ = 0 + 1 + 2b + 3b^2 + 4b^3 \dots + nb^{n-1} \end{aligned}$$

So, instead of the constant a , we have an increasing sequence of coefficients now.

Sum of kb^{k-1}

Idea

Principle

Examples

Summations

Inequalities

There is a cheap trick:

$$\begin{aligned}\frac{d}{db} \sum_{k=0}^n b^k &= \frac{d}{db} (1 + b + b^2 + b^3 + b^4 + \dots + b^n) \\ &= 0 + 1 + 2b + 3b^2 + 4b^3 + \dots + nb^{n-1} = \sum_{k=0}^n kb^{k-1}\end{aligned}$$

On the other hand,

$$\begin{aligned}\frac{d}{db} \sum_{k=0}^n b^k &= \frac{d}{db} \left(\frac{b^{n+1} - 1}{b - 1} \right) = \frac{(n+1)b^n(b-1) - b^{n+1} + 1}{(b-1)^2} \\ &= \frac{nb^{n+1} - (n+1)b^n + 1}{(b-1)^2}\end{aligned}$$

$$\text{Therefore, } \sum_{k=0}^n kb^{k-1} = \frac{nb^{n+1} - (n+1)b^n + 1}{(b-1)^2}.$$

Geometric progression again

Idea

Principle

Examples

Summations

Inequalities

The partial sum of the geometric progression:

$$\sum_{k=0}^n ab^k = ab^0 + ab^1 + ab^2 + \dots + ab^n = \frac{a(b^{n+1} - 1)}{b - 1}$$

Well, what if we want to add up an infinite sequence?

$$\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \dots$$

Infinite geometric progression

Idea

Principle

Examples

Summations

Inequalities

The partial sum of the geometric progression is

$$\sum_{k=0}^n ab^k = ab^0 + ab^1 + ab^2 + \dots + ab^n = \frac{a(b^{n+1} - 1)}{b - 1}$$

If b is a small real number, specifically, if the absolute value $|b| < 1$, then

$$|b| > |b^2| > |b^3| > \dots$$

In the limit, $\lim_{n \rightarrow \infty} b^n = 0$

$$\sum_{n=0}^{\infty} ab^n = \lim_{n \rightarrow \infty} ab^n = \lim_{n \rightarrow \infty} \frac{a(b^{n+1} - 1)}{b - 1} = \frac{a(-1)}{b - 1} = \frac{a}{1 - b}$$

Proving inequalities

Idea

Principle

Examples

Summations

Inequalities

Using mathematical induction, prove that for $n \geq 1$:

$$n < 2^n$$

Proving inequalities

- Idea
- Principle
- Examples
- Summations
- Inequalities

Using mathematical induction, prove that for $n \geq 1$:

$$n < 2^n$$

The base case:

$n = 1$. $1 < 2$ is true.

The inductive step:

Assume that $n < 2^n$ for $n \geq 1$. We have to prove that $n + 1 < 2^{n+1}$.

$$n + 1 < 2^n + 1 < 2^n + 2^n = 2^{n+1}$$

Therefore, *by induction*, $n < 2^n$ is true for $n \geq 1$.

One more proof



Idea
Principle
Examples
Summations
Inequalities

Theorem. All horses are the same color.

We can prove this by induction on the number of horses in a given set. Here's how: If there's just one horse then it's the same color as itself, so the base case is trivial.

All horses are the same color

Idea


Principle

Examples

Summations

Inequalities

For the induction step, assume that there are $n+1$ horses numbered 1 to $n+1$. By the induction hypothesis, horses 1 through n are the same color, and similarly horses 2 through $n+1$ are the same color.



But the middle horses, 2 through n , can't change color when they're in different groups; these are horses, not chameleons. So horses 1 and $n+1$ must be the same color as well. Thus all $n+1$ horses are the same color.

What, if anything, is wrong with this reasoning?