

Discrete Structures, CSCI-150.

Information

Information

Propositions

Operators

Complex propositions

Equivalence

Mon, Wed, 7:00 – 8:15 pm. Hunter North C107.

Instructor: Alexey Nikolaev.

Website: <http://a-nikolaev.github.io/ds/>

Grading policy

No late homeworks accepted. Homeworks every week. Due at the beginning of the class.

Final grade:

HWs: 25%

Midterm I: 25%

Midterm II: 25%

Final: 25%

Course content

Propositional Logic. Operators. Truth tables. Logical equivalence. Rules of inference. Satisfiability. Predicates and quantifiers. Proofs.

Induction. Hanoi towers. Summation of series. Recurrence. Fibonacci numbers. Catalan numbers. Solving linear recurrence.

Number theory. Divisibility and primes. Modulo-arithmetics. GCD and Euclid's algorithm. Cryptography. RSA.

Counting. Sum and product rules. Pigeonhole principle. Permutations, $n!$ Binomial coefficients, n choose k . Selection with replacement.

Sets. Operations, empty set, singleton set, powerset. Natural, rational, real numbers. Diagonalization. Relations and Functions. Counting and Bijection. Partial orders.

Graphs. Bridges of Königsberg. Eulerian and Hamiltonian cycles. Trees, spanning trees. Huffman coding.

Probability. Bernoulli Trials. Random variables. Expected value.

Information

Propositions

Operators

Complex propositions

Equivalence

Literature

Information

Propositions

Operators

Complex propositions

Equivalence

Primary books:

- *Rosen*
“Discrete Mathematics and its Applications” edition 6 or 7.
(you can find a used or new 6th edition for less than \$40).
- *Lehman and Leighton*
Lecture notes “Mathematics for Computer Science” (2004).
(free, but this is not a complete textbook)

Our first object

Information

Propositions

Operators

Complex propositions

Equivalence

Something that is either

true or *false*

Propositions

Information

Propositions

Operators

Complex propositions

Equivalence

Def. A **proposition** is a declarative sentence that is either **true** or **false**, but not both.

A good test for a proposition is to ask “*Is it true that ... ?*” If that makes sense, it is a proposition.

- One plus two equals three.
- Washington, D.C., is the capital of the US.
- The Moon is a satellite of the Earth.
- Albany is the capital of Canada.
- The Sun is a planet.

Propositions

Information

Propositions

Operators

Complex propositions

Equivalence

Def. A **proposition** is a declarative sentence that is either **true** or **false**, but not both.

A good test for a proposition is to ask “*Is it true that ... ?*” If that makes sense, it is a proposition.

- One plus two equals three. *true* ✓
- Washington, D.C., is the capital of the US. *true* ✓
- The Moon is a satellite of the Earth. *true* ✓
- Albany is the capital of Canada. *false* ✓
- The Sun is a planet. *false* ✓ all are propositions

Propositions

Information

Propositions

Operators

Complex propositions

Equivalence

Def. A **proposition** is a declarative sentence that is either **true** or **false**, but not both.

A good test for a proposition is to ask “*Is it true that ... ?*” If that makes sense, it is a proposition.

- Three plus four.
- Consider these sentences.
- Does anyone have any questions?
- The largest planet in the Solar System.
- n in a prime number.

Propositions

Information

Propositions

Operators

Complex propositions

Equivalence

Def. A **proposition** is a declarative sentence that is either **true** or **false**, but not both.

A good test for a proposition is to ask “*Is it true that ... ?*” If that makes sense, it is a proposition.

- Three plus four. ✗ neither one is a proposition
- Consider these sentences. ✗
- Does anyone have any questions? ✗
- The largest planet in the Solar System. ✗
- n in a prime number. ✗

Propositions

Information

Propositions

Operators

Complex propositions

Equivalence

Instead of writing sentences, we will abbreviate them by using *propositional variables*.

It is standard practice to use the lower-case letters: p, q, r, \dots

Then, if

$$p = \text{"It is raining"},$$
$$q = \text{"I have an umbrella"},$$

we can construct *compound propositions* using logical operators:

$$p \text{ and } q = \text{"It is raining, and I have an umbrella"}.$$
$$\text{not } q = \text{"I don't have an umbrella"}.$$

Logical Operators

Information

Propositions

Operators

Complex propositions

Equivalence

And (called Conjunction)

p *and* q

$p \wedge q$ is true when both p and q are true, otherwise false.

Or (called Disjunction)

p *or* q

$p \vee q$ is true when p or q or both are true, otherwise false.

Negation

not p

$\neg p$ is true when p is false, otherwise false.

Truth tables

Information

Propositions

Operators

Complex propositions

Equivalence

And (Conjunction)

p	q	$p \wedge q$
T	T	T
F	T	F
T	F	F
F	F	F

Or (Disjunction)

p	q	$p \vee q$
T	T	T
F	T	T
T	F	T
F	F	F

Negation

p	$\neg p$
T	F
F	T

Think of the truth tables as our ultimate definition of the logical connectives (operators).

More Operators. Implication

Information

Propositions

Operators

Complex propositions

Equivalence

Implication

if p then q

$p \rightarrow q$ is true if whenever p is true, so is q , otherwise false.

Truth table:

*An implication is true when
the if-part is false or the then-part is true.*

p	q	$p \rightarrow q$
T	T	T
F	T	T
T	F	F
F	F	T

So, $p \rightarrow q$ is equivalent to $(\neg p) \vee q$.

“I need an umbrella, if it’s raining”.

“If the Earth is flat, my brother is a physicist”.

More Operators. Implication


Information

Propositions

Operators

Complex propositions

Equivalence



1. *"If he is hungry, he is grumpy".*
2. *"He is hungry".*

Is he grumpy?

More Operators. Implication

Information

Propositions

Operators

Complex propositions

Equivalence

“If he is hungry, he is grumpy”.

$$h \rightarrow g = T$$

If an implication is true, we can make conclusions about g , if we know h .

We know that *“he is hungry”*,

$$h = T,$$

it's only possible that he is grumpy

$$g = T.$$

h	g	$h \rightarrow g$
T	T	T
F	T	T
T	F	F
F	F	T

More Operators. Implication

Information

Propositions

Operators

Complex propositions

Equivalence



1. *"If he is hungry, he is grumpy".*
2. *"He is not hungry".*

Is he happy?

More Operators. Implication

Information

Propositions

Operators

Complex propositions

Equivalence

“If he is hungry, he is grumpy”.

$$h \rightarrow g = T$$

If an implication is true, we can make conclusions about g , if we know h .

If we know that *“he is not hungry”*,

$$h = F,$$

then

g can be T or F .

h	g	$h \rightarrow g$
T	T	T
F	T	T
T	F	F
F	F	T

More Operators. Implication

Information

Propositions

Operators

Complex propositions

Equivalence

p	q	$p \rightarrow q$
T	T	T
F	T	T
T	F	F
F	F	T



Big Al told these guys that dogs can't look up.

Their thoughts:

$p \rightarrow q$ = "If dogs can look up, Big Al is a liar".

p = "Dogs can look up"

q = "Big Al is a liar"


$(\neg p) \vee q$ = "Dogs can't look up, or Big Al is a liar".

$p \rightarrow q$ is equivalent to $(\neg p) \vee q$

More Operators. Implication

p	q	$p \rightarrow q$
T	T	T
F	T	T
T	F	F
F	F	T

Mathematical theorems are often formulated as implications. For example:



Theorem. *If a continuous function defined on an interval is sometimes positive and sometimes negative, it must be 0 at some point.*

Information

Propositions

Operators

Complex propositions

Equivalence

More Operators. Biconditional

Biconditional

p if and only if q

$p \leftrightarrow q$

is true when p and q have the same truth values, otherwise false.

p	q	$p \leftrightarrow q$
T	T	T
F	T	F
T	F	F
F	F	T

Often, “if and only if” is abbreviated to *iff*:

$p \text{ iff } q$

“You can take the flight if and only if you buy a ticket.”

Theorems are often formulated as implications or biconditionals.

Information

Propositions

Operators

Complex propositions

Equivalence

Combined truth tables for connectives \neg , \wedge , \vee , \rightarrow , and \leftrightarrow

Information

Propositions

Operators

Complex propositions

Equivalence

p	q	$\neg p$	$p \wedge q$	$p \vee q$	$p \rightarrow q$	$p \leftrightarrow q$
T	T	F	T	T	T	T
F	T	T	F	T	T	F
T	F	F	F	T	F	F
F	F	T	F	F	T	T

Complex compound propositions?

Information

Propositions

Operators

Complex propositions

Equivalence

Let's take a complex compound proposition:

$$q \vee ((\neg q) \wedge r)$$

q or ((not q) and r)

q	r	\dots
T	T	\dots
F	T	\dots
T	F	\dots
F	F	\dots

Complex compound propositions?

Information

Propositions

Operators

Complex propositions

Equivalence

Let's take a complex compound proposition:

$$q \vee ((\neg q) \wedge r)$$

q or ((not q) and r)

q	r	$\neg q$	\dots
T	T	F	\dots
F	T	T	\dots
T	F	F	\dots
F	F	T	\dots

Complex compound propositions?

Information

Propositions

Operators

Complex propositions

Equivalence

Let's take a complex compound proposition:

$$q \vee ((\neg q) \wedge r)$$

q or ((not q) and r)

q	r	$\neg q$	$(\neg q) \wedge r$	\dots
T	T	F	F	\dots
F	T	T	T	\dots
T	F	F	F	\dots
F	F	T	F	\dots

Complex compound propositions?

Information

Propositions

Operators

Complex propositions

Equivalence

Let's take a complex compound proposition:

$$q \vee ((\neg q) \wedge r)$$

q or ((not q) and r)

q	r	$\neg q$	$(\neg q) \wedge r$	$q \vee ((\neg q) \wedge r)$
T	T	F	F	T
F	T	T	T	T
T	F	F	F	T
F	F	T	F	F

Complex compound propositions?

The number of rows in the truth table of a compound proposition is equal to 2^n , where n is the number of used propositional variables.

$$(\neg p) \vee ((q \rightarrow r) \wedge p)$$

p	q	r	\dots
T	T	T	\dots
F	T	T	\dots
T	F	T	\dots
F	F	T	\dots
T	T	F	\dots
F	T	F	\dots
T	F	F	\dots
F	F	F	\dots

Each of the three variables can take two possible values, so the system has $2 \cdot 2 \cdot 2 = 8$ possible states.

Information

Propositions

Operators

Complex propositions

Equivalence

Equivalence

Information

Propositions

Operators

Complex propositions

Equivalence

Two compound propositions are equivalent if they have the same truth values for all possible cases (have the same truth tables).

q	r	$q \vee ((\neg q) \wedge r)$	$q \vee r$
T	T	T	T
F	T	T	T
T	F	T	T
F	F	F	F

Therefore, these two propositions are logically equivalent!

We write it as follows

$$q \vee ((\neg q) \wedge r) \equiv q \vee r$$

Note that the statement of the equivalence of two compound propositions, $a \equiv b$, is not a proposition itself.

Equivalent formulae

Information

Propositions

Operators

Complex propositions

Equivalence

$$(a \wedge b) \equiv (b \wedge a) \quad \text{commutativity of } \wedge$$

$$(a \vee b) \equiv (b \vee a) \quad \text{commutativity of } \vee$$

$$((a \wedge b) \wedge c) \equiv (a \wedge (b \wedge c)) \quad \text{associativity of } \wedge$$

$$((a \vee b) \vee c) \equiv (a \vee (b \vee c)) \quad \text{associativity of } \vee$$

$$\neg(\neg a) \equiv a \quad \text{double-negation elimination}$$

$$(a \rightarrow b) \equiv (\neg b \rightarrow \neg a) \quad \text{contraposition}$$

$$(a \rightarrow b) \equiv (\neg a \vee b) \quad \text{implication elimination}$$

$$(a \leftrightarrow b) \equiv (a \rightarrow b) \wedge (b \rightarrow a) \quad \text{biconditional elimination}$$

$$\neg(a \wedge b) \equiv (\neg a \vee \neg b) \quad \text{De Morgan's Law}$$

$$\neg(a \vee b) \equiv (\neg a \wedge \neg b) \quad \text{De Morgan's Law}$$

$$(a \wedge (b \vee c)) \equiv (a \wedge b) \vee (a \wedge c) \quad \text{distributivity of } \wedge \text{ over } \vee$$

$$(a \vee (b \wedge c)) \equiv (a \vee b) \wedge (a \vee c) \quad \text{distributivity of } \vee \text{ over } \wedge$$

Equivalent formulae

Information

Propositions

Operators

Complex propositions

Equivalence

If **True** is a compound propositions that is always true, and **False** is a proposition that is always false:

$$a \wedge \mathbf{True} \equiv a \quad \text{identity}$$

$$a \vee \mathbf{False} \equiv a \quad \text{identity}$$

$$a \vee \mathbf{True} \equiv \mathbf{True} \quad \text{domination}$$

$$a \wedge \mathbf{False} \equiv \mathbf{False} \quad \text{domination}$$

$$a \vee \neg a \equiv \mathbf{True} \quad \text{complementation (excluded middle)}$$

$$a \wedge \neg a \equiv \mathbf{False} \quad \text{complementation (non-contradiction)}$$

Examples of such always true compound propositions:

True : $p \vee \neg p$, $p \leftrightarrow p$, $p \rightarrow (p \vee q)$, etc.

False : ... find a few examples of always false propositions.

Proving logical equivalences without truth tables

Information

Propositions

Operators

Complex propositions

Equivalence

We can prove a logical equivalence using a sequence of known equivalences. The goal is to show that the left-hand-side is equivalent to the right-hand-side.

For example, prove that

$$q \vee ((\neg q) \wedge r) \equiv q \vee r$$

$$\begin{aligned} q \vee ((\neg q) \wedge r) &\equiv (q \vee \neg q) \wedge (q \vee r) \\ &\equiv \mathbf{True} \wedge (q \vee r) \\ &\equiv q \vee r \end{aligned}$$

We used the distributivity of \vee over \wedge , then the complementation equivalence, then the identity equivalence.