

Scientific Computing Lab

Partial Differential Equations

Instationary Equations

Tobias Neckel

Instationary PDEs

- Stationary PDEs (time-independent):
 - Only spatial derivatives

$$\frac{\partial^2 T}{\partial x^2} + \frac{\partial^2 T}{\partial y^2} = rhs(x, y)$$

Instationary PDEs

- Stationary PDEs (time-independent):
 - Only spatial derivatives

$$\frac{\partial^2 T}{\partial x^2} + \frac{\partial^2 T}{\partial y^2} = rhs(x, y)$$

- Instationary PDEs (time-dependent):
 - Derivatives in space and time

$$\frac{\partial T}{\partial t} = \frac{\partial^2 T}{\partial x^2} + \frac{\partial^2 T}{\partial y^2}$$

Discretization

- Spatial discretization
 - Finite differences
 - See worksheet 4
- Time discretization
 - Compare ODE
 - See worksheet 2 & 3

Explicit/Implicit

- explicit time steps
 - restricted time step sizes
 - dependent on spatial step
 - no equations to solve
- implicit time step
 - unrestricted time step sizes
 - large systems of equations

Accuracy Issues

- Discretization error in time and space

$$e = e_{space} + e_{time}$$

- High-order space disc. $\rightarrow e_{time}$ dominates

- High-order time disc. $\rightarrow e_{space}$ dominates

Accuracy Issues

- Take into account:
 - Order of discretization
 - Region of stability
 - Maximum allowed timestep size


Accuracy Issues

- Take into account:
 - Order of discretization
 - Region of stability
 - Maximum allowed timestep size
- Higher-order methods do not harm (in terms of accuracy; but be aware of comput. costs!)

Explicit/Implicit

- Separation in space/time-discretization
 - Compute spatial derivatives
 - Consider function value on each grid-point as an ODE
- Sketch: 2.5 ways to a full discretization

Example




Example - Continuous

Hook's Law:

$$\sigma = E \varepsilon$$

Normal tension:

$$\sigma = \frac{F_u}{A} \quad \text{with} \quad \varepsilon = \frac{du}{dx} \quad \rightarrow \quad F_u = E A \frac{du}{dx}$$



Newton's Law:

$$dF_u = dm \frac{d^2 u}{dt^2} \quad \text{with} \quad dm = \rho A dx \quad \rightarrow \quad \frac{dF_u}{dx} = \rho A \frac{d^2 u}{dt^2}$$

Example - Continuous


$$\frac{dF_u}{dx} = \rho A \frac{d^2 u}{dt^2} \quad \text{and} \quad F_u = E A \frac{du}{dx}$$

Differentiate with regard to x:

$$E A \frac{\partial^2 u}{dx^2} = \rho A \frac{\partial^2 u}{\partial t^2}$$

$$\rightarrow E \frac{\partial^2 u}{dx^2} = \rho \frac{\partial^2 u}{\partial t^2}$$


In the following, we set $E = 1, \rho = 1$



Example - Discretization

Continuous
Equation:

$$\frac{\partial^2 u(x, t)}{\partial x^2} = \frac{\partial^2 u(x, t)}{\partial t^2}$$



Discretized
Equation:


$$\begin{aligned} & \frac{u(i+1, t) - 2u(i, t) + u(i-1, t))}{h_0 \cdot h_1} \\ &= \frac{u(i, t+1) - 2u(i, t) + u(i, t-1))}{\delta t^2} \end{aligned}$$

Example - Discretization


Discretized
Equation:

$$\frac{u(i+1,t) - 2u(i,t) + u(i-1,t))}{h_0 \cdot h_1} = \frac{u(i,t+1) - 2u(i,t) + u(i,t-1))}{\delta t^2}$$

$$u(i,t+1) = \frac{\delta t^2}{h_0 \cdot h_1} \cdot (u(i+1,t) - 2u(i,t) + u(i-1,t)) + 2u(i,t) - u(i,t-1)$$



Example – Initial/Boundary Conditions




Example – Initial/Boundary Conditions

Initial Conditions:

$$\begin{pmatrix} u(1) \\ \vdots \\ u(N) \end{pmatrix}$$

$$\begin{pmatrix} v(1) \\ \vdots \\ v(N) \end{pmatrix} = \begin{pmatrix} 0 \\ \vdots \\ 0 \end{pmatrix}$$




Boundary Conditions:

Example – Initial/Boundary Conditions

Initial Conditions:

$$\begin{pmatrix} u(1) \\ \vdots \\ u(N) \end{pmatrix}$$

$$\begin{pmatrix} v(1) \\ \vdots \\ v(N) \end{pmatrix} = \begin{pmatrix} 0 \\ \vdots \\ 0 \end{pmatrix}$$



Boundary Conditions:

$$u(0)=0$$

$$\frac{\partial u(N)}{\partial x}=0$$