

# **Vanishing results**

Alexander R. Miller

## Part 1. Character vanishing

## Part 1. Character vanishing

**Burnside** Nonlinear irreducible characters have zeros.

## Part 1. Character vanishing

**Burnside** Nonlinear irreducible characters have zeros.

**Question** What is the chance that  $\chi(g)$  equals 0?

## Part 1. Character vanishing

**Burnside** Nonlinear irreducible characters have zeros.

**Question** What is the chance that  $\chi(g)$  equals 0?

**Two natural ways to pick a random character value  $\chi(g)$**

## Part 1. Character vanishing

**Burnside** Nonlinear irreducible characters have zeros.

**Question** What is the chance that  $\chi(g)$  equals 0?

**Two natural ways to pick a random character value  $\chi(g)$**

$S_4$	1	6	3	8	6
	id	(12)	(12)(34)	(123)	(1234)
$\chi_1$	1	1	1	1	1
$\chi_2$	3	1	-1	0	-1
$\chi_3$	2	0	2	-1	0
$\chi_4$	3	-1	-1	0	1
$\chi_5$	1	-1	1	1	-1

## Part 1. Character vanishing

**Burnside** Nonlinear irreducible characters have zeros.

**Question** What is the chance that  $\chi(g)$  equals 0?

**Two natural ways to pick a random character value  $\chi(g)$**

$S_4$	1	6	3	8	6
	id	(12)	(12)(34)	(123)	(1234)
$\chi_1$	1	1	1	1	1
$\chi_2$	3	1	-1	0	-1
$\chi_3$	2	0	2	-1	0
$\chi_4$	3	-1	-1	0	1
$\chi_5$	1	-1	1	1	-1

1.

## Part 1. Character vanishing

**Burnside** Nonlinear irreducible characters have zeros.

**Question** What is the chance that  $\chi(g)$  equals 0?

**Two natural ways to pick a random character value  $\chi(g)$**

$S_4$	1	6	3	8	6
	id	(12)	(12)(34)	(123)	(1234)
$\chi_1$	1	1	1	1	1
$\chi_2$	3	1	-1	0	-1
$\chi_3$	2	0	2	-1	0
$\chi_4$	3	-1	-1	0	1
$\chi_5$	1	-1	1	1	-1

1. Choose  $\chi \in \text{Irr}(G)$  and  $g \in G$ , and then evaluate  $\chi(g)$ .



## Part 1. Character vanishing

**Burnside** Nonlinear irreducible characters have zeros.

**Question** What is the chance that  $\chi(g)$  equals 0?

**Two natural ways to pick a random character value  $\chi(g)$**

$S_4$	1	6	3	8	6
	id	(12)	(12)(34)	(123)	(1234)
$\chi_1$	1	1	1	1	1
$\chi_2$	3	1	-1	0	-1
$\chi_3$	2	0	2	-1	0
$\chi_4$	3	-1	-1	0	1
$\chi_5$	1	-1	1	1	-1

1. Choose  $\chi \in \text{Irr}(G)$  and  $g \in G$ , and then evaluate  $\chi(g)$ .

$$\text{Prob}(\chi(g) = 0)$$

## Part 1. Character vanishing

**Burnside** Nonlinear irreducible characters have zeros.

**Question** What is the chance that  $\chi(g)$  equals 0?

**Two natural ways to pick a random character value  $\chi(g)$**

$S_4$	1	6	3	8	6
	id	(12)	(12)(34)	(123)	(1234)
$\chi_1$	1	1	1	1	1
$\chi_2$	3	1	-1	0	-1
$\chi_3$	2	0	2	-1	0
$\chi_4$	3	-1	-1	0	1
$\chi_5$	1	-1	1	1	-1

1. Choose  $\chi \in \text{Irr}(G)$  and  $g \in G$ , and then evaluate  $\chi(g)$ .

$$\text{Prob}(\chi(g) = 0) =$$

## Part 1. Character vanishing

**Burnside** Nonlinear irreducible characters have zeros.

**Question** What is the chance that  $\chi(g)$  equals 0?

**Two natural ways to pick a random character value  $\chi(g)$**

$S_4$	1	6	3	8	6
	id	(12)	(12)(34)	(123)	(1234)
$\chi_1$	1	1	1	1	1
$\chi_2$	3	1	-1	0	-1
$\chi_3$	2	0	2	-1	0
$\chi_4$	3	-1	-1	0	1
$\chi_5$	1	-1	1	1	-1

1. Choose  $\chi \in \text{Irr}(G)$  and  $g \in G$ , and then evaluate  $\chi(g)$ .

$$\text{Prob}(\chi(g) = 0) = \frac{|\{(\chi, g) \in \text{Irr}(G) \times G : \chi(g) = 0\}|}{|\text{Irr}(G) \times G|}$$

## Part 1. Character vanishing

**Burnside** Nonlinear irreducible characters have zeros.

**Question** What is the chance that  $\chi(g)$  equals 0?

**Two natural ways to pick a random character value  $\chi(g)$**

$S_4$	1	6	3	8	6
	id	(12)	(12)(34)	(123)	(1234)
$\chi_1$	1	1	1	1	1
$\chi_2$	3	1	-1	0	-1
$\chi_3$	2	0	2	-1	0
$\chi_4$	3	-1	-1	0	1
$\chi_5$	1	-1	1	1	-1

1. Choose  $\chi \in \text{Irr}(G)$  and  $g \in G$ , and then evaluate  $\chi(g)$ .

$$\text{Prob}(\chi(g) = 0) = \frac{|\{(\chi, g) \in \text{Irr}(G) \times G : \chi(g) = 0\}|}{|\text{Irr}(G) \times G|} =$$

## Part 1. Character vanishing

**Burnside** Nonlinear irreducible characters have zeros.

**Question** What is the chance that  $\chi(g)$  equals 0?

**Two natural ways to pick a random character value  $\chi(g)$**

$S_4$	1	6	3	8	6
	id	(12)	(12)(34)	(123)	(1234)
$\chi_1$	1	1	1	1	1
$\chi_2$	3	1	-1	0	-1
$\chi_3$	2	0	2	-1	0
$\chi_4$	3	-1	-1	0	1
$\chi_5$	1	-1	1	1	-1

1. Choose  $\chi \in \text{Irr}(G)$  and  $g \in G$ , and then evaluate  $\chi(g)$ .

$$\text{Prob}(\chi(g) = 0) = \frac{|\{(\chi, g) \in \text{Irr}(G) \times G : \chi(g) = 0\}|}{|\text{Irr}(G) \times G|} = \frac{28}{120} \approx 0.194$$

## Part 1. Character vanishing

**Burnside** Nonlinear irreducible characters have zeros.

**Question** What is the chance that  $\chi(g)$  equals 0?

**Two natural ways to pick a random character value  $\chi(g)$**

$S_4$	1	6	3	8	6
	id	(12)	(12)(34)	(123)	(1234)
$\chi_1$	1	1	1	1	1
$\chi_2$	3	1	-1	0	-1
$\chi_3$	2	0	2	-1	0
$\chi_4$	3	-1	-1	0	1
$\chi_5$	1	-1	1	1	-1

1. Choose  $\chi \in \text{Irr}(G)$  and  $g \in G$ , and then evaluate  $\chi(g)$ .

$$\text{Prob}(\chi(g) = 0) = \frac{|\{(\chi, g) \in \text{Irr}(G) \times G : \chi(g) = 0\}|}{|\text{Irr}(G) \times G|} = \frac{28}{120} \approx 0.194$$

- 2.

## Part 1. Character vanishing

**Burnside** Nonlinear irreducible characters have zeros.

**Question** What is the chance that  $\chi(g)$  equals 0?

**Two natural ways to pick a random character value  $\chi(g)$**

$S_4$	1	6	3	8	6
	id	(12)	(12)(34)	(123)	(1234)
$\chi_1$	1	1	1	1	1
$\chi_2$	3	1	-1	0	-1
$\chi_3$	2	0	2	-1	0
$\chi_4$	3	-1	-1	0	1
$\chi_5$	1	-1	1	1	-1

1. Choose  $\chi \in \text{Irr}(G)$  and  $g \in G$ , and then evaluate  $\chi(g)$ .

$$\text{Prob}(\chi(g) = 0) = \frac{|\{(\chi, g) \in \text{Irr}(G) \times G : \chi(g) = 0\}|}{|\text{Irr}(G) \times G|} = \frac{28}{120} \approx 0.194$$

2. Choose  $\chi \in \text{Irr}(G)$  and  $K \in \text{Cl}(G)$ , and then evaluate  $\chi(K)$ .

## Part 1. Character vanishing

**Burnside** Nonlinear irreducible characters have zeros.

**Question** What is the chance that  $\chi(g)$  equals 0?

**Two natural ways to pick a random character value  $\chi(g)$**

$S_4$	1	6	3	8	6
	id	(12)	(12)(34)	(123)	(1234)
$\chi_1$	1	1	1	1	1
$\chi_2$	3	1	-1	0	-1
$\chi_3$	2	0	2	-1	0
$\chi_4$	3	-1	-1	0	1
$\chi_5$	1	-1	1	1	-1

1. Choose  $\chi \in \text{Irr}(G)$  and  $g \in G$ , and then evaluate  $\chi(g)$ .

$$\text{Prob}(\chi(g) = 0) = \frac{|\{(\chi, g) \in \text{Irr}(G) \times G : \chi(g) = 0\}|}{|\text{Irr}(G) \times G|} = \frac{28}{120} \approx 0.194$$

2. Choose  $\chi \in \text{Irr}(G)$  and  $K \in \text{Cl}(G)$ , and then evaluate  $\chi(K)$ .

$$\text{Prob}(\chi(K) = 0)$$



## Part 1. Character vanishing

**Burnside** Nonlinear irreducible characters have zeros.

**Question** What is the chance that  $\chi(g)$  equals 0?

**Two natural ways to pick a random character value  $\chi(g)$**

$S_4$	1	6	3	8	6
	id	(12)	(12)(34)	(123)	(1234)
$\chi_1$	1	1	1	1	1
$\chi_2$	3	1	-1	0	-1
$\chi_3$	2	0	2	-1	0
$\chi_4$	3	-1	-1	0	1
$\chi_5$	1	-1	1	1	-1

1. Choose  $\chi \in \text{Irr}(G)$  and  $g \in G$ , and then evaluate  $\chi(g)$ .

$$\text{Prob}(\chi(g) = 0) = \frac{|\{(\chi, g) \in \text{Irr}(G) \times G : \chi(g) = 0\}|}{|\text{Irr}(G) \times G|} = \frac{28}{120} \approx 0.194$$

2. Choose  $\chi \in \text{Irr}(G)$  and  $K \in \text{Cl}(G)$ , and then evaluate  $\chi(K)$ .

$$\text{Prob}(\chi(K) = 0) =$$

## Part 1. Character vanishing

**Burnside** Nonlinear irreducible characters have zeros.

**Question** What is the chance that  $\chi(g)$  equals 0?

**Two natural ways to pick a random character value  $\chi(g)$**

$S_4$	1	6	3	8	6
	id	(12)	(12)(34)	(123)	(1234)
$\chi_1$	1	1	1	1	1
$\chi_2$	3	1	-1	0	-1
$\chi_3$	2	0	2	-1	0
$\chi_4$	3	-1	-1	0	1
$\chi_5$	1	-1	1	1	-1

1. Choose  $\chi \in \text{Irr}(G)$  and  $g \in G$ , and then evaluate  $\chi(g)$ .

$$\text{Prob}(\chi(g) = 0) = \frac{|\{(\chi, g) \in \text{Irr}(G) \times G : \chi(g) = 0\}|}{|\text{Irr}(G) \times G|} = \frac{28}{120} \approx 0.194$$

2. Choose  $\chi \in \text{Irr}(G)$  and  $K \in \text{Cl}(G)$ , and then evaluate  $\chi(K)$ .

$$\text{Prob}(\chi(K) = 0) = \frac{|\{(\chi, K) \in \text{Irr}(G) \times \text{Cl}(G) : \chi(K) = 0\}|}{|\text{Irr}(G) \times \text{Cl}(G)|}$$

## Part 1. Character vanishing

**Burnside** Nonlinear irreducible characters have zeros.

**Question** What is the chance that  $\chi(g)$  equals 0?

**Two natural ways to pick a random character value  $\chi(g)$**

$S_4$	1	6	3	8	6
	id	(12)	(12)(34)	(123)	(1234)
$\chi_1$	1	1	1	1	1
$\chi_2$	3	1	-1	0	-1
$\chi_3$	2	0	2	-1	0
$\chi_4$	3	-1	-1	0	1
$\chi_5$	1	-1	1	1	-1

1. Choose  $\chi \in \text{Irr}(G)$  and  $g \in G$ , and then evaluate  $\chi(g)$ .

$$\text{Prob}(\chi(g) = 0) = \frac{|\{(\chi, g) \in \text{Irr}(G) \times G : \chi(g) = 0\}|}{|\text{Irr}(G) \times G|} = \frac{28}{120} \approx 0.194$$

2. Choose  $\chi \in \text{Irr}(G)$  and  $K \in \text{Cl}(G)$ , and then evaluate  $\chi(K)$ .

$$\text{Prob}(\chi(K) = 0) = \frac{|\{(\chi, K) \in \text{Irr}(G) \times \text{Cl}(G) : \chi(K) = 0\}|}{|\text{Irr}(G) \times \text{Cl}(G)|} =$$

## Part 1. Character vanishing

**Burnside** Nonlinear irreducible characters have zeros.

**Question** What is the chance that  $\chi(g)$  equals 0?

**Two natural ways to pick a random character value  $\chi(g)$**

$S_4$	1	6	3	8	6
	id	(12)	(12)(34)	(123)	(1234)
$\chi_1$	1	1	1	1	1
$\chi_2$	3	1	-1	0	-1
$\chi_3$	2	0	2	-1	0
$\chi_4$	3	-1	-1	0	1
$\chi_5$	1	-1	1	1	-1

1. Choose  $\chi \in \text{Irr}(G)$  and  $g \in G$ , and then evaluate  $\chi(g)$ .

$$\text{Prob}(\chi(g) = 0) = \frac{|\{(\chi, g) \in \text{Irr}(G) \times G : \chi(g) = 0\}|}{|\text{Irr}(G) \times G|} = \frac{28}{120} \approx 0.194$$

2. Choose  $\chi \in \text{Irr}(G)$  and  $K \in \text{Cl}(G)$ , and then evaluate  $\chi(K)$ .

$$\text{Prob}(\chi(K) = 0) = \frac{|\{(\chi, K) \in \text{Irr}(G) \times \text{Cl}(G) : \chi(K) = 0\}|}{|\text{Irr}(G) \times \text{Cl}(G)|} = \frac{4}{5^2} = 0.16$$

$\text{Prob}(\chi(g) = 0)$  for  $S_n$

$\text{Prob}(\chi(g) = 0)$  for  $S_n$

**Theorem (M.)** If  $\chi \in \text{Irr}(S_n)$  and  $g \in S_n$  are chosen uniformly at random, then  $\chi(g) = 0$  with probability  $\rightarrow 1$  as  $n \rightarrow \infty$ .

$\text{Prob}(\chi(g) = 0)$  for  $S_n$

**Theorem (M.)** If  $\chi \in \text{Irr}(S_n)$  and  $g \in S_n$  are chosen uniformly at random, then  $\chi(g) = 0$  with probability  $\rightarrow 1$  as  $n \rightarrow \infty$ .

**One reason** A vanishingly small proportion of classes covers almost all of  $S_n$ .

$\text{Prob}(\chi(g) = 0)$  for  $S_n$

**Theorem (M.)** If  $\chi \in \text{Irr}(S_n)$  and  $g \in S_n$  are chosen uniformly at random, then  $\chi(g) = 0$  with probability  $\rightarrow 1$  as  $n \rightarrow \infty$ .

**One reason** A vanishingly small proportion of classes covers almost all of  $S_n$ .

**Lemma (M.)** For any  $\mathcal{K} \subseteq \text{Cl}(G)$ ,

$$\text{Prob}(\chi(g) = 0) \geq \frac{|\{g \in G : g^G \in \mathcal{K}\}|}{|G|} - \frac{|\mathcal{K}|}{|\text{Cl}(G)|}.$$



$\text{Prob}(\chi(g) = 0)$  and  $\text{Prob}(\chi(K) = 0)$  for some other groups

$\text{Prob}(\chi(g) = 0)$  and  $\text{Prob}(\chi(K) = 0)$  for some other groups

**Lemma (Gallagher–Larsen–M.)** For each finite group  $G$  and  $\epsilon > 0$ ,

$$\text{Prob}(\chi(g) \neq 0) \leq \frac{|\{(\chi, g) : \gcd(\chi(1), |g^G|) \geq \epsilon \chi(1)\}|}{|\text{Irr}(G) \times G|} + \epsilon^2.$$

$\text{Prob}(\chi(g) = 0)$  and  $\text{Prob}(\chi(K) = 0)$  for some other groups

**Lemma (Gallagher–Larsen–M.)** For each finite group  $G$  and  $\epsilon > 0$ ,

$$\text{Prob}(\chi(g) \neq 0) \leq \frac{|\{(\chi, g) : \gcd(\chi(1), |g^G|) \geq \epsilon \chi(1)\}|}{|\text{Irr}(G) \times G|} + \epsilon^2.$$

**Theorem (Gallagher–Larsen–M.)** For  $G = \text{GL}(n, q)$ , the proportion  $P_{n,q}$  of pairs  $(\chi, g) \in \text{Irr}(G) \times G$  with  $\chi(g) = 0$  satisfies

$$\inf_q P_{n,q} \rightarrow 1 \text{ as } n \rightarrow \infty.$$

So for any sequence of prime powers  $q_1, q_2, \dots$ , we have  $P_{n,q_n} \rightarrow 1$  as  $n \rightarrow \infty$ .

$\text{Prob}(\chi(g) = 0)$  and  $\text{Prob}(\chi(K) = 0)$  for some other groups

**Lemma (Gallagher–Larsen–M.)** For each finite group  $G$  and  $\epsilon > 0$ ,

$$\text{Prob}(\chi(g) \neq 0) \leq \frac{|\{(\chi, g) : \gcd(\chi(1), |g^G|) \geq \epsilon\chi(1)\}|}{|\text{Irr}(G) \times G|} + \epsilon^2.$$

**Theorem (Gallagher–Larsen–M.)** For  $G = \text{GL}(n, q)$ , the proportion  $P_{n,q}$  of pairs  $(\chi, g) \in \text{Irr}(G) \times G$  with  $\chi(g) = 0$  satisfies

$$\inf_q P_{n,q} \rightarrow 1 \text{ as } n \rightarrow \infty.$$

So for any sequence of prime powers  $q_1, q_2, \dots$ , we have  $P_{n,q_n} \rightarrow 1$  as  $n \rightarrow \infty$ .

**Theorem (Larsen–M.)** If  $G_n$  is any sequence of finite simple groups of Lie type with rank tending to  $\infty$ , then almost every entry in the character table of  $G_n$  is zero as  $n$  tends to  $\infty$ .

$\text{Prob}(\chi(K) = 0)$  for  $S_n$

**Question** What can be said about the limiting behavior of  $\text{Prob}(\chi_\lambda(\mu) = 0)$ ?

$\text{Prob}(\chi(K) = 0)$  for  $S_n$

**Question** What can be said about the limiting behavior of  $\text{Prob}(\chi_\lambda(\mu) = 0)$ ?

$$\frac{n}{\text{Prob}(\chi_\lambda(\mu) = 0)}$$

\_\_\_\_\_

$\text{Prob}(\chi(K) = 0)$  for  $S_n$

**Question** What can be said about the limiting behavior of  $\text{Prob}(\chi_\lambda(\mu) = 0)$ ?

$$\frac{n \quad \text{Prob}(\chi_\lambda(\mu) = 0)}{2}$$

---

$\text{Prob}(\chi(K) = 0)$  for  $S_n$

**Question** What can be said about the limiting behavior of  $\text{Prob}(\chi_\lambda(\mu) = 0)$ ?

$n$	$\text{Prob}(\chi_\lambda(\mu) = 0)$
2	0.0000

---



$\text{Prob}(\chi(K) = 0)$  for  $S_n$

**Question** What can be said about the limiting behavior of  $\text{Prob}(\chi_\lambda(\mu) = 0)$ ?

$n$	$\text{Prob}(\chi_\lambda(\mu) = 0)$
2	0.0000
3	

$\text{Prob}(\chi(K) = 0)$  for  $S_n$

**Question** What can be said about the limiting behavior of  $\text{Prob}(\chi_\lambda(\mu) = 0)$ ?

---

$n$	$\text{Prob}(\chi_\lambda(\mu) = 0)$
2	0.0000
3	0.1111

---

$\text{Prob}(\chi(K) = 0)$  for  $S_n$

**Question** What can be said about the limiting behavior of  $\text{Prob}(\chi_\lambda(\mu) = 0)$ ?

---

$n$	$\text{Prob}(\chi_\lambda(\mu) = 0)$
2	0.0000
3	0.1111
4	

---

$\text{Prob}(\chi(K) = 0)$  for  $S_n$

**Question** What can be said about the limiting behavior of  $\text{Prob}(\chi_\lambda(\mu) = 0)$ ?

---

$n$	$\text{Prob}(\chi_\lambda(\mu) = 0)$
2	0.0000
3	0.1111
4	0.1600

---

$\text{Prob}(\chi(K) = 0)$  for  $S_n$

**Question** What can be said about the limiting behavior of  $\text{Prob}(\chi_\lambda(\mu) = 0)$ ?

$n$	$\text{Prob}(\chi_\lambda(\mu) = 0)$
2	0.0000
3	0.1111
4	0.1600
5	0.2041

$\text{Prob}(\chi(K) = 0)$  for  $S_n$


**Question** What can be said about the limiting behavior of  $\text{Prob}(\chi_\lambda(\mu) = 0)$ ?

$n$	$\text{Prob}(\chi_\lambda(\mu) = 0)$
2	0.0000
3	0.1111
4	0.1600
5	0.2041
...	...
37	0.3642
38	0.3659

$\text{Prob}(\chi(K) = 0)$  for  $S_n$

**Question** What can be said about the limiting behavior of  $\text{Prob}(\chi_\lambda(\mu) = 0)$ ?


$n$	$\text{Prob}(\chi_\lambda(\mu) = 0)$
2	0.0000
3	0.1111
4	0.1600
5	0.2041
...	...
37	0.3642
38	0.3659



$\text{Prob}(\chi(K) = 0)$  for  $S_n$

**Question** What can be said about the limiting behavior of  $\text{Prob}(\chi_\lambda(\mu) = 0)$ ?

$n$	$\text{Prob}(\chi_\lambda(\mu) = 0)$
2	0.0000
3	0.1111
4	0.1600
5	0.2041
...	...
37	0.3642
38	0.3659



**Best known bound**  $\text{Prob}(\chi_\lambda(\mu) = 0) \geq C / \log n$



Part 2.

## Part 2. Modular vanishing for $S_n$

## Part 2. Modular vanishing for $S_n$

---

$n$

$$\text{Prob}(\chi_\lambda(\mu) \equiv 0 \pmod{2})$$

---

## Part 2. Modular vanishing for $S_n$

---

$n$	4
$\text{Prob}(\chi_\lambda(\mu) \equiv 0 \pmod{2})$	.24

---

## Part 2. Modular vanishing for $S_n$

---

$n$	4	5
$\text{Prob}(\chi_\lambda(\mu) \equiv 0 \pmod{2})$	.24	.33

---

## Part 2. Modular vanishing for $S_n$

---

$n$	4	5	6
$\text{Prob}(\chi_\lambda(\mu) \equiv 0 \pmod{2})$	.24	.33	.36

---

## Part 2. Modular vanishing for $S_n$

---

$n$	4	5	6	7
$\text{Prob}(\chi_\lambda(\mu) \equiv 0 \pmod{2})$	.24	.33	.36	.40

---

## Part 2. Modular vanishing for $S_n$

---

$n$	4	5	6	7	8
$\text{Prob}(\chi_\lambda(\mu) \equiv 0 \pmod{2})$	.24	.33	.36	.40	.55

---



## Part 2. Modular vanishing for $S_n$

$n$	4	5	6	7	8	9
$\text{Prob}(\chi_\lambda(\mu) \equiv 0 \pmod{2})$	.24	.33	.36	.40	.55	.56

## Part 2. Modular vanishing for $S_n$

$n$	4	5	6	7	8	9	10
$\text{Prob}(\chi_\lambda(\mu) \equiv 0 \pmod{2})$	.24	.33	.36	.40	.55	.56	.55

## Part 2. Modular vanishing for $S_n$


$n$	4	5	6	7	8	9	10	...
$\text{Prob}(\chi_\lambda(\mu) \equiv 0 \pmod{2})$	.24	.33	.36	.40	.55	.56	.55	

## Part 2. Modular vanishing for $S_n$

$n$	4	5	6	7	8	9	10	...	76
$\text{Prob}(\chi_\lambda(\mu) \equiv 0 \pmod{2})$	.24	.33	.36	.40	.55	.56	.55		.87


## Part 2. Modular vanishing for $S_n$

$n$	4	5	6	7	8	9	10	...	76
$\text{Prob}(\chi_\lambda(\mu) = 0 \bmod 2)$	.24	.33	.36	.40	.55	.56	.55		.87



## Part 2. Modular vanishing for $S_n$


$n$	4	5	6	7	8	9	10	...	76
$\text{Prob}(\chi_\lambda(\mu) \equiv 0 \pmod{2})$	.24	.33	.36	.40	.55	.56	.55		.87



- The character table of  $S_{76}$  has about 86 trillion entries.

## Part 2. Modular vanishing for $S_n$


$n$	4	5	6	7	8	9	10	...	76
$\text{Prob}(\chi_\lambda(\mu) = 0 \bmod 2)$	.24	.33	.36	.40	.55	.56	.55		.87



- The character table of  $S_{76}$  has about 86 trillion entries.
- Several months of computations on super computers used for earth science.

## Part 2. Modular vanishing for $S_n$

$n$	4	5	6	7	8	9	10	...	76
$\text{Prob}(\chi_\lambda(\mu) = 0 \bmod 2)$	.24	.33	.36	.40	.55	.56	.55		.87




- The character table of  $S_{76}$  has about 86 trillion entries.
- Several months of computations on super computers used for earth science.
- Similar computations for other primes and prime powers.



## Part 2. Modular vanishing for $S_n$

$n$	4	5	6	7	8	9	10	...	76
$\text{Prob}(\chi_\lambda(\mu) \equiv 0 \pmod{2})$	.24	.33	.36	.40	.55	.56	.55		.87




- The character table of  $S_{76}$  has about 86 trillion entries.
- Several months of computations on super computers used for earth science.
- Similar computations for other primes and prime powers.

### Conjecture (M.)

Almost every entry in the character table of  $S_n$  is divisible by any fixed prime  $p$  as  $n \rightarrow \infty$ .

## Part 2. Modular vanishing for $S_n$

$n$	4	5	6	7	8	9	10	...	76
$\text{Prob}(\chi_\lambda(\mu) \equiv 0 \pmod{2})$	.24	.33	.36	.40	.55	.56	.55		.87




- The character table of  $S_{76}$  has about 86 trillion entries.
- Several months of computations on super computers used for earth science.
- Similar computations for other primes and prime powers.

### Conjecture (M.)

Almost every entry in the character table of  $S_n$  is divisible by any fixed prime  $p$  as  $n \rightarrow \infty$ . In other words,  $\text{Prob}(\chi_\lambda(\mu) \equiv 0 \pmod{p}) \rightarrow 1$  as  $n \rightarrow \infty$ .

## Part 2. Modular vanishing for $S_n$

$n$	4	5	6	7	8	9	10	...	76
$\text{Prob}(\chi_\lambda(\mu) \equiv 0 \pmod{2})$	.24	.33	.36	.40	.55	.56	.55		.87



- The character table of  $S_{76}$  has about 86 trillion entries.
- Several months of computations on super computers used for earth science.
- Similar computations for other primes and prime powers.

### Conjecture (M.)

Almost every entry in the character table of  $S_n$  is divisible by any fixed prime  $p$  as  $n \rightarrow \infty$ . In other words,  $\text{Prob}(\chi_\lambda(\mu) \equiv 0 \pmod{p}) \rightarrow 1$  as  $n \rightarrow \infty$ .

### Conjecture (M.)

Almost every entry in the character table of  $S_n$  is divisible by any fixed  $m$  as  $n \rightarrow \infty$ .

## Part 2. Modular vanishing for $S_n$

### **Conjecture (M.)**

Almost every entry in the character table of  $S_n$  is divisible by any fixed prime.

### **Conjecture (M.)**

Almost every entry in the character table of  $S_n$  is divisible by any fixed integer.

## Part 2. Modular vanishing for $S_n$

### **Conjecture (M.)**

Almost every entry in the character table of  $S_n$  is divisible by any fixed prime.

### **Conjecture (M.)**

Almost every entry in the character table of  $S_n$  is divisible by any fixed integer.

**Lemma**  $\chi_\lambda(\mu) \equiv \chi_\lambda(\nu) \pmod{p}$  if  $\nu$  is obtained by joining  $p$  equal parts in  $\mu$ .

## Part 2. Modular vanishing for $S_n$

### **Conjecture (M.)**

Almost every entry in the character table of  $S_n$  is divisible by any fixed prime.

### **Conjecture (M.)**

Almost every entry in the character table of  $S_n$  is divisible by any fixed integer.

**Lemma**  $\chi_\lambda(\mu) \equiv \chi_\lambda(\nu) \pmod{p}$  if  $\nu$  is obtained by joining  $p$  equal parts in  $\mu$ .

**Example** For  $p = 2$ ,  $\chi_\lambda(42111)$

## Part 2. Modular vanishing for $S_n$

### **Conjecture (M.)**

Almost every entry in the character table of  $S_n$  is divisible by any fixed prime.

### **Conjecture (M.)**

Almost every entry in the character table of  $S_n$  is divisible by any fixed integer.

**Lemma**  $\chi_\lambda(\mu) \equiv \chi_\lambda(\nu) \pmod{p}$  if  $\nu$  is obtained by joining  $p$  equal parts in  $\mu$ .

**Example** For  $p = 2$ ,  $\chi_\lambda(42111) \equiv \chi_\lambda(4221)$  .

## Part 2. Modular vanishing for $S_n$

### **Conjecture (M.)**

Almost every entry in the character table of  $S_n$  is divisible by any fixed prime.

### **Conjecture (M.)**

Almost every entry in the character table of  $S_n$  is divisible by any fixed integer.

**Lemma**  $\chi_\lambda(\mu) \equiv \chi_\lambda(\nu) \pmod{p}$  if  $\nu$  is obtained by joining  $p$  equal parts in  $\mu$ .

**Example** For  $p = 2$ ,  $\chi_\lambda(42111) \equiv \chi_\lambda(4221) \equiv \chi_\lambda(441) \pmod{2}$ .



## Part 2. Modular vanishing for $S_n$

### **Conjecture (M.)**

Almost every entry in the character table of  $S_n$  is divisible by any fixed prime.

### **Conjecture (M.)**

Almost every entry in the character table of  $S_n$  is divisible by any fixed integer.

**Lemma**  $\chi_\lambda(\mu) \equiv \chi_\lambda(\nu) \pmod{p}$  if  $\nu$  is obtained by joining  $p$  equal parts in  $\mu$ .

**Example** For  $p = 2$ ,  $\chi_\lambda(42111) \equiv \chi_\lambda(4221) \equiv \chi_\lambda(441) \equiv \chi_\lambda(81) \pmod{2}$ .

## Part 2. Modular vanishing for $S_n$

### **Conjecture (M.)**

Almost every entry in the character table of  $S_n$  is divisible by any fixed prime.

### **Conjecture (M.)**

Almost every entry in the character table of  $S_n$  is divisible by any fixed integer.

**Lemma**  $\chi_\lambda(\mu) \equiv \chi_\lambda(\nu) \pmod{p}$  if  $\nu$  is obtained by joining  $p$  equal parts in  $\mu$ .

**Example** For  $p = 2$ ,  $\chi_\lambda(42111) \equiv \chi_\lambda(4221) \equiv \chi_\lambda(441) \equiv \chi_\lambda(81) \pmod{2}$ .

## Part 2. Modular vanishing for $S_n$

### **Conjecture (M.)**

Almost every entry in the character table of  $S_n$  is divisible by any fixed prime.

### **Conjecture (M.)**

Almost every entry in the character table of  $S_n$  is divisible by any fixed integer.

**Lemma**  $\chi_\lambda(\mu) \equiv \chi_\lambda(\nu) \pmod{p}$  if  $\nu$  is obtained by joining  $p$  equal parts in  $\mu$ .

**Example** For  $p = 2$ ,  $\chi_\lambda(42111) \equiv \chi_\lambda(4221) \equiv \chi_\lambda(441) \equiv \chi_\lambda(81) \pmod{2}$ .

**Corollary**  $\chi_\lambda(\mu) \equiv 0 \pmod{p}$  if  $\lambda$  is a  $\mathfrak{l}_p(\mu)$ -core.

## Part 2. Modular vanishing for $S_n$

### Conjecture (M.)

Almost every entry in the character table of  $S_n$  is divisible by any fixed prime.

### Conjecture (M.)

Almost every entry in the character table of  $S_n$  is divisible by any fixed integer.

**Lemma**  $\chi_\lambda(\mu) \equiv \chi_\lambda(\nu) \pmod{p}$  if  $\nu$  is obtained by joining  $p$  equal parts in  $\mu$ .

**Example** For  $p = 2$ ,  $\chi_\lambda(42111) \equiv \chi_\lambda(4221) \equiv \chi_\lambda(441) \equiv \chi_\lambda(81) \pmod{2}$ .

**Corollary**  $\chi_\lambda(\mu) \equiv 0 \pmod{p}$  if  $\lambda$  is a  $\mathfrak{l}_p(\mu)$ -core.

$n$	30	40	50	60	70	80	90	100	110	120
Prob( $\lambda$ is an $\mathfrak{l}_2(\mu)$ -core)	.45	.44	.45	.47	.48	.48	.48	.47	.47	.46

## Part 2. Modular vanishing for $S_n$

### Conjecture (M.)

Almost every entry in the character table of  $S_n$  is divisible by any fixed prime.

### Conjecture (M.)

Almost every entry in the character table of  $S_n$  is divisible by any fixed integer.

**Lemma**  $\chi_\lambda(\mu) \equiv \chi_\lambda(\nu) \pmod{p}$  if  $\nu$  is obtained by joining  $p$  equal parts in  $\mu$ .

**Example** For  $p = 2$ ,  $\chi_\lambda(42111) \equiv \chi_\lambda(4221) \equiv \chi_\lambda(441) \equiv \chi_\lambda(81) \pmod{2}$ .

**Corollary**  $\chi_\lambda(\mu) \equiv 0 \pmod{p}$  if  $\lambda$  is a  $\mathfrak{l}_p(\mu)$ -core.

$n$	30	40	50	60	70	80	90	100	110	120
Prob( $\lambda$ is an $\mathfrak{l}_2(\mu)$ -core)	.45	.44	.45	.47	.48	.48	.48	.47	.47	.46

**Partial results** M., Gluck, Morotti, Ono, McSpirt, Harman, Peluse, Soundararajan, . . .

## Part 2. Modular vanishing for $S_n$

### Conjecture (M.)

Almost every entry in the character table of  $S_n$  is divisible by any fixed prime.

### Conjecture (M.)

Almost every entry in the character table of  $S_n$  is divisible by any fixed integer.

**Lemma**  $\chi_\lambda(\mu) \equiv \chi_\lambda(\nu) \pmod{p}$  if  $\nu$  is obtained by joining  $p$  equal parts in  $\mu$ .

**Example** For  $p = 2$ ,  $\chi_\lambda(42111) \equiv \chi_\lambda(4221) \equiv \chi_\lambda(441) \equiv \chi_\lambda(81) \pmod{2}$ .

**Corollary**  $\chi_\lambda(\mu) \equiv 0 \pmod{p}$  if  $\lambda$  is a  $\mathfrak{l}_p(\mu)$ -core.

$n$	30	40	50	60	70	80	90	100	110	120
Prob( $\lambda$ is an $\mathfrak{l}_2(\mu)$ -core)	.45	.44	.45	.47	.48	.48	.48	.47	.47	.46

**Partial results** M., Gluck, Morotti, Ono, McSpirt, Harman, Peluse, Soundararajan, . . .

**Lemma (Morotti)**  $\#\{\text{partitions of } n \text{ that are not } t\text{-cores}\} \leq (t+1)p(n-t)$ .

## Part 2. Modular vanishing for $S_n$

### Conjecture (M.)

Almost every entry in the character table of  $S_n$  is divisible by any fixed prime.

### Conjecture (M.)

Almost every entry in the character table of  $S_n$  is divisible by any fixed integer.

**Lemma**  $\chi_\lambda(\mu) \equiv \chi_\lambda(\nu) \pmod{p}$  if  $\nu$  is obtained by joining  $p$  equal parts in  $\mu$ .

**Example** For  $p = 2$ ,  $\chi_\lambda(42111) \equiv \chi_\lambda(4221) \equiv \chi_\lambda(441) \equiv \chi_\lambda(81) \pmod{2}$ .

**Corollary**  $\chi_\lambda(\mu) \equiv 0 \pmod{p}$  if  $\lambda$  is a  $\mathfrak{l}_p(\mu)$ -core.

$n$	30	40	50	60	70	80	90	100	110	120
Prob( $\lambda$ is an $\mathfrak{l}_2(\mu)$ -core)	.45	.44	.45	.47	.48	.48	.48	.47	.47	.46

**Partial results** M., Gluck, Morotti, Ono, McSpirt, Harman, Peluse, Soundararajan, . . .

**Lemma (Morotti)**  $\#\{\text{partitions of } n \text{ that are not } t\text{-cores}\} \leq (t+1)p(n-t)$ .

### Theorem (Peluse–Soundararajan)

Almost every entry in the character table of  $S_n$  is divisible by any fixed prime.

## Part 2. Modular vanishing for $S_n$

### Conjecture (M.)

Almost every entry in the character table of  $S_n$  is divisible by any fixed prime.

### Conjecture (M.)

Almost every entry in the character table of  $S_n$  is divisible by any fixed integer.

**Lemma**  $\chi_\lambda(\mu) \equiv \chi_\lambda(\nu) \pmod{p}$  if  $\nu$  is obtained by joining  $p$  equal parts in  $\mu$ .

**Example** For  $p = 2$ ,  $\chi_\lambda(42111) \equiv \chi_\lambda(4221) \equiv \chi_\lambda(441) \equiv \chi_\lambda(81) \pmod{2}$ .

**Corollary**  $\chi_\lambda(\mu) \equiv 0 \pmod{p}$  if  $\lambda$  is a  $\mathfrak{l}_p(\mu)$ -core.

$n$	30	40	50	60	70	80	90	100	110	120
Prob( $\lambda$ is an $\mathfrak{l}_2(\mu)$ -core)	.45	.44	.45	.47	.48	.48	.48	.47	.47	.46

**Partial results** M., Gluck, Morotti, Ono, McSpirt, Harman, Peluse, Soundararajan, . . .

**Lemma (Morotti)**  $\#\{\text{partitions of } n \text{ that are not } t\text{-cores}\} \leq (t+1)p(n-t)$ .

### Theorem (Peluse–Soundararajan)

Almost every entry in the character table of  $S_n$  is divisible by any fixed prime.

- Progress is being made on prime powers by various groups.



## Part 3. Two variations

### Part 3. Two variations

Let  $P(G)$  be either  $\text{Prob}(\chi(g) = 0)$  or  $\text{Prob}(\chi(K) = 0)$ .

### Part 3. Two variations

Let  $P(G)$  be either  $\text{Prob}(\chi(g) = 0)$  or  $\text{Prob}(\chi(K) = 0)$ .

1. Treat  $P(G)$  as a random variable itself.

### Part 3. Two variations

Let  $P(G)$  be either  $\text{Prob}(\chi(g) = 0)$  or  $\text{Prob}(\chi(K) = 0)$ .

1. Treat  $P(G)$  as a random variable itself.

**Theorem (M.)** The expected value of  $P(S_\lambda)$  tends to 1 as  $n \rightarrow \infty$ .

### Part 3. Two variations

Let  $P(G)$  be either  $\text{Prob}(\chi(g) = 0)$  or  $\text{Prob}(\chi(K) = 0)$ .

1. Treat  $P(G)$  as a random variable itself.

**Theorem (M.)** The expected value of  $P(S_\lambda)$  tends to 1 as  $n \rightarrow \infty$ .

2.

### Part 3. Two variations

Let  $P(G)$  be either  $\text{Prob}(\chi(g) = 0)$  or  $\text{Prob}(\chi(K) = 0)$ .

1. Treat  $P(G)$  as a random variable itself.

**Theorem (M.)** The expected value of  $P(S_\lambda)$  tends to 1 as  $n \rightarrow \infty$ .

2.  $G_1 < G_2 < \dots$

### Part 3. Two variations

Let  $P(G)$  be either  $\text{Prob}(\chi(g) = 0)$  or  $\text{Prob}(\chi(K) = 0)$ .

1. Treat  $P(G)$  as a random variable itself.

**Theorem (M.)** The expected value of  $P(S_\lambda)$  tends to 1 as  $n \rightarrow \infty$ .

2.  $G_1 < G_2 < \dots \longrightarrow P(G_1), P(G_2), \dots \in [0, 1]$

### Part 3. Two variations

Let  $P(G)$  be either  $\text{Prob}(\chi(g) = 0)$  or  $\text{Prob}(\chi(K) = 0)$ .

1. Treat  $P(G)$  as a random variable itself.

**Theorem (M.)** The expected value of  $P(S_\lambda)$  tends to 1 as  $n \rightarrow \infty$ .

2.  $G_1 < G_2 < \dots \longrightarrow P(G_1), P(G_2), \dots \in [0, 1]$   
 $a_1, a_2, \dots$



### Part 3. Two variations

Let  $P(G)$  be either  $\text{Prob}(\chi(g) = 0)$  or  $\text{Prob}(\chi(K) = 0)$ .

1. Treat  $P(G)$  as a random variable itself.

**Theorem (M.)** The expected value of  $P(S_\lambda)$  tends to 1 as  $n \rightarrow \infty$ .

2.  $G_1 < G_2 < \dots \longrightarrow P(G_1), P(G_2), \dots \in [0, 1]$   
          ?                     $\longleftarrow a_1, a_2, \dots$

### Part 3. Two variations

Let  $P(G)$  be either  $\text{Prob}(\chi(g) = 0)$  or  $\text{Prob}(\chi(K) = 0)$ .

1. Treat  $P(G)$  as a random variable itself.

**Theorem (M.)** The expected value of  $P(S_\lambda)$  tends to 1 as  $n \rightarrow \infty$ .

2.  $G_1 < G_2 < \dots \longrightarrow P(G_1), P(G_2), \dots \in [0, 1]$   
                  ?                    $\longleftarrow a_1, a_2, \dots$

**Theorem (M.)** If  $a_1, a_2, \dots \in [0, 1]$  and  $\epsilon_1, \epsilon_2, \dots \in (0, \infty)$ , then for each prime  $p$  there exists a chain of  $p$ -groups  $G_1 < G_2 < \dots$  such that, for each  $i$ ,

$$|P(G_i) - a_i| < \epsilon_i.$$

### Part 3. Two variations

Let  $P(G)$  be either  $\text{Prob}(\chi(g) = 0)$  or  $\text{Prob}(\chi(K) = 0)$ .

1. Treat  $P(G)$  as a random variable itself.

**Theorem (M.)** The expected value of  $P(S_\lambda)$  tends to 1 as  $n \rightarrow \infty$ .

2.  $G_1 < G_2 < \dots \longrightarrow P(G_1), P(G_2), \dots \in [0, 1]$   
                  ?                    $\longleftarrow a_1, a_2, \dots$

**Theorem (M.)** If  $a_1, a_2, \dots \in [0, 1]$  and  $\epsilon_1, \epsilon_2, \dots \in (0, \infty)$ , then for each prime  $p$  there exists a chain of  $p$ -groups  $G_1 < G_2 < \dots$  such that, for each  $i$ ,

$$|P(G_i) - a_i| < \epsilon_i.$$

In particular, the set  $\{P(G) : |G| < \infty\}$  is dense in  $[0, 1]$ .

## Part 4. Zeros and roots of unity

## Part 4. Zeros and roots of unity

	1 id	6 (12)	3 (12)(34)	8 (123)	6 (1234)
$\chi_1$	1	1	1	1	1
$\chi_2$	3	1	-1	0	-1
$\chi_3$	2	0	2	-1	0
$\chi_4$	3	-1	-1	0	1
$\chi_5$	1	-1	1	1	-1

## Part 4. Zeros and roots of unity

	1 id	6 (12)	3 (12)(34)	8 (123)	6 (1234)	24/24
$\chi_1$	1	1	1	1	1	
$\chi_2$	3	1	-1	0	-1	
$\chi_3$	2	0	2	-1	0	
$\chi_4$	3	-1	-1	0	1	
$\chi_5$	1	-1	1	1	-1	

## Part 4. Zeros and roots of unity

	1	6	3	8	6	
	id	(12)	(12)(34)	(123)	(1234)	
$\chi_1$	1	1	1	1	1	24/24
$\chi_2$	3	1	-1	0	-1	23/24
$\chi_3$	2	0	2	-1	0	
$\chi_4$	3	-1	-1	0	1	
$\chi_5$	1	-1	1	1	-1	

## Part 4. Zeros and roots of unity

	1	6	3	8	6	
	id	(12)	(12)(34)	(123)	(1234)	
$\chi_1$	1	1	1	1	1	24/24
$\chi_2$	3	1	-1	0	-1	23/24
$\chi_3$	2	0	2	-1	0	20/24
$\chi_4$	3	-1	-1	0	1	
$\chi_5$	1	-1	1	1	-1	



## Part 4. Zeros and roots of unity

	1	6	3	8	6	
	id	(12)	(12)(34)	(123)	(1234)	
$\chi_1$	1	1	1	1	1	24/24
$\chi_2$	3	1	-1	0	-1	23/24
$\chi_3$	2	0	2	-1	0	20/24
$\chi_4$	3	-1	-1	0	1	23/24
$\chi_5$	1	-1	1	1	-1	

## Part 4. Zeros and roots of unity

	1	6	3	8	6	
	id	(12)	(12)(34)	(123)	(1234)	
$\chi_1$	1	1	1	1	1	24/24
$\chi_2$	3	1	-1	0	-1	23/24
$\chi_3$	2	0	2	-1	0	20/24
$\chi_4$	3	-1	-1	0	1	23/24
$\chi_5$	1	-1	1	1	-1	24/24

## Part 4. Zeros and roots of unity

	1	6	3	8	6	
	id	(12)	(12)(34)	(123)	(1234)	
$\chi_1$	1	1	1	1	1	24/24
$\chi_2$	3	1	-1	0	-1	23/24
$\chi_3$	2	0	2	-1	0	20/24
$\chi_4$	3	-1	-1	0	1	23/24
$\chi_5$	1	-1	1	1	-1	24/24
		5/5				

## Part 4. Zeros and roots of unity

	1	6	3	8	6	
	id	(12)	(12)(34)	(123)	(1234)	
$\chi_1$	1	1	1	1	1	24/24
$\chi_2$	3	1	-1	0	-1	23/24
$\chi_3$	2	0	2	-1	0	20/24
$\chi_4$	3	-1	-1	0	1	23/24
$\chi_5$	1	-1	1	1	-1	24/24
		5/5		5/5		

## Part 4. Zeros and roots of unity

	1	6	3	8	6	
	id	(12)	(12)(34)	(123)	(1234)	
$\chi_1$	1	1	1	1	1	24/24
$\chi_2$	3	1	-1	0	-1	23/24
$\chi_3$	2	0	2	-1	0	20/24
$\chi_4$	3	-1	-1	0	1	23/24
$\chi_5$	1	-1	1	1	-1	24/24
		5/5		5/5	5/5	

## Part 4. Zeros and roots of unity

	1 id	6 (12)	3 (12)(34)	8 (123)	6 (1234)	
$\chi_1$	1	1	1	1	1	24/24
$\chi_2$	3	1	-1	0	-1	23/24
$\chi_3$	2	0	2	-1	0	20/24
$\chi_4$	3	-1	-1	0	1	23/24
$\chi_5$	1	-1	1	1	-1	24/24
		5/5		5/5	5/5	

### Definition

$$\theta(G) = \min_{\chi \in \text{Irr}(G)} \frac{\#\{g \in G : \chi(g) \text{ is 0 or a root of unity}\}}{|G|}$$

$$\theta'(G) = \min_{\text{L.T.A. } K} \frac{\#\{\chi \in \text{Irr}(G) : \chi(K) \text{ is 0 or a root of unity}\}}{|\text{Cl}(G)|}$$

## Part 4. Zeros and roots of unity

	1 id	6 (12)	3 (12)(34)	8 (123)	6 (1234)	
$\chi_1$	1	1	1	1	1	24/24
$\chi_2$	3	1	-1	0	-1	23/24
$\chi_3$	2	0	2	-1	0	20/24
$\chi_4$	3	-1	-1	0	1	23/24
$\chi_5$	1	-1	1	1	-1	24/24
		5/5		5/5	5/5	

### Definition

$$\theta(G) = \min_{\chi \in \text{Irr}(G)} \frac{\#\{g \in G : \chi(g) \text{ is 0 or a root of unity}\}}{|G|}$$

$$\theta'(G) = \min_{\text{L.T.A. } K} \frac{\#\{\chi \in \text{Irr}(G) : \chi(K) \text{ is 0 or a root of unity}\}}{|\text{Cl}(G)|}$$

**Example**  $\theta(S_4) = 20/24$

## Part 4. Zeros and roots of unity

	1 id	6 (12)	3 (12)(34)	8 (123)	6 (1234)	
$\chi_1$	1	1	1	1	1	24/24
$\chi_2$	3	1	-1	0	-1	23/24
$\chi_3$	2	0	2	-1	0	20/24
$\chi_4$	3	-1	-1	0	1	23/24
$\chi_5$	1	-1	1	1	-1	24/24
		5/5		5/5	5/5	

### Definition

$$\theta(G) = \min_{\chi \in \text{Irr}(G)} \frac{\#\{g \in G : \chi(g) \text{ is 0 or a root of unity}\}}{|G|}$$

$$\theta'(G) = \min_{\text{L.T.A. } K} \frac{\#\{\chi \in \text{Irr}(G) : \chi(K) \text{ is 0 or a root of unity}\}}{|\text{Cl}(G)|}$$

**Example**  $\theta(S_4) = 20/24$  and  $\theta'(S_4) = 1$ .



## Part 4. Zeros and roots of unity

	1	6	3	8	6	
	id	(12)	(12)(34)	(123)	(1234)	
$\chi_1$	1	1	1	1	1	24/24
$\chi_2$	3	1	-1	0	-1	23/24
$\chi_3$	2	0	2	-1	0	20/24
$\chi_4$	3	-1	-1	0	1	23/24
$\chi_5$	1	-1	1	1	-1	24/24
		5/5		5/5	5/5	

### Definition

$$\theta(G) = \min_{\chi \in \text{Irr}(G)} \frac{\#\{g \in G : \chi(g) \text{ is 0 or a root of unity}\}}{|G|}$$

$$\theta'(G) = \min_{\text{L.T.A. } K} \frac{\#\{\chi \in \text{Irr}(G) : \chi(K) \text{ is 0 or a root of unity}\}}{|\text{Cl}(G)|}$$

**Example**  $\theta(S_4) = 20/24$  and  $\theta'(S_4) = 1$ .

**Theorem (J.G. Thompson)**  $\theta(G) > 1/3$ .

## Part 4. Zeros and roots of unity

	1 id	6 (12)	3 (12)(34)	8 (123)	6 (1234)	
$\chi_1$	1	1	1	1	1	24/24
$\chi_2$	3	1	-1	0	-1	23/24
$\chi_3$	2	0	2	-1	0	20/24
$\chi_4$	3	-1	-1	0	1	23/24
$\chi_5$	1	-1	1	1	-1	24/24
		5/5		5/5	5/5	

### Definition

$$\theta(G) = \min_{\chi \in \text{Irr}(G)} \frac{\#\{g \in G : \chi(g) \text{ is 0 or a root of unity}\}}{|G|}$$

$$\theta'(G) = \min_{\text{L.T.A. } K} \frac{\#\{\chi \in \text{Irr}(G) : \chi(K) \text{ is 0 or a root of unity}\}}{|\text{Cl}(G)|}$$

**Example**  $\theta(S_4) = 20/24$  and  $\theta'(S_4) = 1$ .

**Theorem (J.G. Thompson)**  $\theta(G) > 1/3$ .

**Theorem (P.X. Gallagher)**  $\theta'(G) > 1/3$ .

## Part 4. Zeros and roots of unity

	1	6	3	8	6	
	id	(12)	(12)(34)	(123)	(1234)	
$\chi_1$	1	1	1	1	1	24/24
$\chi_2$	3	1	-1	0	-1	23/24
$\chi_3$	2	0	2	-1	0	20/24
$\chi_4$	3	-1	-1	0	1	23/24
$\chi_5$	1	-1	1	1	-1	24/24
		5/5		5/5	5/5	

### Definition

$$\theta(G) = \min_{\chi \in \text{Irr}(G)} \frac{\#\{g \in G : \chi(g) \text{ is 0 or a root of unity}\}}{|G|}$$

$$\theta'(G) = \min_{\text{L.T.A. } K} \frac{\#\{\chi \in \text{Irr}(G) : \chi(K) \text{ is 0 or a root of unity}\}}{|\text{Cl}(G)|}$$

**Example**  $\theta(S_4) = 20/24$  and  $\theta'(S_4) = 1$ .

**Theorem (J.G. Thompson)**  $\theta(G) > 1/3$ .

**Theorem (P.X. Gallagher)**  $\theta'(G) > 1/3$ .

**Theorem (C.L. Siegel)** For totally positive  $\alpha \neq 0, 1$ ,  $\tilde{\text{Tr}}(\alpha) \geq 3/2$ .  
(Siegel gets applied to  $\alpha = |\chi(g)|^2$ . Burnside used  $\tilde{\text{Tr}}(\alpha) \geq 1$  for  $\alpha \neq 0$ .)

**Theorem (Thompson)**  $\theta(G) > 1/3$ .

**Theorem (Gallagher)**  $\theta'(G) > 1/3$ .

**Theorem (Thompson)**  $\theta(G) > 1/3$ .

**Theorem (Gallagher)**  $\theta'(G) > 1/3$ .

**Question** What are the greatest lower bounds:  $\inf_G \theta(G)$ ,  $\inf_G \theta'(G)$  ?

**Theorem (Thompson)**  $\theta(G) > 1/3$ .

**Theorem (Gallagher)**  $\theta'(G) > 1/3$ .

**Question** What are the greatest lower bounds:  $\inf_G \theta(G)$ ,  $\inf_G \theta'(G)$  ?

**Conjecture (M.)**  $\inf \theta(G), \inf \theta'(G) = 1/2$ .

**Theorem (Thompson)**  $\theta(G) > 1/3$ .

**Theorem (Gallagher)**  $\theta'(G) > 1/3$ .

**Question** What are the greatest lower bounds:  $\inf_G \theta(G)$ ,  $\inf_G \theta'(G)$  ?

**Conjecture (M.)**  $\inf \theta(G), \inf \theta'(G) = 1/2$ .

**Proposition (M.)** For  $G_n = \text{Suz}(2^{2n+1})$ , we have  $\theta(G_n), \theta'(G_n) \rightarrow 1/2$ .

**Theorem (Thompson)**  $\theta(G) > 1/3$ .

**Theorem (Gallagher)**  $\theta'(G) > 1/3$ .

**Question** What are the greatest lower bounds:  $\inf_G \theta(G)$ ,  $\inf_G \theta'(G)$  ?

**Conjecture (M.)**  $\inf \theta(G), \inf \theta'(G) = 1/2$ .

**Proposition (M.)** For  $G_n = \text{Suz}(2^{2n+1})$ , we have  $\theta(G_n), \theta'(G_n) \rightarrow 1/2$ .

**Theorem (M.)** The conjecture holds for:



**Theorem (Thompson)**  $\theta(G) > 1/3$ .

**Theorem (Gallagher)**  $\theta'(G) > 1/3$ .

**Question** What are the greatest lower bounds:  $\inf_G \theta(G)$ ,  $\inf_G \theta'(G)$  ?

**Conjecture (M.)**  $\inf \theta(G), \inf \theta'(G) = 1/2$ .

**Proposition (M.)** For  $G_n = \text{Suz}(2^{2n+1})$ , we have  $\theta(G_n), \theta'(G_n) \rightarrow 1/2$ .

**Theorem (M.)** The conjecture holds for:

- Finite groups of order  $< 2^9$ .

**Theorem (Thompson)**  $\theta(G) > 1/3$ .

**Theorem (Gallagher)**  $\theta'(G) > 1/3$ .

**Question** What are the greatest lower bounds:  $\inf_G \theta(G)$ ,  $\inf_G \theta'(G)$  ?

**Conjecture (M.)**  $\inf \theta(G), \inf \theta'(G) = 1/2$ .

**Proposition (M.)** For  $G_n = \text{Suz}(2^{2n+1})$ , we have  $\theta(G_n), \theta'(G_n) \rightarrow 1/2$ .

**Theorem (M.)** The conjecture holds for:

- Finite groups of order  $< 2^9$ .
- Simple groups of order  $\leq 10^9$ .

**Theorem (Thompson)**  $\theta(G) > 1/3$ .

**Theorem (Gallagher)**  $\theta'(G) > 1/3$ .

**Question** What are the greatest lower bounds:  $\inf_G \theta(G)$ ,  $\inf_G \theta'(G)$  ?

**Conjecture (M.)**  $\inf \theta(G), \inf \theta'(G) = 1/2$ .

**Proposition (M.)** For  $G_n = \text{Suz}(2^{2n+1})$ , we have  $\theta(G_n), \theta'(G_n) \rightarrow 1/2$ .

**Theorem (M.)** The conjecture holds for:

- Finite groups of order  $< 2^9$ .
- Simple groups of order  $\leq 10^9$ .
- All sporadic groups.

**Theorem (Thompson)**  $\theta(G) > 1/3$ .

**Theorem (Gallagher)**  $\theta'(G) > 1/3$ .

**Question** What are the greatest lower bounds:  $\inf_G \theta(G)$ ,  $\inf_G \theta'(G)$  ?

**Conjecture (M.)**  $\inf \theta(G), \inf \theta'(G) = 1/2$ .

**Proposition (M.)** For  $G_n = \text{Suz}(2^{2n+1})$ , we have  $\theta(G_n), \theta'(G_n) \rightarrow 1/2$ .

**Theorem (M.)** The conjecture holds for:

- Finite groups of order  $< 2^9$ .
- Simple groups of order  $\leq 10^9$ .
- All sporadic groups.
- $A_n, L_2(q), \text{Suz}(2^{2n+1}), \text{Ree}(3^{2n+1})$ .

**Theorem (Thompson)**  $\theta(G) > 1/3$ .

**Theorem (Gallagher)**  $\theta'(G) > 1/3$ .

**Question** What are the greatest lower bounds:  $\inf_G \theta(G)$ ,  $\inf_G \theta'(G)$  ?

**Conjecture (M.)**  $\inf \theta(G), \inf \theta'(G) = 1/2$ .

**Proposition (M.)** For  $G_n = \text{Suz}(2^{2n+1})$ , we have  $\theta(G_n), \theta'(G_n) \rightarrow 1/2$ .

**Theorem (M.)** The conjecture holds for:

- Finite groups of order  $< 2^9$ .
- Simple groups of order  $\leq 10^9$ .
- All sporadic groups.
- $A_n, L_2(q), \text{Suz}(2^{2n+1}), \text{Ree}(3^{2n+1})$ .
- All finite nilpotent groups.

**Theorem (Thompson)**  $\theta(G) > 1/3$ .

**Theorem (Gallagher)**  $\theta'(G) > 1/3$ .

**Question** What are the greatest lower bounds:  $\inf_G \theta(G)$ ,  $\inf_G \theta'(G)$  ?

**Conjecture (M.)**  $\inf \theta(G), \inf \theta'(G) = 1/2$ .

**Proposition (M.)** For  $G_n = \text{Suz}(2^{2n+1})$ , we have  $\theta(G_n), \theta'(G_n) \rightarrow 1/2$ .

**Theorem (M.)** The conjecture holds for:

- Finite groups of order  $< 2^9$ .
- Simple groups of order  $\leq 10^9$ .
- All sporadic groups.
- $A_n, L_2(q), \text{Suz}(2^{2n+1}), \text{Ree}(3^{2n+1})$ .
- All finite nilpotent groups.

**Theorem (M.)** Each nonlinear irreducible character of a finite nilpotent group is zero on more than half the group.

**Theorem (Thompson)**  $\theta(G) > 1/3$ .

**Theorem (Gallagher)**  $\theta'(G) > 1/3$ .

**Question** What are the greatest lower bounds:  $\inf_G \theta(G)$ ,  $\inf_G \theta'(G)$  ?

**Conjecture (M.)**  $\inf \theta(G), \inf \theta'(G) = 1/2$ .

**Proposition (M.)** For  $G_n = \text{Suz}(2^{2n+1})$ , we have  $\theta(G_n), \theta'(G_n) \rightarrow 1/2$ .

**Theorem (M.)** The conjecture holds for:

- Finite groups of order  $< 2^9$ .
- Simple groups of order  $\leq 10^9$ .
- All sporadic groups.
- $A_n, L_2(q), \text{Suz}(2^{2n+1}), \text{Ree}(3^{2n+1})$ .
- All finite nilpotent groups.

**Theorem (M.)** Each nonlinear irreducible character of a finite nilpotent group is zero on more than half the group.

**Theorem (M.)** More than half the irreducible characters of a finite nilpotent group are zero on any given larger-than-average class.

**Theorem (Thompson)**  $\theta(G) > 1/3$ .

**Theorem (Gallagher)**  $\theta'(G) > 1/3$ .

**Question** What are the greatest lower bounds:  $\inf_G \theta(G)$ ,  $\inf_G \theta'(G)$  ?

**Conjecture (M.)**  $\inf \theta(G), \inf \theta'(G) = 1/2$ .

**Proposition (M.)** For  $G_n = \text{Suz}(2^{2n+1})$ , we have  $\theta(G_n), \theta'(G_n) \rightarrow 1/2$ .

**Theorem (M.)** The conjecture holds for:

- Finite groups of order  $< 2^9$ .
- Simple groups of order  $\leq 10^9$ .
- All sporadic groups.
- $A_n, L_2(q), \text{Suz}(2^{2n+1}), \text{Ree}(3^{2n+1})$ .
- All finite nilpotent groups.

**Theorem (M.)** Each nonlinear irreducible character of a finite nilpotent group is zero on more than half the group.

**Theorem (M.)** More than half the irreducible characters of a finite nilpotent group are zero on any given larger-than-average class.

**Theorem (M.)** Let  $G$  be finite nilpotent, let  $\chi \in \text{Irr}(G)$ , and let  $g \in G$ . Then

$$\chi(g) = 0 \quad \text{or} \quad \tilde{\text{Tr}} \left( |\chi(g)|^2 \right) \geq 2^{\#\{\text{primes dividing } \chi(1)\}}$$



A closing question

## A closing question

Let  $P(G)$  be either  $\text{Prob}(\chi(g) = 0)$  or  $\text{Prob}(\chi(K) = 0)$ .

## A closing question

Let  $P(G)$  be either  $\text{Prob}(\chi(g) = 0)$  or  $\text{Prob}(\chi(K) = 0)$ .

**Conjecture (M.)** For  $G_{p,n} \in \text{Syl}_p(S_n)$ , we have  $P(G_{p,n}) \rightarrow 1$  as  $n \rightarrow \infty$ .