

# The Hessian of tall-skinny networks is easy to invert

Ali Rahimi

November 21, 2025

## Abstract

We show how to solve linear systems that involve the Hessian of a deep net. Compared to the naive approach of first computing the Hessian, then solving the linear system, which takes storage that's quadratic in the number of parameters and cubically many operations, the method scales roughly like Pearlmutter's algorithm for computing Hessian-vector products. That means Hessian-inverse-vector products can be computed in time and storage that scale linearly in the number of parameters.

## 1 Introduction

The Hessian of a deep net is the matrix of second-order mixed partial derivatives of its loss with respect to its parameters. Decades ago, when deep nets had only hundreds or thousands of parameters, the Hessian matrix could be inverted to implement optimizers that converged much faster than gradient descent [9, 1]. However, much larger modern deep nets, using the Hessian has become increasingly less practical: The Hessian of a model with a billion parameters would have a quintillion entries, which is far larger than can be stored, multiplied, factorized, or inverted, even in the largest data centers. A common workaround is to approximate the Hessian as a low-rank matrix [10, 6] or as a diagonal matrix [2, 3, 5]. Such approximations makes it easier to apply the inverse of the Hessian to a vector.

It has long been known that the product of the Hessian with a fixed vector (the so-called Hessian-vector product) can be computed in time and storage that scales linearly with the number of parameters in the network. This is dramatically less than the cubic scaling of the naive algorithm that first computes the Hessian matrix and then multiplies it by the vector. The trick, which is due to Pearlmutter [8], is to modify the original network to cause the gradient of the modified network to become the desired Hessian-vector product. Given a way to compute the Hessian-vector product, one can then compute the Hessian-inverse-vector product by, say Krylov iterations like Conjugate Gradient. However, the quality of the result would then depend on the conditioning of the Hessian, which is notoriously poor for deep nets [4]. Unfortunately, there seems to exist no variant of Pearlmutter's trick to compute the Hessian-inverse-vector products directly.

This article offers a way to compute Hessian-inverse-vector products using roughly as much time as Pearlmutter's algorithm requires to compute Hessian-vector products. The Hessian of a deep net has a regular structure that makes it possible to compute Hessian-inverse-vector products exactly and efficiently. Regardless of the specific operations in each layer, the Hessian of a layerwise deep net is a low-order matrix polynomial that involves the first order and second order mixed derivatives of each layer, and the inverse of a block-bi-diagonal operator that represents the backpropagation algorithm. Exploiting this structure offers a direct way to compute Hessian-vector products using exactly the same operations as Pearlmutter's algorithm (see Appendix A). It also leads to a way to compute Hessian-inverse-vector product that does not require forming or storing the full Hessian, and without incurring quadratically (let alone cubically) many FLOPs. For an  $L$ -layer deep net where each layer has  $p$  parameters and produces  $a$  activations, naively storing the Hessian would require  $O(L^2 p^2)$  memory, and multiplying it by a vector or solving a linear system would require  $O(L^2 p^2 + L \max(a, p)^3)$  and  $O(L^3 p^3)$  operations, respectively. In contrast, we will show how to perform these operations using only  $O(L \max(a, p)^3)$  computations. The dependence on the parameters on the number of activations and parameters in each layer is still cubic, but the dependence on the number of layers is only linear. This makes operating on the Hessian of tall and skinny networks more efficient than the Hessian of short and fat networks.

## 2 Overview

Our objective is to efficiently solve linear systems of the form  $Hx = g$ , where  $H$  is the Hessian of a deep neural network, without forming  $H$  explicitly. To do this, we employ the following strategy:

1. Write down the gradient of the deep net in matrix form, as a bi-diagonal system of linear equations. Solving this system uses back-substitution, which in this case coincides exactly with the computations carried out by backpropagation.
2. Differentiate this matrix form to obtain the second-order derivatives. Write the result in matrix form, which involves a second-order polynomial in the inverse of the aforementioned bi-diagonal matrix.
3. Lift this polynomial into a higher-dimensional matrix form, and use the Woodbury formula to invert the resulting matrix.
4. Derive a fast algorithm to apply the inverse of such polynomials.

## 3 Notation

We write a deep net as a pipeline of functions  $\ell = 1, \dots, L$ ,

$$\begin{aligned} z_1 &= f_1(z_0; x_1) \\ &\dots \\ z_\ell &= f_\ell(z_{\ell-1}; x_\ell) \\ &\dots \\ z_L &= f_L(z_{L-1}; x_L) \end{aligned} \tag{1}$$

The vectors  $x_1, \dots, x_L$  are the parameters of the pipeline. The vectors  $z_1, \dots, z_L$  are its intermediate activations, and  $z_0$  is the input to the pipeline. The last layer  $f_L$  computes the final activations and their training loss, so the scalar  $z_L$  is the loss of the model on the input  $z_0$ . To make this loss's dependence on  $z_0$  and the parameters explicit, we sometimes write it as  $z_L(z_0; x)$ . This formalization deviates slightly from the traditional deep net formalism in two ways: First, the training labels are subsumed in  $z_0$ , and are propagated through the layers until they're used in the loss. Second, the last layer fuses the loss (which has no parameters) and the last layer (which does).

We'll assume the first and partial derivatives of each layer with respect to its parameters and its inputs exist. This poses some complications with ReLU activations and other non-differentiable operations in modern networks. Notably, for the Hessian to be symmetric, it must be differentiable everywhere. We'll largely ignore this complication and assume that difapproximations to these operations are used.

At the end of each section, we'll count the number of floating point operations required to compute various expressions. While the derivations do not impose any restrictions on the shape of the layers, for the purpose of this accounting, we'll assume all but the last  $L$  layers have  $a$ -dimensional activations ( $z_\ell \in \mathbb{R}^a$ ) and  $p$ -dimensional parameters ( $x_\ell \in \mathbb{R}^p$ ).

## 4 Backpropagation, the matrix way

We would like to fit the vector of parameters  $x = (x_1, \dots, x_L)$  given a training dataset, which we represent by a stochastic input  $z_0$  to the pipeline. Training the model proceeds by gradient descent steps along the stochastic gradient  $\partial z_L(z_0; x)/\partial x$ . The components of this direction can be computed by the chain rule with

a backward recursion:

$$\frac{\partial z_L}{\partial x_\ell} = \underbrace{\frac{\partial z_L}{\partial z_\ell}}_{b_\ell} \underbrace{\frac{\partial z_\ell}{\partial x_\ell}}_{\nabla_x f_\ell} \quad (2)$$

$$\frac{\partial z_L}{\partial z_\ell} = \underbrace{\frac{\partial z_L}{\partial z_{\ell+1}}}_{b_{\ell+1}} \underbrace{\frac{\partial z_{\ell+1}}{\partial z_\ell}}_{\nabla_z f_{\ell+1}}. \quad (3)$$

The identification  $b_\ell \equiv \frac{\partial z_L}{\partial z_\ell}$ ,  $\nabla_x f_\ell \equiv \frac{\partial z_\ell}{\partial x_\ell}$ , and  $\nabla_z f_\ell \equiv \frac{\partial z_\ell}{\partial z_{\ell-1}}$  turns this recurrence into

$$\frac{\partial z_L}{\partial x_\ell} = b_\ell \cdot \nabla_x f_\ell \quad (4)$$

$$b_\ell = b_{\ell+1} \cdot \nabla_z f_{\ell+1}, \quad (5)$$

with the base case  $b_L = 1$ , a scalar. These two equations can be written in vector form as

$$\frac{\partial z_L}{\partial x} = \begin{bmatrix} \frac{\partial z_L}{\partial x_1} & \cdots & \frac{\partial z_L}{\partial x_L} \end{bmatrix} = \underbrace{\begin{bmatrix} b_1 & \cdots & b_L \end{bmatrix}}_{\equiv b} \underbrace{\begin{bmatrix} \nabla_x f_1 & & \\ & \ddots & \\ & & \nabla_x f_L \end{bmatrix}}_{\equiv D_x}, \quad (6)$$

and

$$\underbrace{\begin{bmatrix} b_1 & b_2 & b_3 & \cdots & b_{L-1} & b_L \end{bmatrix} \begin{bmatrix} I & & & & \\ -\nabla_z f_2 & I & & & \\ & -\nabla_z f_3 & I & & \\ & & \ddots & \ddots & \\ & & & -\nabla_z f_L & 1 \end{bmatrix}}_{\equiv M} = \underbrace{\begin{bmatrix} 0 & \cdots & 1 \end{bmatrix}}_{\equiv e_L}. \quad (7)$$

Solving for  $b$  and substituting back gives

$$\frac{\partial z_L}{\partial x} = e_L M^{-1} D_x. \quad (8)$$

The matrix  $M$  is block bi-diagonal. Its diagonal entries are identity matrices, and its off-diagonal matrices are the gradient of the intermediate activations with respect to the layer's parameters. The matrix  $D_x$  is block diagonal, with the block as the derivative of each layer's activations with respect to its inputs.  $M$  is invertible because the spectrum of a triangular matrix can be read off its diagonal, which in this case is all ones.

The number of operations required to compute

## 5 The Hessian

The gradient we computed above is the unique vector  $v$  such that  $dz_L \equiv z_L(x+dx) - z_L(dx) \rightarrow v(x) \cdot dx$  as  $dx \rightarrow 0$ . In this section, we compute the Hessian  $H$  of  $z_L$  with respect to the parameters. This is the unique matrix  $H(x)$  such that  $dv^\top \equiv v^\top(x+dx) - v^\top(x) \rightarrow H(x) dx$  as  $dx \rightarrow 0$ . We use the facts that  $dM^{-1} = -M^{-1}(dM)M^{-1}$  and  $b = e_L M^{-1}$  to write

$$\begin{aligned} dv &= d(e_L M^{-1} D_x) \\ &= e_L M^{-1} (dD_x) + e_L (dM^{-1}) D_x \\ &= b \cdot dD_x - e_L M^{-1} (dM) M^{-1} D_x \\ &= b \cdot dD_x - b \cdot (dM) M^{-1} D_x \end{aligned} \quad (9)$$

We compute each of these terms separately. As part of this agenda, we will rely on the gradient of tensor-valued functions  $g : \mathbb{R}^d \rightarrow \mathbb{R}^{o_1 \times \dots \times o_k}$ . Define this gradient  $\nabla_x g(x) \in \mathbb{R}^{(o_1 \dots o_k) \times d}$  as the unique matrix-valued function that satisfies

$$\text{vec}(g(x + dx) - g(x)) \rightarrow \nabla_x g(x) \cdot dx \quad (10)$$

as  $dx \rightarrow 0$ . This convention readily implies the Hessian of a vector-valued function: If  $g : \mathbb{R}^d \rightarrow \mathbb{R}^o$ , then  $\nabla_{xx} g(x) \in \mathbb{R}^{o \times d^2}$  is the unique matrix such that  $\text{vec}(\nabla_x g(x + dx) - \nabla_x g(x)) \rightarrow \nabla_{xx} g(x) dx$ . This convention also readily accommodates the chain rule. For example, the gradient of  $h(x) \equiv f(g(x))$  for matrix-valued  $f$  and  $g$  can be written as  $\nabla f \nabla g$  as expected. It also implies partial derivatives like  $\nabla_{yz} g$  for  $g : \mathbb{R}^{|x|} \rightarrow \mathbb{R}^{|g|}$ . If  $y \in \mathbb{R}^{|y|}$  and  $z \in \mathbb{R}^{|z|}$  are restrictions of  $x \in \mathbb{R}^{|x|}$  to some  $|y|$  and  $|z|$ -dimensional subsets, then  $\nabla_z g(x) \in \mathbb{R}^{|g| \times |z|}$ , and  $\nabla_{yz} g(x) = \nabla_y \nabla_z g(x) \in \mathbb{R}^{|g| \times |z| \times |y|}$ . See Chapter 6 of Magnus & Neudecker [7] for a deeper treatment of higher order derivatives of vector-valued functions.

## 5.1 The term involving $dD_x$

The matrix  $D_x$  is block-diagonal with its  $\ell$ th diagonal block containing the matrix  $D_\ell \equiv \nabla_x f_\ell$ . Using the facts that  $\text{vec}(ABC) = (C^\top \otimes A) \text{vec}(B)$ , and  $(A \otimes B)^\top = A^\top \otimes B^\top$ , we get

$$\begin{aligned} b \cdot (dD_x) &= [b_1 \quad \dots \quad b_L] \begin{bmatrix} dD_1 & & \\ & \ddots & \\ & & dD_L \end{bmatrix} \\ &= [b_1 \cdot dD_1 \quad \dots \quad b_L \cdot dD_L] \\ &= [\text{vec}(dD_1)^\top (I \otimes b_1^\top) \quad \dots \quad \text{vec}(dD_L)^\top (I \otimes b_L^\top)] \\ &= \begin{bmatrix} \text{vec}(dD_1) \\ \vdots \\ \text{vec}(dD_L) \end{bmatrix}^\top \begin{bmatrix} I \otimes b_1^\top & & \\ & \ddots & \\ & & I \otimes b_L^\top \end{bmatrix} \end{aligned} \quad (11)$$

Observe that  $\text{vec}(dD_\ell) = d \text{vec} \nabla_x f_\ell(z_{\ell-1}; x_\ell)$  varies with  $dx$  through both its arguments  $x_\ell$  and  $z_{\ell-1}$ . Using mixed partials of vector-valued functions described above, we get

$$\text{vec}(dD_\ell) = d \text{vec}(\nabla_x f_\ell) = (\nabla_{xx} f_\ell) dx_\ell + (\nabla_{zx} f_\ell) dz_{\ell-1}. \quad (12)$$

Stacking these equations gives

$$\begin{bmatrix} \text{vec}(dD_1) \\ \vdots \\ \text{vec}(dD_L) \end{bmatrix} = \begin{bmatrix} \nabla_{xx} f_1 & & \\ & \ddots & \\ & & \nabla_{xx} f_L \end{bmatrix} dx + \begin{bmatrix} \nabla_{zx} f_1 & & \\ & \ddots & \\ & & \nabla_{zx} f_L \end{bmatrix} \begin{bmatrix} dz_0 \\ \vdots \\ dz_{L-1} \end{bmatrix}. \quad (13)$$

Each vector  $dz_\ell$  in turn varies with  $dx$  via  $dz_\ell = (\nabla_x f_\ell) dx_\ell + (\nabla_z f_\ell) dz_{\ell-1}$ , with the base case  $dz_0 = 0$ , since the input  $z_0$  does not vary with  $dx$ . Stacking up this recurrence gives

$$\begin{bmatrix} I & & \\ -\nabla_z f_2 & I & \\ & \ddots & \\ -\nabla_z f_L & & 1 \end{bmatrix} \begin{bmatrix} dz_1 \\ \vdots \\ dz_{L-1} \\ dz_L \end{bmatrix} = \begin{bmatrix} \nabla_x f_1 & & \\ & \ddots & \\ & & \nabla_x f_L \end{bmatrix} dx. \quad (14)$$

We can solve for the vector  $\begin{bmatrix} dz_1 \\ \vdots \\ dz_L \end{bmatrix} = M^{-1} D_x dx$  and use the downshifting matrix

$$P \equiv \begin{bmatrix} 0 & & \\ I & 0 & \\ & \ddots & \\ & & I & 0 \end{bmatrix} \quad (15)$$

to plug back the vector  $\begin{bmatrix} dz_0 \\ \vdots \\ dz_{L-1} \end{bmatrix} = PM^{-1}D_x dx$ :

$$\begin{bmatrix} \text{vec}(dD_1) \\ \vdots \\ \text{vec}(dD_L) \end{bmatrix} = \left( \begin{bmatrix} \nabla_{xx} f_1 & & \\ & \ddots & \\ & & \nabla_{xx} f_L \end{bmatrix} + \begin{bmatrix} \nabla_{zx} f_1 & & \\ & \ddots & \\ & & \nabla_{zx} f_L \end{bmatrix} PM^{-1}D_x \right) dx. \quad (16)$$

## 5.2 The term involving $dM$

The matrix  $dM$  is lower-block-diagonal with  $dM_2, \dots, dM_L$ , and  $dM_\ell \equiv d\nabla_z f_\ell$ . Similar to the above, we can write

$$b \cdot (dM)M^{-1}D_x = [b_1 \quad \dots \quad b_{L-1} \quad b_L] \begin{bmatrix} 0 & & \\ -dM_2 & 0 & \\ & \ddots & \\ & & -dM_L \quad 0 \end{bmatrix} M^{-1}D_x \quad (17)$$

$$= -[b_2 \cdot dM_2 \quad \dots \quad b_L \cdot dM_L \quad 0] M^{-1}D_x \quad (18)$$

$$= -[\text{vec}(dM_2)^\top (I \otimes b_2^\top) \quad \dots \quad \text{vec}(dM_L)^\top (I \otimes b_L^\top) \quad 0] M^{-1}D_x \quad (19)$$

$$= -\begin{bmatrix} \text{vec}(dM_1) \\ \vdots \\ \text{vec}(dM_L) \end{bmatrix}^\top \begin{bmatrix} 0 & & \\ I \otimes b_2^\top & 0 & \\ & \ddots & \\ & & I \otimes b_L^\top & 0 \end{bmatrix} M^{-1}D_x \quad (20)$$

$$= -\begin{bmatrix} \text{vec}(dM_1) \\ \vdots \\ \text{vec}(dM_L) \end{bmatrix}^\top \begin{bmatrix} I \otimes b_1^\top & & \\ & \ddots & \\ & & I \otimes b_L^\top \end{bmatrix} PM^{-1}D_x. \quad (21)$$

Each matrix  $dM_\ell = d\nabla_z f_\ell(z_{\ell-1}; x_\ell)$  varies with  $dx$  through both  $x_\ell$  and  $z_{\ell-1}$  as  $d \text{vec}(M_\ell) = (\nabla_{xz} f_\ell) dx_\ell + (\nabla_{zz} f_\ell) dz_{\ell-1}$ . Following the steps of the previous section gives

$$\begin{bmatrix} \text{vec}(dM_1) \\ \vdots \\ \text{vec}(dM_L) \end{bmatrix} = \left( \begin{bmatrix} \nabla_{xz} f_1 & & \\ & \ddots & \\ & & \nabla_{xz} f_L \end{bmatrix} + \begin{bmatrix} \nabla_{zz} f_1 & & \\ & \ddots & \\ & & \nabla_{zz} f_L \end{bmatrix} PM^{-1}D_x \right) dx. \quad (22)$$

## 5.3 Putting it all together

We have just shown that the Hessian of the deep net has the form

$$H \equiv \frac{\partial^2 z_L}{\partial x^2} = D_D (D_{xx} + D_{zx} PM^{-1}D_x) + D_x^\top M^{-T} P^\top D_M (D_{xz} + D_{zz} PM^{-1}D_x) \quad (23)$$

$$= D_D D_{xx} + D_D D_{zx} PM^{-1}D_x + D_x^\top M^{-T} P^\top D_M D_{xz} + D_x^\top M^{-T} P^\top D_M D_{zz} PM^{-1}D_x. \quad (24)$$

The various matrices involved are recapitulated below:

$$\begin{aligned}
D_D &\equiv \begin{bmatrix} \underbrace{I \otimes b_1}_{p \times ap} & & \\ & \ddots & \\ & & I \otimes b_L \end{bmatrix}, & D_M &\equiv \begin{bmatrix} \underbrace{I \otimes b_1}_{a \times a^2} & & \\ & \ddots & \\ & & I \otimes b_L \end{bmatrix}, \\
P &\equiv \begin{bmatrix} 0 & & & \\ I & 0 & & \\ & \ddots & & \\ & & I & 0 \end{bmatrix}, & M &\equiv \begin{bmatrix} I & & & & \\ \underbrace{-\nabla_z f_2}_{a \times a} & I & & & \\ & -\nabla_z f_3 & I & & \\ & & \ddots & \ddots & \\ & & & -\nabla_z f_L & 1 \end{bmatrix}, \\
D_x &\equiv \begin{bmatrix} \underbrace{\nabla_x f_1}_{a \times p} & & \\ & \ddots & \\ & & \nabla_x f_L \end{bmatrix}, & D_{xx} &\equiv \begin{bmatrix} \underbrace{\nabla_{xx} f_1}_{ap \times p} & & \\ & \ddots & \\ & & \nabla_{xx} f_L \end{bmatrix}, & D_{xz} &\equiv \begin{bmatrix} \underbrace{\nabla_{xz} f_1}_{a^2 \times p} & & \\ & \ddots & \\ & & \nabla_{xz} f_L \end{bmatrix}, \\
D_{zx} &\equiv \begin{bmatrix} \underbrace{\nabla_{zx} f_1}_{ap \times a} & & \\ & \ddots & \\ & & \nabla_{zx} f_L \end{bmatrix}, & D_{zz} &\equiv \begin{bmatrix} \underbrace{\nabla_{zz} f_1}_{a^2 \times a} & & \\ & \ddots & \\ & & \nabla_{zz} f_L \end{bmatrix}.
\end{aligned}$$

## 5.4 Multiplying a vector by the Hessian

Given a vector  $g \in \mathbb{R}^{Lp}$ , the formula above allows us to compute  $Hg$  in  $O(Lap^2 + La^2p + La^3)$  operations without forming  $H$ . This cost is dominated by multiplying by the  $D_{xx}$ ,  $D_{zx}$ , and  $D_{zz}$  matrices. Appendix A shows that these operations are exactly the operations involved in Pearlmutter's trick to compute the Hessian-vector product.

It's tempting to use this insight to use Krylov methods to solve systems of the form  $Hx = b$  without forming  $H$ . This would require compute  $Hg$  some number of times that depends on the condition number of  $H$ . However, the next section shows how to compute  $H^{-1}b$  with roughly only as many operations as are needed to compute  $Hg$ .

## 6 The inverse of the Hessian

The above shows that the Hessian is a second order matrix polynomial in  $M^{-1}$ . While  $M$  itself is block-bidiagonal,  $M^{-1}$  is dense, so  $H$  is dense. Nevertheless, this polynomial can be lifted into a higher order object whose inverse is easy to compute:

$$H = D_D D_{xx} + D_D D_{zx} P M^{-1} D_x + D_x^\top M^{-\top} P^\top D_M D_{xz} + D_x^\top M^{-\top} P^\top D_M D_{zz} P M^{-1} D_x$$

We wish to solve the system  $Hx = g$  for  $x$ . We can convert this dense system involving inverses into a larger, sparse system by introducing auxiliary variables. Define

$$y \equiv M^{-1} D_x x, \tag{25}$$

which implies  $My - D_x x = 0$ . Define a second auxiliary variable

$$z \equiv M^{-\top} (P^\top D_M D_{xz} x + P^\top D_M D_{zz} P y), \tag{26}$$

which implies  $M^\top z - P^\top D_M D_{xz} x - P^\top D_M D_{zz} P y = 0$ . The original equation  $Hx = g$  becomes

$$D_D D_{xx} x + D_D D_{zx} P y + D_x^\top z = g. \tag{27}$$

Collecting these three linear equations gives us a unified system

$$\underbrace{\begin{bmatrix} D_D D_{xx} & D_D D_{zx} P & D_x^\top \\ -D_x & M & 0 \\ -P^\top D_M D_{xz} & -P^\top D_M D_{zz} P & M^\top \end{bmatrix}}_{\mathcal{K}} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} g \\ 0 \\ 0 \end{bmatrix}. \quad (28)$$

Finding  $x = H^{-1}g$  is equivalent to solving this following block-linear system for  $x, y, z$ . The matrix  $\mathcal{K}$  has dimension roughly  $(Lp + 2La) \times (Lp + 2La)$ .  $D_D D_{xx}$ ,  $D_D D_{zx} P$ ,  $P^\top D_M D_{xz}$ ,  $P^\top D_M D_{zz} P$ , and  $D_x$  are block-diagonal or block-off-diagonal, and  $M$  is block-bidiagonal.

If we reorder the variables layer-by-layer as  $(x_1, y_1, z_1, \dots, x_L, y_L, z_L)$ , the matrix  $\mathcal{K}$  becomes a block-banded matrix with bandwidth proportional to  $(p + a)$ . Solving a linear system with a banded matrix of size  $N$  and bandwidth  $w$  requires  $O(Nw^2)$  operations. Here,  $N \propto L$  and  $w \propto (p + a)$ . Therefore, we can compute  $H^{-1}g$  in  $O(L(p + a)^3)$  time. This confirms that for tall-skinny networks, inverting the Hessian is linear in the depth of the network.

## References

- [1] Etienne Barnard. Optimization for training neural nets. *IEEE Transactions on Neural Networks*, 3(2):232–240, March 1992.
- [2] Suzanna Becker and Yann LeCun. Improving the convergence of back-propagation learning with second order methods. *Proceedings of the 1988 Connectionist Models Summer School*, pages 29–37, 1989.
- [3] John Duchi, Elad Hazan, and Yoram Singer. Adaptive subgradient methods for online learning and stochastic optimization. *Journal of Machine Learning Research*, 12(61):2121–2159, 2011.
- [4] Behrooz Ghorbani, Shankar Krishnan, and Ying Xiao. An investigation into neural net optimization via hessian eigenvalue density. *CoRR*, abs/1901.10159, 2019.
- [5] Diederik P. Kingma and Jimmy Ba. Adam: A method for stochastic optimization. *arXiv preprint arXiv:1412.6980*, 2014.
- [6] Quoc V. Le, Jiquan Ngiam, Adam Coates, Ahbik Lahiri, Bobby Prochnow, and Andrew Y. Ng. On optimization methods for deep learning. In *Proceedings of the 28th International Conference on Machine Learning (ICML)*, pages 265–272, Bellevue, Washington, USA, 2011. Omnipress.
- [7] Jan R Magnus and Heinz Neudecker. *Matrix Differential Calculus with Applications in Statistics and Econometrics*. John Wiley & Sons, 3rd edition, 2019.
- [8] Barak A. Pearlmutter. Fast exact multiplication by the hessian. *Neural Computation*, 6(1):147–160, 1994.
- [9] Richard L. Watrous. Learning algorithms for connectionist networks: Applied gradient methods of nonlinear optimization. In *Proceedings of the First IEEE International Conference on Neural Networks*, volume 2, pages 619–627, San Diego, CA, June 1987. IEEE.
- [10] Andrew R. Webb and David Lowe. A hybrid optimisation strategy for adaptive feed-forward layered networks. Technical Report RSRE Memorandum 4193, Royal Signals and Radar Establishment (RSRE), Malvern, UK, 1988.

## A Equation (24) is Pearlmutter’s Hessian-vector multiplication algorithm

We showed that Equation (24) makes it possible to compute Hessian-vector product  $Hv$  in time linear in  $L$ . These operations are equivalent to Pearlmutter’s [8] algorithm, a framework to compute Hessian-vector

products in networks with arbitrary topologies. This section specializes Pearlmutter’s machinery to the pipeline topology, and shows that the operations it produces coincide exactly with those of Equation (24).

Consider a set of vectors  $v_1, \dots, v_L$  that match the dimensions of the parameter vectors  $x_1, \dots, x_L$ . Just as  $z_L(x_1, \dots, x_L)$  denotes the loss under the parameters  $w$ , we’ll consider the perturbed loss  $z_L(x_1 + \alpha v_1, \dots, x_L + \alpha v_L)$  with a scalar  $\alpha$ . By the chain rule,

$$\left. \frac{\partial}{\partial \alpha} z_L(x_1 + \alpha_1 v_1, \dots, x_L + \alpha_L v_L) \right|_{\alpha=0} = \nabla_x z_L(x_1, \dots, x_L) \cdot v. \quad (29)$$

Applying  $\nabla_x$  to both sides gives

$$\nabla_x \left. \frac{\partial}{\partial \alpha} z_L(x_1 + \alpha_1 v_1, \dots, x_L + \alpha_L v_L) \right|_{\alpha=0} = \nabla_x^2 z_L(x_1, \dots, x_L) \cdot v. \quad (30)$$

In other words, to compute the Hessian-vector product  $\nabla_x^2 z_L \cdot v$ , it suffices to compute the gradient of  $\frac{\partial z_L}{\partial \alpha}$  with respect to  $x$ . We can do this by applying standard backpropagation to  $\frac{\partial z_L}{\partial \alpha}$ . At each stage  $\ell$  during its backward pass, backpropagation produces  $\frac{\partial}{\partial x_\ell} \frac{\partial z_L}{\partial \alpha} = \nabla_{x_\ell, x} z_L \cdot v$ , yielding a block of rows in  $\nabla_x^2 z_L \cdot v$ .

To see that this generates the same operations as applying Equation (24) to  $v$ , we’ll write the backprop operations from Equation (2) against  $\frac{\partial z_L}{\partial \alpha}$  explicitly. We’ll use again the fact that  $z_\ell$  depends on  $\alpha$  through both  $z_{\ell-1}$  and  $x_\ell + \alpha v_\ell$  to massage the backward recursion of for  $\frac{\partial z_L}{\partial \alpha}$  into a format that matches Equation (2):

$$b'_\ell \equiv \frac{\partial}{\partial z_\ell} \frac{\partial z_L}{\partial \alpha} = \frac{\partial}{\partial \alpha} \frac{\partial z_L}{\partial z_\ell} = \frac{\partial}{\partial \alpha} b_\ell = \frac{\partial}{\partial \alpha} [b_{\ell+1} \cdot \nabla_z f_{\ell+1}] \quad (31)$$

$$= b'_{\ell+1} \cdot \nabla_z f_{\ell+1} + \left[ (I \otimes b_{\ell+1}) \frac{\partial}{\partial \alpha} \text{vec}(\nabla_z f_{\ell+1}) \right]^\top \quad (32)$$

$$= b'_{\ell+1} \cdot \nabla_z f_{\ell+1} + \left[ (I \otimes b_{\ell+1}) \left( \nabla_{zz} f_{\ell+1} \cdot \frac{\partial z_\ell}{\partial \alpha} + \nabla_{xz} f_{\ell+1} \cdot v_{\ell+1} \right) \right]^\top. \quad (33)$$

During the backward pass, from  $b_\ell$  and  $b'_\ell$ , we compute

$$\frac{\partial}{\partial x_\ell} \frac{\partial z_L}{\partial \alpha} = (\nabla_{x_\ell, x} z_L) \cdot v = \frac{\partial}{\partial \alpha} \frac{\partial z_L}{\partial x_\ell} = \frac{\partial}{\partial \alpha} \left[ \frac{\partial z_L}{\partial z_\ell} \frac{\partial z_\ell}{\partial x_\ell} \right] = \frac{\partial}{\partial \alpha} [b_\ell \cdot \nabla_x f_\ell] \quad (34)$$

$$= b'_\ell \cdot \nabla_x f_\ell + \left[ (I \otimes b_\ell) \frac{\partial}{\partial \alpha} \text{vec}(\nabla_x f_\ell) \right]^\top \quad (35)$$

$$= b'_\ell \cdot \nabla_x f_\ell + \left[ (I \otimes b_\ell) \left( \nabla_{zx} f_\ell \cdot \frac{\partial z_{\ell-1}}{\partial \alpha} + \nabla_{xx} f_\ell \cdot v_\ell \right) \right]^\top. \quad (36)$$

Stacking these backward equations horizontally with  $b' \equiv [b'_1 \dots b'_L]$ ,  $g_\ell^\alpha \equiv \frac{\partial z_\ell}{\partial \alpha}$  and  $g^\alpha \equiv \begin{bmatrix} g_1^\alpha \\ \vdots \\ g_L^\alpha \end{bmatrix}$ , then transposing, gives

$$\begin{aligned} M^\top (b')^\top &= P D_M (D_{zz} g^\alpha + D_{xz} v) \\ \nabla_x^2 z_L \cdot v &= D_x^\top (b')^\top + D_D (D_{zx} P g^\alpha + D_{xx} v). \end{aligned} \quad (37)$$

$g_\ell^\alpha$  can be computed during the forward pass via

$$g_\ell^\alpha \equiv \frac{\partial z_\ell}{\partial \alpha} = \nabla_z f_\ell \cdot \frac{\partial z_{\ell-1}}{\partial \alpha} + \nabla_x f_\ell \cdot v_\ell = \nabla_z f_\ell \cdot g_{\ell-1}^\alpha + \nabla_x f_\ell \cdot v_\ell, \quad (38)$$

which when stacked up, gives  $M g^\alpha = D_x v$ . Plugging  $g^\alpha$  back into Equation (37) and solving for  $b'$  gives

$$\nabla_x^2 z_L \cdot v = D_x^\top M^{-\top} P D_M (D_{zz} M^{-1} D_x v + D_{xz} v) + D_D (D_{zx} P M^{-1} D_x v + D_{xx} v). \quad (39)$$

This coincides with Equation (24), showing that the two algorithms are equivalent.