

ADVERTISEMENT

simethicone (OTC)

Brand and Other Names: Mylicon, Phazyme, Gas X Extra Strength, Gas X Softgels, Gas X Thin Strips, Children's Gas X Tongue Twisters Thin Strips, Gas X, PediaCare Infant's Gas Relief Drops, St. Joseph Infant's Gas Relief

Classes: Gastrointestinal Agents, Other



Dosing & Uses

ADULT

Dosage Forms & Strengths

capsule

- 125mg
- 180mg
- 250mg
- 500mg

tablet, chewable

- 80mg
- 125mg
- 250mg

liquid, oral

- 20mg/0.3mL

oral suspension

- 20mg/0.3mL
- 40mg/0.6mL

strip, orally disintegrating

- 62.5mg

Gas Retention in Gastrointestinal Tract

40-360 mg PO q6hr after meals and at bedtime PRN; not to exceed 500 mg/day

PEDIATRIC**Dosage Forms & Strengths****capsule**

- 125mg
- 180mg
- 250mg
- 500mg

tablet, chewable

- 80mg
- 125mg
- 250mg

liquid, oral

- 20mg/0.3mL

oral suspension

- 20mg/0.3mL
- 40mg/0.6mL

strip, orally disintegrating

- 62.5mg

Gas Retention in Gastrointestinal Tract

<2 years: 20 mg (0.3 mL) PO q6hr after meals and at bedtime; not to exceed 240 mg/day

2-12 years: 40 mg (0.6 mL) PO q6hr after meals and at bedtime; not to exceed 480 mg/day

>12 years: 40-360 mg PO q6hr after meals and at bedtime; not to exceed 500 mg/day

Adverse Effects

Frequency Not Defined

Loose stools

Warnings

Contraindications

Hypersensitivity to simethicone

Cautions

Chewable tablet should be chewed thoroughly before swallowing; liquid-filled capsules should not be chewed

Can cause false-negative gastric guaiac test

Pregnancy & Lactation

Pregnancy category: C

Lactation: Not absorbed systemically; not expected to be excreted in milk

Pregnancy Categories

A: Generally acceptable. Controlled studies in pregnant women show no evidence of fetal risk.

B: May be acceptable. Either animal studies show no risk but human studies not available or animal studies showed minor risks and human studies done and showed no risk.

C: Use with caution if benefits outweigh risks. Animal studies show risk and human studies not available or neither animal nor human studies done.

D: Use in LIFE-THREATENING emergencies when no safer drug available. Positive evidence of human fetal risk.

X: Do not use in pregnancy. Risks involved outweigh potential benefits. Safer alternatives exist.

NA: Information not available.

Pharmacology

Mechanism of Action

Changes surface tension of gas bubbles and causes collapse of foam bubbles, thus allowing easier passage of gas and preventing gas pockets in gastrointestinal tract

Elimination

Excretion: Feces (as unchanged drug); complete excretion within 24-48 hr

Medscape prescription drug monographs are based on FDA-approved labeling information, unless otherwise noted, combined with additional data derived from primary medical literature.
