Grade sheet for Final Exam Problem Two Part 1: Pearson's Correlation Coefficient

Part 1: Ignoring exposure status, is there a relationship between the Day 1 and Day 15 body weight measures? What is the strength and direction of the relationship? Is it possible to illustrate the relationship graphically? If yes, show how

Topic:

Previous studies have shown a link between pesticide exposure and thyroid disease, which can lead to increased weight gain, a symptom of thyroid disease.

Population:

Rats exposed to a certain pesticide

Research Question:

Whether or not there is a strong association between body weights measurements of day 1 and day 15.

Methods

Description of Outcome: The weights of the rats day on 1 and day 15

Description of Data Summary for Each Variable:


110 rats were a part of a research study to test the effects of a certain pesticide on weight gain. The rats were randomly selected, and divide among two groups: rats that either receive low exposure of this pesticide or high exposure of this pesticide. The trial lasted for 15 days and the weights of the rats were recorded on day 1 and day 15.

Verification of normality:

Per the central limit theorem, if the sample size is above 20 the sample mean will follow a normal distribution.

Verification of Linear Association:

We will verify that the variables have a linear association by looking at the scatter plot of the weights on day 1 and day 15. Scatter plot below:



Statement of Null Hypothesis:

There is no association between day 1 of trial and day 15 of trial of body weight measurements (H_o: p=0)

Statement of Alternative Hypothesis:

There is an association between day 1 of trial and day 15 of trial of body weight measurements (H₀: p≠0)

Statistical Method for testing Association:

Pearson's correlation coefficient

Decision Rule:

Reject H_0 in favor of H_A if p-value is less than alpha (p-value< α) otherwise fail to reject the null H_0

Method of Computation:

R statistical software version 2.11.1

Significance Level:

 $\alpha = 0.05$

Results

Data Summary for Day 1 weights and Day 15 Weights included in table 1.0

Table 1.0 Summary table of Weights of Rats on Day 1 and Day 15

Group	n	mean	SD	95% CI
Day 1 Weights	110	97.54	10.22	95.6088, 99.4721
Day 15 Weights	110	127.25	10.51	125.2620, 129.2353

Normality is assumed per the CLT based on adequate sample size for each group

Assumptions:

Sample is representative

Sample is large enough

Sample measurements for each subject are independent of one another

Measurements of each group are independent of each other

Linear relationship

Heteroskedasticity

Test Results:

r: 0.76 | p-value: <0.0001

P-value Results

Since p-value < 0.05, we reject the H_0 in favor of H_A

Description of Association:

The correlation coefficient for weights is 0.76, thus the relationship of body weight measurements on day 1 and day 15 of the sample is significant and positively associated.

Discussion

Evidence suggests, that on day 1 and day 15 of the trial there is a significant and positive association between body weight measurements of the rats.

Implication of Results

Moving forward, researchers should close high quality of results.	ely monitor the we	ight of the rats bef	ore, during, and aft	ter testing to ensure a

R-code

T4 Problem2<- read.csv(file.choose(),header=TRUE)

Et<-T4 Problem2[T4 Problem2\$Exposure=='low' & T4 Problem2\$BWDay1,1]

Ft<-T4 Problem2[T4 Problem2\$Exposure=='low' & T4 Problem2\$BWDay15,2]

Gt<-T4 Problem2[T4 Problem2\$Exposure=='high' & T4 Problem2\$BWDay1,1]

Ht<-T4 Problem2[T4 Problem2\$Exposure=='high' & T4 Problem2\$BWDay15,2]

Day1 lo<-c(Et) #49

Day15 lo<-c(Ft) #49

Day1 hi<-c(Gt) #61

Day15 hi<-c(Ht)#61

just day 1<-T4 Problem2[T4 Problem2\$BWDay1,1]

mean_just_day_1<-mean(just_day_1)

qqPlot((Day1 lo))

qqPlot(Day15 lo)

#low

MDay1 lo<-mean(Day1 lo) #98.01

MDay15 lo<-mean(Day15 lo) #126.57

#hi

MDay15 hi<-mean(Day15 hi) #127.78

MDay1 hi<-mean(Day1 hi) #97.15

#question one correlation. #Ignoring amount of radiation, is there a relationship between

#weight of day one, and day 15.

#what is strength and direction?

#per scatter plot, there is a positive association in weights on day 1 vs day 2

#null hypothesis, there is no relationship(H0: p=0). alternative hypothesis, there is a relationship(H0: p/=0) #assumptions:

#sample is representative.subjects are indepedntly measured, relationship is linear, sample size is above 20 #variability of one measure to be more/less contast throughout the values of the other measure and vice/versa (heteroskedasticity)

plot(T4 Problem2\$BWDay1, T4 Problem2\$BWDay15,

xlab="Day 1 Weights", ylab="Day 15 Weights",

main = "Wegith of Rats: Day 1 vs Day 15")

#summerize the two measurements with mean sizes, means, SD 95CI

ddply(T4 Problem2, .(Exposure), summarize, BWDay1=sd(BWDay1))

#Exposure BWDay1

high 10.777890

low 9.573986

ddply(T4 Problem2, .(Exposure), summarize, BWDay15=sd(BWDay15))

#Exposure BWDay15

high 10.71949

low 10.32063

CI(Day1 lo, ci=0.95)

#upper mean lower

#100.76221 98.01224 95.26228

CI(Day15_lo, ci=0.95)
#upper mean lower
#129.5434 126.5790 123.6145
CI(Day15_hi, ci=0.95)
#upper mean lower
#130.5319 127.7866 125.0412
CI(Day1_hi, ci=0.95)
#upper mean lower
#99.92182 97.16148 94.40113

CI(T4 Problem2\$BWDay1, ci=0.95)

#upper mean lower #99.47213 97.54045 95.60878

CI(T4_Problem2\$BWDay15, ci=0.95) #upper mean lower #129.2353 127.2486 125.2620

sd(T4_Problem2\$BWDay15)
sd(T4_Problem2\$BWDay1)

cor.test(T4_Problem2\$BWDay1, T4_Problem2\$BWDay15,

alternative = "two.sided", method = "pearson", conf.level = 0.95)

#Pearson's product-moment correlation

#data: T4_Problem2\$BWDay1 and T4_Problem2\$BWDay15

#t = 12.25, df = 108, p-value < 2.2e-16

#alternative hypothesis: true correlation is not equal to 0

#95 percent confidence interval:

#0.6711525 0.8311335

#sample estimates:

#cor

#0.7625644

#r value is positive number close to 1, therefore there is a strong relationship