

# Exploring Periodic Boundary Conditions: A Study on 2D and 3D Spheres and Continuum Limit

Alireza Same<sup>1</sup>, Hanie Hatami<sup>2</sup>, Mehrnaz Rabiee<sup>3</sup>

---

## Abstract

We analyze systems with periodic boundary conditions in two and three dimensions. Beginning with an  $N$ -particle system confined to a circle, we derive the governing ordinary differential equation and solve it both analytically using circulant matrices and numerically. We then take the continuum limit as  $N$  goes to infinity to obtain a wave equation on the circle with periodic boundary conditions. This partial differential equation is solved exactly, allowing us to construct visualizations of the propagating waves. Next we study an  $N$ -particle system confined to the surface of a regular polyhedron, deriving the discrete equations and solving numerically using symmetry arguments. We then discuss statistical mechanics approaches for this system. Finally, we analyze longitudinal elastic waves propagating on a sphere, deriving the continuum wave equation in spherical coordinates and discussing analytical and numerical solution methods based on spectral decomposition. Throughout we emphasize the mathematical physics, solving equations analytically where possible and numerically otherwise. Results for all systems are illustrated through figures.

**Keywords:** Particle Systems, Mathematical Physics, Fourier Analysis, Differential Equations, Continuum Limit, Periodic Symmetry, Platonic Solids

---

## 1. Introduction

Periodic boundary conditions arise naturally in the study of waves and particles constrained to closed topologies. The mathematics of such systems, though conceptually simple, reveals rich physical behaviors. In one dimension, systems with periodic boundaries can be solved exactly. As the dimension increases, new challenges emerge. On a circle, the topology is uniform and easily parametrized, yielding to straightforward analysis via Fourier decomposition. On a sphere, the additional dimension introduces variability in the surface curvature, requiring more advanced methods like spherical harmonics. In three dimensions, modeling particles or waves constrained to regular convex polyhedron brings deeper obstacles, as these Platonic solids resist exact geometric parameterization.

Approximation methods must therefore complement exact mathematical techniques in exploring these systems. The interplay of analytical, numerical, and geometric ideas is what makes periodic systems fascinating to study. When symmetry allows clean solutions, the mathematics elegantly explains the physics. And when intricacies of higher dimensions obstruct analysis, numerical methods and geometric intuition guide the way.

In this Article, we embark on a mathematical tour of periodic boundaries in multiple dimensions. We begin with approachable problems, deriving and solving ordinary differential

equations through matrix techniques. Moving to continuous systems, we encounter tractable partial differential equations which yield to Fourier analysis. Finally, we contend with intricate geometries, wrestling with equations which only numerical computation can conquer. What ties these investigations together is the quest to understand systems constrained to live within themselves, endowed with a symmetry as beautiful as it is challenging.

## 2. 2D Spheres Revisited

### 2.1. Analytical Description

As established in foundational texts on wave phenomena (see [Georgi](#)), a prototypical model for cyclic wave propagation considers a discrete set of point particles confined to a circular domain. This framework relies on two key approximations: first, that interactions are strictly local, with each particle only directly influencing its immediate neighbors. Second, that the inter-particle forces respond linearly to changes in separation relative to an equilibrium length. While undeniably reductionist, these assumptions prove sufficient to capture the salient physics of transverse wave transmission through the medium.

Specifically, we examine a system of  $N$  identical particles of mass  $m$  regularly spaced around a ring of radius  $R$ . Identical harmonic springs, characterized by force constant  $k$ , link adjacent particles along the circumference. This geometry exhibits a continuous rotational symmetry - rotations and reflections by  $2\pi/N$  map the configuration onto itself. Our theoretical description should therefore not differentiate between the  $2N$  symmetry-equivalent orientations. The sole measurable degree of freedom lies in the particles' relative displacements, as

---

Email addresses: alirezasame@gmail.com (Alireza Same), alirezasame@gmail.com (Hanie Hatami), alirezasame@gmail.com (Mehrnaz Rabiee)

<sup>1</sup>99100709

<sup>2</sup>99100614

<sup>3</sup>99100669

absolute positions along the circle can not be experimentally determined without first arbitrarily fixing a reference point.

The simplicity of this model, alongside its inherent periodicity, motivate its study before tackling more intricate geometries in higher dimensions. Even within the confines of localized, linear interactions, analytical and computational techniques reveal rich wave behaviors arising from the cyclic topology. The mathematics provides profound and elegant insight into the underlying physics.

As an elementary example, Figure 1 illustrates the case for  $N = 4$  particles. The fourfold rotational symmetry is evident, as is the periodic nature of the circular boundary. With the model established, we now embark on an in-depth analysis.




Figure 1: 4 identically distributed point masses on a circle

The configuration of the system can be fully specified by the central angle displacement  $\theta_i$  of each particle  $i$  relative to its initial angular position. As per the rotational symmetry, the labeling of individual particles carries no physical significance. We therefore characterize the state through the  $N$ -dimensional vector  $\theta = (\theta_1, \theta_2, \dots, \theta_N)$  encompassing each particle's angular shift. For small oscillations, we expect the time evolution of  $\theta$  to follow a linear equation of the form:

$$M\ddot{\theta} + K\theta = 0 \Rightarrow \ddot{\theta} + W\theta = 0 \quad (1)$$

The matrix  $W$  belongs to the special class of circulant matrices which respect circular permutation symmetry. To derive its structure, consider the  $i$ th particle's interactions: since it only couples to its nearest neighbors, its motion is governed by:

$$\ddot{\theta}_i = a\theta_{i-1} + b\theta_i + c\theta_{i+1}$$

By symmetry, the coefficient  $a$  equals  $c$ . For a system of  $N$  particles, this yields an interaction matrix  $W$  of the form:

$$W = \begin{pmatrix} A & B & 0 & \dots & B \\ B & A & B & \dots & 0 \\ 0 & B & A & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ B & 0 & 0 & \dots & A \end{pmatrix}_{N \times N} \quad (2)$$

While the precise values of  $A$  and  $B$  do not qualitatively impact the dynamics, specific models may impose further constraints such as  $A = -2B$  (see Georgi). Analytical solutions rely on the circulant structure, as discussed in Gray.

As discussed, the normal modes of this system should respect the circular symmetry. This mandates mode shapes of the form:

$$\beta^{(k)} = (1, \omega^k, \omega^{2k}, \dots, \omega^{(N-1)k}) \quad \omega = e^{2\pi i/N}$$

where  $\omega$  denotes the complex  $N$ th root of unity. The angular frequency  $\omega_k$  of the  $k$ th mode takes the form:

$$\omega_k^2 = A + 2B \cos\left(\frac{2k\pi}{N}\right) \quad k = 0, 1, \dots, N-1$$

This reveals intrinsic degeneracies arising from the system's symmetry. For even  $N$ , the  $k = 0$  and  $k = N/2$  modes have real-valued eigenvectors, while the remaining  $N - 2$  modes come in complex-conjugate pairs which can be combined into degenerate real modes. The connection to Fourier analysis will become transparent in examining the continuum limit. The normal modes indeed correspond to a Discrete Fourier Transform, as elaborated in Gray. The circulant structure enables analytical decomposition into these symmetry-dictated eigenmodes.

In the Case where  $A = -2B$  the relation simplifies to the simpler case below:

$$\omega_k^2 = \omega_0^2 \sin^2\left(\frac{k\pi}{N}\right)$$

## 2.2. Numerical Description and Comparison with the Analytical Solution

Thus far, we have relied on linearized equations to enable an analytical treatment. To complement this, we now numerically simulate the full nonlinear particle interactions without simplifications. Details of the model and algorithms are in the supplementary simulation [Jupyter Notebook](#). In short, we integrate the coupled second-order ODEs for the  $N$  particles using an adaptive timestep fourth-order Runge-Kutta Python routine. By systematically increasing the imposed oscillation amplitude, we can validate the accuracy of our small amplitude analytical solutions and probe changes at higher displacements. These computations provide a flexible framework to incorporate future refinements like stochastic forces or dissipation. Most importantly, the nonlinear simulations both test the assumptions made earlier and better represent the true dynamics.

## 3. Discussion

## 4. Summary and conclusions

### Acknowledgements

Thanks to ...

### References

- Georgi, H., . The Physics of Waves.
- Gray, R.M., . Toeplitz and Circulant Matrices: A Review 2, 155–239. URL: <https://www.nowpublishers.com/article/Details/CIT-006>, doi:10.1561/0100000006.