

Application of learning algorithms to nonlinear filtering and MCMC

PhD Dissertation defense

Nov 6, 2019

Anand Radhakrishnan



Department of Electrical and Computer Engineering — University of Florida

Committee chair: Prof. Sean Meyn

Committee members : Prof. José Principe, Prof. Kamran Mohseni, Prof. James Hobert

Thanks to the National Science Foundation and Army Research Office

Thanks to friends & family

Outline

- 1 PhD Proposal - Recap
- 2 Poisson's Equation
- 3 Differential LSTD learning
- 4 Applications to Nonlinear filtering
- 5 Applications to MCMC
- 6 Conclusions and Future Work

great that you
went through
this fast

PhD Proposal - Recap

Work till then

- Presented the **feedback particle filter (FPF)** - an approximation of the nonlinear filter that requires computing a gain function by solving a *Poisson's equation*.
version of
- Proposed a **differential TD-learning** algorithm to approximate the gradient of the solution to Poisson's equation. [R et al. 16]

PhD Proposal - Recap

Work till then

- Presented the **feedback particle filter (FPF)** - an approximation of the nonlinear filter that requires computing a gain function by solving a *Poisson's equation*.
- Proposed a **differential TD-learning** algorithm to approximate the gradient of the solution to Poisson's equation. [R et al. 16]
- Presented numerical examples for gain approximation and state estimation for scalar systems.

PhD Proposal - Recap

Tasks promised and accomplished

- Extend the algorithm to a multidimensional setting.

PhD Proposal - Recap

Tasks promised and accomplished

- Extend the algorithm to a multidimensional setting. ~~✓~~ [✓] [R et al. 18]
- Explore appropriate basis selection for these algorithms.

PhD Proposal - Recap

Tasks promised and accomplished

- Extend the algorithm to a multidimensional setting. ✓ [R et al. 18] *delete*
- Explore appropriate basis selection for these algorithms. ✓ *delete*
- Impact of gain approximation error on filtering performance.

PhD Proposal - Recap

Tasks promised and accomplished

- Extend the algorithm to a multidimensional setting. - ✓ [R et al. 18]
- Explore appropriate basis selection for these algorithms. - ✓
- Impact of gain approximation error on filtering performance. - ✗
[Taghvaei et al. 18]

PhD Proposal - Recap

Tasks promised and accomplished

- Extend the algorithm to a multidimensional setting. - ✓ [R et al. 18]
- Explore appropriate basis selection for these algorithms. - ✓
- Impact of gain approximation error on filtering performance. - X
[Taghvaei et al. 18]

New direction *kernel methods*

- Simplified differential LSTD algorithm for Langevin diffusion. [R et al. 18]
- Improvements for online gain estimation. [R et al. 19]
- Explored the application to MCMC algorithms. [Brosse et al. 19]

Poisson's Equation

$$0 = \tilde{c} + \mathcal{D}h$$

$$h(x) = \mathbb{E} \left[\int_0^{\tau} \tilde{c}(X(t)) dt \right]$$

with $X(0) = x$

Optimal FPF Gain

$$\mathbf{K} = \nabla h$$

$$= 2\|\Delta h_{\theta}\|_2$$

$$\mathcal{L}_2^{\text{CLT}} = 2\langle h_{\theta}, e_{\theta} \rangle$$

$$\mathcal{L}_{\text{CLT}}^2 = \langle 2\Delta h, e \rangle$$

$$\phi(x) = \max_u \left\{ \mathbb{E} \left[\tilde{c}(x+u) \right] - \frac{1}{2} \|u\|^2 \right\}$$

Optimal MCMC CV

Optimal Control

Poisson's Equation

Poisson's Equation

- Second order partial differential equation with applications in various fields.
- General form in stochastic systems

~~delete~~ (no "equal by definition")

$$\mathcal{D}h = -f$$

\mathcal{D} - differential operator

f - forcing function, usually centered

Poisson's Equation

- Second order partial differential equation with applications in various fields.
- General form in stochastic systems

$$\mathcal{D}h := -f$$

\mathcal{D} - differential operator

f - forcing function, usually centered

h - solution to Poisson's equation

Poisson's Equation

Stochastic optimal control

~~del w.c?~~
or introduce P_*

Example: In average cost optimal control problems,

$$\min_u \{c(x, u) + P_u h^*(x)\} = h^*(x) + \eta^*$$

$c(x, u)$ - Cost function associated with state x and action u .

P_u - Transition kernel of the controlled Markov chain.

η^* - Optimal average cost.

$h^*(\cancel{x})$ - Relative value function

delete

Where is D ? f ?

$$Dh^* = -\tilde{c}$$

Poisson's Equation

Stochastic optimal control

Example: In average cost optimal control problems,

$$\min_u \{c(x, u) + P_u h^*(x)\} = h^*(x) + \eta^*$$

$c(x, u)$ - Cost function associated with state x and action u .

P_u - Transition kernel of the controlled Markov chain.

η^* - Optimal average cost.

$h^*(x)$ - **Relative value function**

Poisson's equation is the dynamic programming equation.

Poisson's Equation

Langevin Diffusion

Langevin diffusion is given by the SDE,

$$d\Phi_t = \underbrace{-\nabla U(\Phi_t) dt}_{\text{Drift term}} + \underbrace{\sqrt{2} dW_t}_{\text{Diffusion term}}, \quad \Phi \in \mathbb{R}^d$$

$U \in C^1$ is called the *potential function*.

$W = \{W_t : t \geq 0\}$ is a standard Brownian motion on \mathbb{R}^d .

Poisson's Equation

Langevin Diffusion

Langevin diffusion is given by the SDE,

$$d\Phi_t = \underbrace{-\nabla U(\Phi_t) dt}_{\text{Drift term}} + \underbrace{\sqrt{2} dW_t}_{\text{Diffusion term}}, \quad \Phi \in \mathbb{R}^d$$

$U \in C^1$ is called the *potential function*.

$\mathbf{W} = \{W_t : t \geq 0\}$ is a standard Brownian motion on \mathbb{R}^d .

- May be regarded as a d -dimensional gradient flow with “noise”.

Poisson's Equation

Langevin Diffusion

Langevin diffusion is given by the SDE,

$$d\Phi_t = \underbrace{-\nabla U(\Phi_t) dt}_{\text{Drift term}} + \underbrace{\sqrt{2} dW_t}_{\text{Diffusion term}}, \quad \Phi \in \mathbb{R}^d$$

$U \in C^1$ is called the *potential function*.

$\mathbf{W} = \{W_t : t \geq 0\}$ is a standard Brownian motion on \mathbb{R}^d .

- May be regarded as a d -dimensional gradient flow with “noise”.
- Diffusion is reversible, with unique *invariant density* $\rho = e^{-U+\Lambda}$, where Λ is a normalizing constant.

Poisson's Equation

Langevin Diffusion

Differential generator \mathcal{D}_t :

$$\mathcal{D}f := \lim_{t \rightarrow 0} \frac{\mathbb{E}[f(\Phi_t) | \Phi_0 = x] - f(x)}{t}$$

Poisson's Equation

Langevin Diffusion

Differential generator \mathcal{D} ,

$$\begin{aligned}\mathcal{D}f &:= \lim_{t \rightarrow 0} \frac{\mathbb{E}[f(\Phi_t) | \Phi_0 = x] - f(x)}{t} \\ &= -\nabla U \cdot \nabla f + \Delta f, \quad f \in C^2,\end{aligned}$$

where ∇ is the gradient and Δ is the Laplacian.

Poisson's Equation

Langevin Diffusion

Differential generator \mathcal{D} ,

$$\begin{aligned}\mathcal{D}f &:= \lim_{t \rightarrow 0} \frac{\mathbb{E}[f(\Phi_t) | \Phi_0 = x] - f(x)}{t} \\ &= -\nabla U \cdot \nabla f + \Delta f, \quad f \in C^2,\end{aligned}$$

where ∇ is the gradient and Δ is the Laplacian.

Let $c: \mathbb{R}^d \rightarrow \mathbb{R}$ be a function of interest, and

$$\eta = \int c(x) \rho(x) dx = \langle c, 1 \rangle_{L^2}.$$

Poisson's Equation

Langevin Diffusion

Differential generator \mathcal{D} ,

$$\begin{aligned}\mathcal{D}f &:= \lim_{t \rightarrow 0} \frac{\mathbb{E}[f(\Phi_t) | \Phi_0 = x] - f(x)}{t} \\ &= -\nabla U \cdot \nabla f + \Delta f, \quad f \in C^2,\end{aligned}$$

where ∇ is the gradient and Δ is the Laplacian.

Let $c: \mathbb{R}^d \rightarrow \mathbb{R}$ be a function of interest, and

$$\eta = \int c(x) \rho(x) dx = \langle c, 1 \rangle_{L^2}.$$

Function $h \in C^2$ solves Poisson's equation with forcing function c if

*need this
earlier
with
sum dis.
for time*

$$\left. \begin{aligned} \mathcal{D}h &:= -\tilde{c}, & \tilde{c} &= c - \eta. \\ h &:= \int_0^\infty \mathbb{E}[\tilde{c}(\Phi_t)] dt \end{aligned} \right\}$$

Poisson's Equation

Existence of a solution

- A solution exists under weak assumptions on U and c [Glynn 96, Kontoyiannis 12].
- Representations for the gradient of h and bounds are obtained in [Laugesen 15, Devraj 18].
- A smooth solution $h \in C^2$ exists under stronger conditions in [Pardoux 01], subject to growth conditions on c similar to [Glynn 96].

Poisson's Equation

Approximate solution to Poisson's equation

Obtaining an analytical solution for h is difficult outside special cases.
Hence approximation.

Poisson's Equation

Approximate solution to Poisson's equation

Obtaining an analytical solution for h is difficult outside special cases.
Hence approximation.

Goal: For a given function class \mathcal{H} , find the minimizer of

$$g^* := \arg \min_{g \in \mathcal{H}} \|h - g\|_{L^2}^2$$

Such minimum norm optimization problems can be solved using *TD learning* [Tsitsikilis 99, CTCN].

Poisson's Equation

Approximate solution to Poisson's equation

Obtaining an analytical solution for h is difficult outside special cases.
Hence approximation.

Goal: For a given function class \mathcal{H} , find the minimizer of

$$g^* := \arg \min_{g \in \mathcal{H}} \|h - g\|_{L^2}^2$$


Such minimum norm optimization problems can be solved using *TD learning* [Tsitsiklis 99, CTCN].

to solve  if the process

Challenge - No algorithm exists for state spaces of dimension > 1

does not regenerate (for diffusions, dim > 1 ruled out).

LSTD learning

Discounted cost case

Discounted-cost value function:

$$h^\gamma(x) := \int_0^\infty e^{-\gamma t} \mathsf{E}_x[c(\Phi_t)] dt, \quad \gamma > 0$$

discount factor

Discounted-cost optimality equation:

$$\gamma h^\gamma = c + \mathcal{D}h^\gamma$$

I prefer color,
but o.k.

LSTD learning

Discounted cost case

Discounted-cost value function:

$$h^\gamma(x) := \int_0^\infty e^{-\gamma t} \mathsf{E}_x[c(\Phi_t)] dt, \quad \gamma > 0 - \text{discount factor}$$

Discounted-cost optimality equation:

$$\gamma h^\gamma = c + \mathcal{D}h^\gamma$$

LSTD goal $\hat{g}^* := \arg \min_{g \in \mathcal{H}} \|h^\gamma - g\|_{L^2}^2$

LSTD learning

Discounted cost case

$$\text{LSTD goal} - g^* := \arg \min_{g \in \mathcal{H}} \|h^\gamma - g\|_{L^2}^2$$

For a linear parameterization

$$g = h^\theta := \sum_{i=1}^{\ell} \theta_i \psi_i$$
$$\theta^* = M^{-1} b$$

$$M_{ij} = \langle \psi_i, \psi_j \rangle_{L^2}, \quad b_i = \langle \psi_i, h^\gamma \rangle_{L^2}$$

LSTD learning

Discounted cost case

$$\text{LSTD goal} - g^* := \arg \min_{g \in \mathcal{H}} \|h^\gamma - g\|_{L^2}^2$$

For a linear parameterization

$$g = h^\theta := \sum_{i=1}^{\ell} \theta_i \psi_i$$
$$\theta^* = M^{-1} b$$

$$M_{ij} = \langle \psi_i, \psi_j \rangle_{L^2}, \quad b_i = \langle \psi_i, h^\gamma \rangle_{L^2}$$

LSTD learning

Discounted cost case

$$\text{LSTD goal} - g^* := \arg \min_{g \in \mathcal{H}} \|h^\gamma - g\|_{L^2}^2$$

For a linear parameterization

$$g = h^\theta := \sum_{i=1}^{\ell} \theta_i \psi_i$$

$$\theta^* = M^{-1}b$$

$$M_{ij} = \langle \psi_i, \psi_j \rangle_{L^2}, \quad b_i = \langle \psi_i, h^\gamma \rangle_{L^2}$$

$$= \langle \psi_i, R_\gamma c \rangle_{L^2}$$

Resolvent kernel \bullet

$$R_\gamma c(x) := \int_0^\infty \mathbb{E}_x \left[e^{-\gamma t} c(\Phi_t) \right] dt$$

$$R_\gamma c = (I_\gamma - \mathcal{D})^{-1} c$$

why subscript?
 $(I_\gamma - \mathcal{D})$

LSTD learning

Discounted cost case

$$\text{LSTD goal} - g^* := \arg \min_{g \in \mathcal{H}} \|h^\gamma - g\|_{L^2}^2$$

For a linear parameterization

$$g = h^\theta := \sum_{i=1}^{\ell} \theta_i \psi_i$$

$$\theta^* = M^{-1} b$$

$$M_{ij} = \langle \psi_i, \psi_j \rangle_{L^2}, \quad b_i = \langle \psi_i, R_\gamma c \rangle_{L^2}$$

$$= \langle R_\gamma^\dagger \psi_i, c \rangle_{L^2}$$

Using an adjoint operation and applying the stationarity of Φ .

LSTD learning

Discounted cost case

$$\text{LSTD goal} - g^* := \arg \min_{g \in \mathcal{H}} \|h^\gamma - g\|_{L^2}^2$$

For a linear parameterization

$$g = h^\theta := \sum_{i=1}^{\ell} \theta_i \psi_i$$

$$\theta^* = M^{-1} b$$

$$M_{ij} = \langle \psi_i, \psi_j \rangle_{L^2}, \quad b_i = \langle R_\gamma^\dagger \psi_i, c \rangle_{L^2}$$

Eligibility vector $\varnothing \varphi_6(t) := \int_0^\infty e^{-\gamma r} \psi(\Phi_{t-r}) dr$

$\varnothing(+)$

$$R_\gamma^\dagger \psi_i(x) = E[\varnothing_i(t) | \Phi_t = x]$$

\varnothing_i

LSTD learning

Discounted cost case

$$\text{LSTD goal } \underset{g \in \mathcal{H}}{\arg \min} \|h^\gamma - g\|_{L^2}^2$$

For a linear parameterization

$$g = h^\theta := \sum_{i=1}^{\ell} \theta_i \psi_i$$

$$\theta^* = M^{-1}b$$

$$\begin{aligned} M_{ij} &= \langle \psi_i, \psi_j \rangle_{L^2}, \quad b_i = \langle R_\gamma^\dagger \psi_i, c \rangle_{L^2} \\ &= \mathbb{E}[\varphi_{\psi_i}(t) c(\Phi_t)] \end{aligned}$$

LSTD learning

Discounted cost case

ODE formulation of the LSTD algorithm:

$$\begin{aligned}\frac{d}{dt}M(t) &= \psi(\Phi_t)\psi^T(\Phi_t) \\ \frac{d}{dt}\varphi_\psi(t) &= -\gamma \varphi_\psi(t) + \psi(\Phi_t) \\ \frac{d}{dt}b(t) &= \varphi_\psi(t)c(\Phi_t) \\ \theta(t) &:= M(t)^{-1}b(t)\end{aligned}$$

LSTD learning

Discounted cost case

ODE formulation of the LSTD algorithm:

$$\begin{aligned}\frac{d}{dt}M(t) &= \psi(\Phi_t)\psi^T(\Phi_t) \\ \frac{d}{dt}\varphi_\psi(t) &= -\gamma \varphi_\psi(t) + \psi(\Phi_t) \\ \frac{d}{dt}b(t) &= \varphi_\psi(t)c(\Phi_t) \\ \theta(t) &:= M(t)^{-1}b(t)\end{aligned}$$

By law of large numbers,

$$\lim_{t \rightarrow \infty} \theta(t) = \theta^*$$

LSTD learning

Discounted cost case

ODE formulation of the LSTD algorithm:

$$\frac{d}{dt} M(t) = \psi(\Phi_t) \psi^T(\Phi_t)$$

$$\frac{d}{dt} \varphi_\psi(t) = -\gamma \varphi_\psi(t) + \psi(\Phi_t)$$

$$\frac{d}{dt} b(t) = \varphi_\psi(t) c(\Phi_t)$$

$$\theta(t) := M(t)^{-1} b(t)$$

By law of large numbers,

be able to defend this.

$$\lim_{t \rightarrow \infty} \theta(t) = \theta^*$$

~~Drawback~~ For average-cost, LSTD requires the existence of a regenerating state.

Differential LSTD learning (∇ -LSTD)

Poisson's equation

Idea: Approximate the gradient of h directly [R et al. 16, Devraj et al. 16]:

$$g^* := \arg \min_{g \in \mathcal{H}} \|\nabla h - \nabla g\|_{L^2}^2$$

Differential LSTD learning (∇ -LSTD)

Poisson's equation

Idea: Approximate the gradient of h directly [R et al. 16, Devraj et al. 16]:

$$g^* := \arg \min_{g \in \mathcal{H}} \|\nabla h - \nabla g\|_{L^2}^2$$

Need to choose a function class \mathcal{H} for g (or ∇g)

- A finitely parameterized family of functions.
- A reproducing kernel Hilbert space (RKHS).

Differential LSTD learning (∇ -LSTD)

Poisson's equation

Idea: Approximate the gradient of h directly [R et al. 16, Devraj et al. 16]:


$$g^* := \arg \min_{g \in \mathcal{H}} \|\nabla h - \nabla g\|_{L^2}^2$$

Need to choose a function class \mathcal{H} for g (or ∇g)

- A finitely parameterized family of functions.
- A reproducing kernel Hilbert space (RKHS).

Choice of basis is not an easy task

\implies RKHS framework is far easier to implement.



Differential LSTD learning (∇ -LSTD)

Poisson's equation

$$\nabla\text{-LSTD goal} - g^* := \arg \min_{g \in \mathcal{H}} \|\nabla h - \nabla g\|_{L^2}^2$$

Challenge: the function h is not known,
and hence the objective function is not observable

Differential LSTD learning (∇ -LSTD)

Poisson's equation

$$\nabla\text{-LSTD goal} - g^* := \arg \min_{g \in \mathcal{H}} \|\nabla h - \nabla g\|_{L^2}^2$$

Challenge: the function h is not known,
and hence the objective function is not observable

∇ -LSTD: For a linear parameterization

$$g = h^\theta := \sum_{i=1}^{\ell} \theta_i \psi_i \implies \nabla g = \sum_{i=1}^{\ell} \theta_i \nabla \psi_i$$

$$\theta^* = M^{-1} b$$

$$M_{ij} = \langle \nabla \psi_i, \nabla \psi_j \rangle_{L^2}, \quad b_i = \langle \nabla \psi_i, \nabla h \rangle_{L^2}$$

Differential LSTD learning (∇ -LSTD)

Poisson's equation

$$\nabla\text{-LSTD goal } \overset{\circ}{g} := \arg \min_{g \in \mathcal{H}} \|\nabla h - \nabla g\|_{L^2}^2$$

Challenge: the function h is not known,
and hence the objective function is not observable

∇ -LSTD: For a linear parameterization

$$g = h^\theta := \sum_{i=1}^{\ell} \theta_i \psi_i \implies \nabla g = \sum_{i=1}^{\ell} \theta_i \nabla \psi_i$$

$$\theta^* = M^{-1} b$$

$$M_{ij} = \langle \nabla \psi_i, \nabla \psi_j \rangle_{L^2}, \quad b_i = \langle \nabla \psi_i, \nabla h \rangle_{L^2}$$

Differential LSTD learning (∇ -LSTD)

Poisson's equation

$$\nabla\text{-LSTD goal} - g^* := \arg \min_{g \in \mathcal{H}} \|\nabla h - \nabla g\|_{L^2}^2$$

Challenge: the function h is not known,
and hence the objective function is not observable

∇ -LSTD: For a linear parameterization

$$g = h^\theta := \sum_{i=1}^{\ell} \theta_i \psi_i \implies \nabla g = \sum_{i=1}^{\ell} \theta_i \nabla \psi_i$$

$$\theta^* = M^{-1} b$$

$$M_{ij} = \langle \nabla \psi_i, \nabla \psi_j \rangle_{L^2}, \quad b_i = \langle \nabla \psi_i, R_{U''} \nabla c \rangle_{L^2}$$

Resolvent kernel - $R_{U''} \nabla c(x) := \int_0^\infty \mathsf{E}_x \left[\exp \left(- \int_0^t U''(\Phi_s) \, ds \right) \nabla c(\Phi_t) \right] dt$

Differential LSTD learning (∇ -LSTD)

Poisson's equation

$$\nabla\text{-LSTD goal} - g^* := \arg \min_{g \in \mathcal{H}} \|\nabla h - \nabla g\|_{L^2}^2$$

Challenge: the function h is not known,
and hence the objective function is not observable

∇ -LSTD: For a linear parameterization

$$g = h^\theta := \sum_{i=1}^{\ell} \theta_i \psi_i \implies \nabla g = \sum_{i=1}^{\ell} \theta_i \nabla \psi_i$$

$$\theta^* = M^{-1} b$$

$$M_{ij} = \langle \nabla \psi_i, \nabla \psi_j \rangle_{L^2}, \quad b_i = \langle \nabla \psi_i, R_{U''} \nabla c \rangle_{L^2}$$

$$= \langle R_{U''}^\dagger \nabla \psi_i, \nabla c \rangle_{L^2}$$

Differential LSTD learning (∇ -LSTD)

Poisson's equation

$$\nabla\text{-LSTD goal } \underset{\cancel{g^*}}{\cancel{\theta}} g^* := \arg \min_{g \in \mathcal{H}} \|\nabla h - \nabla g\|_{L^2}^2$$

Challenge: the function h is not known,
and hence the objective function is not observable

∇ -LSTD-L: For Langevin diffusion, if $f, g \in L^2(\rho)$

$$\langle \nabla f, \nabla g \rangle_{L^2} = -\langle f, \mathcal{D}g \rangle_{L^2} = -\langle \mathcal{D}f, g \rangle_{L^2}.$$

Differential LSTD learning (∇ -LSTD)

Poisson's equation

$$\nabla\text{-LSTD goal} - g^* := \arg \min_{g \in \mathcal{H}} \|\nabla h - \nabla g\|_{L^2}^2$$

Challenge: the function h is not known,
and hence the objective function is not observable

∇ -LSTD-L: For Langevin diffusion, if $f, g \in L^2(\rho)$

$$\langle \nabla f, \nabla g \rangle_{L^2} = -\langle f, \mathcal{D}g \rangle_{L^2} = -\langle \mathcal{D}f, g \rangle_{L^2}.$$

Applying this and Poisson's equation $\mathcal{D}h = -\tilde{c}$:

$$\begin{aligned}\|\nabla h - \nabla g\|_{L^2}^2 &= \|\nabla h\|_{L^2}^2 + \|\nabla g\|_{L^2}^2 - 2\langle \nabla h, \nabla g \rangle_{L^2} \\ &= \|\nabla h\|_{L^2}^2 + \|\nabla g\|_{L^2}^2 + 2\langle \mathcal{D}h, g \rangle_{L^2}\end{aligned}$$

use color transformation is clear

Differential LSTD learning (∇ -LSTD)

Poisson's equation

$$\nabla\text{-LSTD goal} - g^* := \arg \min_{g \in \mathcal{H}} \|\nabla h - \nabla g\|_{L^2}^2$$

Challenge: the function h is not known,
and hence the objective function is not observable

∇ -LSTD-L: For Langevin diffusion, if $f, g \in L^2(\rho)$

$$\langle \nabla f, \nabla g \rangle_{L^2} = -\langle f, \mathcal{D}g \rangle_{L^2} = -\langle \mathcal{D}f, g \rangle_{L^2}.$$

Applying this and Poisson's equation $\mathcal{D}h = -\tilde{c}$:

$$\begin{aligned}\|\nabla h - \nabla g\|_{L^2}^2 &= \|\nabla h\|_{L^2}^2 + \|\nabla g\|_{L^2}^2 - 2\langle \nabla h, \nabla g \rangle_{L^2} \\ &= \|\nabla h\|_{L^2}^2 + \|\nabla g\|_{L^2}^2 - 2\langle \tilde{c}, g \rangle_{L^2}\end{aligned}$$

Differential LSTD learning (∇ -LSTD – L)

Poisson's equation

$$\text{ ∇ -LSTD-L goal} - g^* := \arg \min_{g \in \mathcal{H}} \left\{ \|\nabla g\|_{L^2}^2 - 2\langle \tilde{c}, g \rangle_{L^2} \right\}$$

∇ -LSTD-L: For a linear parameterization

$$g = h^\theta := \sum_{i=1}^{\ell} \theta_i \psi_i \implies \nabla g = \sum_{i=1}^{\ell} \theta_i \nabla \psi_i$$
$$\theta^* = M^{-1} b$$

Differential LSTD learning (∇ -LSTD – L)

Poisson's equation

$$\text{ ∇ -LSTD-L goal} - g^* := \arg \min_{g \in \mathcal{H}} \left\{ \|\nabla g\|_{L^2}^2 - 2\langle \tilde{c}, g \rangle_{L^2} \right\}$$

∇ -LSTD-L: For a linear parameterization

$$g = h^\theta := \sum_{i=1}^{\ell} \theta_i \psi_i \implies \nabla g = \sum_{i=1}^{\ell} \theta_i \nabla \psi_i$$

$$\theta^* = M^{-1} b$$

$$M_{ij} = \langle \nabla \psi_i, \nabla \psi_j \rangle_{L^2}$$

$$b_i = \langle \nabla \psi_i, \nabla \textcolor{red}{h} \rangle_{L^2} = \langle \psi_i, \tilde{c} \rangle_{L^2}$$

Differential LSTD learning (∇ -LSTD – L)

Poisson's equation

$$\nabla\text{-LSTD-L goal} - g^* := \arg \min_{g \in \mathcal{H}} \left\{ \|\nabla g\|_{L^2}^2 - 2\langle \tilde{c}, g \rangle_{L^2} \right\}$$

$\nabla\text{-LSTD-L}$: For a linear parameterization

$$g = h^\theta := \sum_{i=1}^{\ell} \theta_i \psi_i \implies \nabla g = \sum_{i=1}^{\ell} \theta_i \nabla \psi_i$$

$$\theta^* = M^{-1}b$$

$$M_{ij} = \langle \nabla \psi_i, \nabla \psi_j \rangle_{L^2}$$

↙

$$\approx \frac{1}{t} \int_0^t \nabla \psi(\Phi_s) \nabla \psi^T(\Phi_s) ds$$

↙
small step

$$b_i = \langle \nabla \psi_i, \nabla h \rangle_{L^2} = \langle \psi_i, \tilde{c} \rangle_{L^2}$$

$$\approx \frac{1}{t} \int_0^t \psi_i(\Phi_s) \tilde{c}(\Phi_s) ds$$

Differential LSTD learning on RKHS (∇ -LSTD-RKHS)

Basics of RKHS

Choose a kernel function $K(x, y)$ that is

- *Symmetric*: $K(x, y) = K(y, x)$ for any $x, y \in \mathbb{R}^d$
- *Positive definite*: For any finite subset $\{x^i\} \subset \mathbb{R}^d$, the matrix $\{M_{ij} := K(x^i, x^j)\}$ is positive definite.
- *Smooth*: $K \in C^2$ *raises questions*

K defines a ~~unique~~ reproducing kernel Hilbert space (RKHS) \mathcal{H} [Moore-Aronszajn theorem].

Inner product: If $g_\alpha = \sum_i \alpha_i K(x^i, \cdot)$ and $g_\beta = \sum_j \beta_j K(y^j, \cdot)$,

$$\langle g_\alpha, g_\beta \rangle_{\mathcal{H}} := \sum_{i,j} \alpha_i \beta_j K(x^i, y^j)$$

Reproducing property: $g_\alpha(x) = \langle g_\alpha, K(x, \cdot) \rangle_{\mathcal{H}}, \quad x \in \mathbb{R}^d$.

Differential LSTD learning on RKHS

Empirical risk minimization (ERM)

Recall ∇ -LSTD goal: $\boxed{\nabla\text{-LSTD-L goal}}?$

$$g^* = \arg \min_{g \in \mathcal{H}} \left\{ \|\nabla g\|_{L^2}^2 - 2\langle \tilde{c}, g \rangle_{L^2} \right\}$$

Approximation via empirical risk minimization (ERM):

$$\arg \min_{g \in \mathcal{H}} \underbrace{\frac{1}{N} \sum_{i=1}^N \left[\|\nabla g(x^i)\|^2 - 2\tilde{c}_N(x^i)g(x^i) \right]}_{\text{Empirical risk}} + \underbrace{\lambda \|g\|_{\mathcal{H}}^2}_{\text{Regularization}}$$

where $\tilde{c}_N(x) = c(x) - \frac{1}{N} \sum_{i=1}^N c(x^i)$, $x \in \mathbb{R}^d$

Differential LSTD learning on RKHS

Empirical risk minimization (ERM)

Extended Representer Theorem [Zhou 08]

If loss function $L(x, \cdot, \cdot)$ is convex on \mathbb{R}^{d+1} for each $x \in \mathbb{R}^d$, then the optimizer g^* over $g \in \mathcal{H}$ exists:

$$g^*(\cdot) = \sum_{i=1}^N \left[\beta_i^{0*} K(x^i, \cdot) + \sum_{k=1}^d \beta_i^{k*} \frac{\partial}{\partial x_k} K(x^i, \cdot) \right]$$

Differential LSTD learning on RKHS

Empirical risk minimization (ERM)

Extended Representer Theorem [Zhou 08]

If loss function $L(x, \cdot, \cdot)$ is convex on \mathbb{R}^{d+1} for each $x \in \mathbb{R}^d$, then the optimizer g^* over $g \in \mathcal{H}$ exists:

$$g^*(\cdot) = \sum_{i=1}^N \left[\beta_i^{0*} K(x^i, \cdot) + \sum_{k=1}^d \beta_i^{k*} \frac{\partial}{\partial x_k} K(x^i, \cdot) \right]$$

Our loss function is convex: $L(x, g, \nabla g) = \|\nabla g(x)\|^2 - 2\tilde{c}(x)g(x)$

Differential LSTD learning on RKHS

Optimal solution in one dimension

∇ -LSTD-RKHS ERM:

$$g^* = \arg \min_{g \in \mathcal{H}} \frac{1}{N} \sum_{i=1}^N \left\{ (g'(x^i))^2 - 2\tilde{c}_N(x^i)g(x^i) \right\} + \lambda \|g\|_{\mathcal{H}}^2$$

$$\text{Solution: } g^*(y) = \sum_{i=1}^N \left\{ \beta_i^{0*} K(x^i, y) + \beta_i^{1*} \partial_x K(x^i, y) \right\}, \quad y \in \mathbb{R}$$

Differential LSTD learning on RKHS

Optimal solution in one dimension

∇ -LSTD-RKHS ERM:

$$g^* = \arg \min_{g \in \mathcal{H}} \frac{1}{N} \sum_{i=1}^N \left\{ (g'(x^i))^2 - 2\tilde{c}_N(x^i)g(x^i) \right\} + \lambda \|g\|_{\mathcal{H}}^2$$

Solution: $g^*(y) = \sum_{i=1}^N \left\{ \beta_i^{0*} K(x^i, y) + \beta_i^{1*} \partial_x K(x^i, y) \right\}, \quad y \in \mathbb{R}$

Notation:

$$M_{00}(i, j) := K(x^i, x^j), \quad M_{10}(i, j) := \frac{\partial K}{\partial x}(x^i, x^j)$$

$$M_{01}(i, j) := \frac{\partial K}{\partial y}(x^i, x^j), \quad M_{11}(i, j) := \frac{\partial^2 K}{\partial x \partial y}(x^i, x^j)$$

$$\tilde{c}_j := \tilde{c}_N(x^j), \quad \beta^T = [\beta_1^0, \dots, \beta_N^0, \beta_1^1, \dots, \beta_N^1]$$

Differential LSTD learning on RKHS

Optimal solution in one dimension

∇ -LSTD-RKHS ERM:

$$g^* = \arg \min_{g \in \mathcal{H}} \frac{1}{N} \sum_{i=1}^N \left\{ (g'(x^i))^2 - 2\tilde{c}_N(x^i)g(x^i) \right\} + \lambda \|g\|_{\mathcal{H}}^2$$

Solution: $g^*(y) = \sum_{i=1}^N \left\{ \beta_i^{0*} K(x^i, y) + \beta_i^{1*} \partial_x K(x^i, y) \right\}, \quad y \in \mathbb{R}$

Computation: $\beta^* = M^{-1}b$ Boring

$$M = \frac{1}{N} \left[\begin{array}{c|c} M_{01} & \\ \hline M_{11} & \end{array} \right] [M_{10} \mid M_{11}] + \lambda \left[\begin{array}{c|c} M_{00} & M_{01} \\ \hline M_{10} & M_{11} \end{array} \right]$$

$$b = \frac{1}{N} \left[\begin{array}{c} M_{00} \\ \hline M_{10} \end{array} \right]$$

Differential LSTD learning on RKHS

Simplified solution

Drawback : Complexity grows linearly with d , since $\beta^* \in \mathbb{R}^{(d+1) \times N}$

Simplified solution : By considering the finite-dimensional function class -
 $\mathcal{H}_N := \text{span}\{K_{x^j} : 1 \leq j \leq N\}$

$$g^*(y) = \sum_{j=1}^N \beta_j^* K(x^j, y)$$

$$\beta^* = M^{-1}b$$

~~where, $M := N^{-1}M_{01}M_{10} + \lambda M_{00}$, $b := N^{-1}M_{00}\tilde{s}$~~

Surprising empirical observation : Simplified solution does as good as the optimal solution for $d \leq 5$.

Applications to Nonlinear filtering

Feedback Particle Filter

needs color
A figure

Goal : To obtain estimates of the state of a stochastic dynamical system based on noisy partial observations.

← figure here of a plane or ...

Kalman filter is optimal for a linear Gaussian system.

For nonlinear systems, conditional distribution fails to be Gaussian, cannot be captured by a finite set of parameters.

Particle filters are Monte-Carlo approximations of the nonlinear filter.

Applications to Nonlinear filtering

Feedback Particle Filter

Problem:

Signal: $dX_t = a(X_t)dt + dB_t, \quad X_0 \sim \rho_0^*,$

Observation: $dZ_t = c(X_t)dt + dW_t,$

- $X_t \in \mathbb{R}^d$ is the state at time t .
- $\{Z_t : t \geq 0\}$ is the observation process.
- $a(\cdot), c(\cdot)$ are C^1 functions.
- $\{B_t\}, \{W_t\}$ are mutually independent Wiener processes.

Applications to Nonlinear filtering

Feedback Particle Filter

Problem:

Signal: $dX_t = a(X_t)dt + dB_t, \quad X_0 \sim \rho_0^*,$

Observation: $dZ_t = c(X_t)dt + dW_t,$

- $X_t \in \mathbb{R}^d$ is the state at time t .
- $\{Z_t : t \geq 0\}$ is the observation process.
- $a(\cdot), c(\cdot)$ are C^1 functions.
- $\{B_t\}, \{W_t\}$ are mutually independent Wiener processes.
- $\rho_t^* := P(X_t | Z_s : s \leq t)$ is the posterior distribution.

↑ no brackets

Applications to Nonlinear filtering

Feedback Particle Filter

et. al. (mainly Roshan)

Feedback particle filter (FPF) [Yang 13] is motivated by techniques from mean-field optimal control.

Applications to Nonlinear filtering

Feedback Particle Filter

Feedback particle filter (FPF) [Yang 13] is motivated by techniques from mean-field optimal control.

N particles are propagated in the form of a controlled system.

$$dX_t^i = \underbrace{a(X_t^i)dt + dB_t^i}_{\text{Propagation}} + \underbrace{dU_t^i}_{\text{Update}}, \quad i = 1 \text{ to } N$$

- $X_t^i \in \mathbb{R}$ is the state of the i^{th} particle at time t
- U_t^i is the control input applied to i^{th} particle
- $\{B_t^i\}$ are mutually independent standard Wiener processes.

Applications to Nonlinear filtering

Feedback Particle Filter

Feedback particle filter (FPF) [Yang 13] is motivated by techniques from mean-field optimal control.

N particles are propagated in the form of a controlled system.

$$dX_t^i = \underbrace{a(X_t^i)dt + dB_t^i}_{\text{Propagation}} + \underbrace{dU_t^i}_{\text{Update}}, \quad i = 1 \text{ to } N$$

- $X_t^i \in \mathbb{R}$ is the state of the i^{th} particle at time t
- U_t^i is the control input applied to i^{th} particle
- $\{B_t^i\}$ are mutually independent standard Wiener processes.

Approximation of ρ_t^* :

$$\rho_t^* \approx \rho_t^{(N)}(A) = \frac{1}{N} \sum_{i=1}^N \mathbb{I}\{X_t^i \in A\}, \quad A \subset \mathbb{R}.$$

} block or
color or
something
to make
distinctive

Applications to Nonlinear filtering

Feedback Particle Filter

Asymptotically exact filter obtained by minimizing the KL divergence between ρ_t^* and ρ_t (see [Yang 13]):

$$dU_t^i = K_t(X_t^i) \circ \overbrace{(dZ_t - \frac{1}{2}[c(X_t^i) + \hat{c}_t]dt)}^{dI_t^i},$$

I_t^i : Innovations process.

K_t : FPF gain, similar in nature to the Kalman gain.

Applications to Nonlinear filtering

Feedback Particle Filter

Asymptotically exact filter obtained by minimizing the KL divergence between ρ_t^* and ρ_t (see [Yang 13]):

$$dU_t^i = K_t(X_t^i) \circ \overbrace{(dZ_t - \frac{1}{2}[c(X_t^i) + \hat{c}_t]dt)}^{dI_t^i} \otimes$$

I_t^i : Innovations process.

K_t : FPF gain, similar in nature to the Kalman gain.





Figure: Finite- N implementation

Applications to Nonlinear filtering

Feedback Particle Filter

KF:

$$d\mu_t = a(\mu_t)dt + \underbrace{K_{\text{kal}}(dZ_t - c(\mu_t)dt)}_{\text{update}}$$



Kalman filter


Applications to Nonlinear filtering

Feedback Particle Filter


go through this fast unnecessary review.

KF: $d\mu_t = a(\mu_t)dt + \underbrace{K_{kal}(dZ_t - c(\mu_t)dt)}_{\text{update}}$

FPF: $dX_t^i = a(X_t^i)dt + dB_t^i + \underbrace{K_t(X_t^i) \circ (dZ_t - \frac{1}{2}[c(X_t^i) + \hat{c}_t]dt)}_{\text{update}}$



Kalman filter



Feedback particle filter (FPF)

Applications to Nonlinear filtering

FPF Gain function

Representation: $K_t = \nabla h$

h solves Poisson's equation: $\mathcal{D}h = -\nabla U \cdot \nabla h + \Delta h = -\tilde{c}$.

Applications to Nonlinear filtering

FPF Gain function

Representation: $\mathbf{K}_t = \nabla h$

h solves Poisson's equation: $\mathcal{D}h = -\nabla U \cdot \nabla h + \Delta h = -\tilde{c}$.

Approximations to \mathbf{K} can be obtained by

$$\min_{g \in \mathcal{H}} \|\mathbf{K} - \hat{\mathbf{K}}\|_{L^2}^2 = \min_{g \in \mathcal{H}} \|\nabla h - \nabla g\|_{L^2}^2$$

Applications to Nonlinear filtering

FPF Gain function

Representation: $K_t = \nabla h$

h solves Poisson's equation: $\mathcal{D}h = -\nabla U \cdot \nabla h + \Delta h = -\tilde{c}$.

Approximations to K can be obtained by

$$\min_{g \in \mathcal{H}} \|K - \hat{K}\|_{L^2}^2 = \min_{g \in \mathcal{H}} \|\nabla h - \nabla g\|_{L^2}^2$$

Can be solved using ∇ -LSTD learning

Applications to Nonlinear filtering

FPF Gain function

Representation: $K_t = \nabla h$

h solves Poisson's equation: $\mathcal{D}h = -\nabla U \cdot \nabla h + \Delta h = -\tilde{c}$.

Approximations to K can be obtained by

$$\min_{g \in \mathcal{H}} \|K - \hat{K}\|_{L^2}^2 = \min_{g \in \mathcal{H}} \|\nabla h - \nabla g\|_{L^2}^2$$

FPF implementation requires online gain estimation for each t .

- ∇ -LSTD-RKHS with optimal mean
- ∇ -LSTD-RKHS with memory

Applications to Nonlinear filtering

∇ -LSTD-RKHS-OM

Constant gain approximation for K is the minimizer obtained over all deterministic vectors:

$$\hat{K}^* := \arg \min_{\hat{K} \in \mathbb{R}^d} \|K - \hat{K}\|_{L^2}^2$$

Applications to Nonlinear filtering

∇ -LSTD-RKHS-OM

Constant gain approximation for K is the minimizer obtained over all deterministic vectors:

$$\hat{K}^* := \arg \min_{\hat{K} \in \mathbb{R}^d} \|K - \hat{K}\|_{L^2}^2$$

Solution is evidently the mean, $\hat{K}^* = E[K]$.

$$\hat{K}_k^* = \langle K, e_k \rangle_{L^2}$$

Applications to Nonlinear filtering

∇ -LSTD-RKHS-OM

Constant gain approximation for K is the minimizer obtained over all deterministic vectors:

$$\hat{K}^* := \arg \min_{\hat{K} \in \mathbb{R}^d} \|K - \hat{K}\|_{L^2}^2$$

Solution is evidently the mean, $\hat{K}^* = E[K]$.

$$\begin{aligned}\hat{K}_k^* &= \langle K, e_k \rangle_{L^2} \\ &= \langle \nabla h, e_k \rangle_{L^2}\end{aligned}$$

Applications to Nonlinear filtering

∇ -LSTD-RKHS-OM

Constant gain approximation for K is the minimizer obtained over all deterministic vectors:

$$\hat{K}^* := \arg \min_{\hat{K} \in \mathbb{R}^d} \|K - \hat{K}\|_{L^2}^2$$

Solution is evidently the mean, $\hat{K}^* = E[K]$.

$$\begin{aligned}\hat{K}_k^* &= \langle K, e_k \rangle_{L^2} \\ &= \langle \nabla h, e_k \rangle_{L^2} \\ &= -\langle Dh, x_k \rangle_{L^2} = \langle \tilde{c}, x_k \rangle_{L^2}\end{aligned}$$

∇x_k

Applications to Nonlinear filtering

∇ -LSTD-RKHS-OM

Constant gain approximation for K is the minimizer obtained over all deterministic vectors:

$$\hat{K}^* := \arg \min_{\hat{K} \in \mathbb{R}^d} \|K - \hat{K}\|_{L^2}^2$$

Solution is evidently the mean, $\hat{K}^* = E[K]$.

$$\begin{aligned}\hat{K}_k^* &= \langle K, e_k \rangle_{L^2} \\ &= \langle \nabla h, e_k \rangle_{L^2} \\ &= -\langle \mathcal{D}h, x_k \rangle_{L^2} = \langle \tilde{c}, x_k \rangle_{L^2}\end{aligned}$$

Empirical approximation:

$$\hat{K}_k^* \approx \frac{1}{N} \sum_{i=1}^N [c(x^i) - \hat{c}] x_k^i$$

Applications to Nonlinear filtering

∇ -LSTD-RKHS-OM

Redefine the approximation to K as,

$$\nabla g = \hat{K}^* + \nabla \tilde{g}$$

Modified ERM with constraints is:

$$\begin{aligned}\tilde{g}^* &:= \arg \min_{\tilde{g} \in \mathcal{H}} \|\nabla h - \hat{K}^* - \nabla \tilde{g}\|_{L_2}^2 \\ \text{s.t. } &\langle \partial_{x_k} \tilde{g}, 1 \rangle_{L_2} = 0, \quad 1 \leq k \leq d\end{aligned}$$

Applications to Nonlinear filtering

∇ -LSTD-RKHS-OM

Redefine the approximation to K as,

$$\nabla g = \hat{K}^* + \nabla \tilde{g}$$

Modified ERM with constraints is:

$$\begin{aligned}\tilde{g}^* := \arg \min_{\tilde{g} \in \mathcal{H}} \quad & \| \nabla h - \hat{K}^* - \nabla \tilde{g} \|_{L_2}^2 \\ \text{s.t.} \quad & \langle \partial_{x_k} \tilde{g}, 1 \rangle_{L_2} = 0, \quad 1 \leq k \leq d\end{aligned}$$

Solution obtained by finding a saddle point for the Lagrangian

$$L(\tilde{g}, \mu) := \| \nabla h - \hat{K}^* - \nabla \tilde{g} \|_{L_2}^2 + \langle \mu, \nabla \tilde{g} \rangle_{L_2}$$

where $\mu \in \mathbb{R}^d$ are the Lagrange multipliers.

Applications to Nonlinear filtering

∇ -LSTD-RKHS-OM

Redefine the approximation to K as,

$$\nabla g = \hat{K}^* + \nabla \tilde{g}$$

Modified ERM with constraints is:

$$\begin{aligned}\tilde{g}^* := \arg \min_{\tilde{g} \in \mathcal{H}} & \| \nabla h - \hat{K}^* - \nabla \tilde{g} \|_{L_2}^2 \\ \text{s.t. } & \langle \partial_{x_k} \tilde{g}, 1 \rangle_{L_2} = 0, \quad 1 \leq k \leq d\end{aligned}$$

Using $\mathcal{H}_N := \text{span}\{K_{x^j} : 1 \leq j \leq N\}$, β and μ can be obtained by solving $N + d$ linear equations

$$K = \hat{K}^* + \nabla \tilde{g}^*$$

Applications to Nonlinear filtering

∇ -LSTD-RKHS-memory

Gain updates are done at $t = n\delta$, where δ is the inter-sampling time.

Continuity : $K_n = K_{t_n} \approx K_{t_{n-1}}$ if $\delta \approx 0$.

Applications to Nonlinear filtering

∇ -LSTD-RKHS-memory

Gain updates are done at $t = n\delta$, where δ is the inter-sampling time.

Continuity : $K_n = K_{t_n} \approx K_{t_{n-1}}$ if $\delta \approx 0$.

Adding a regularizer term to the loss function:

$$g_n^* := \arg \min_{g \in \mathcal{H}} \frac{1}{N} \sum_{j=1}^N L_n(x_n^j, g, \nabla g) + \lambda \|g\|_{\mathcal{H}}^2$$

$$L_n(x, g, \nabla g) := \|\nabla g(x)\|^2 - 2\tilde{c}_N(x)g(x) + \underbrace{\lambda_{mem} \|\nabla g(x) - \nabla g_{n-1}(x)\|^2}_{\text{continuity penalty}}$$

Applications to Nonlinear filtering

∇ -LSTD-RKHS-memory

Gain updates are done at $t = n\delta$, where δ is the inter-sampling time.

Continuity : $K_n = K_{t_n} \approx K_{t_{n-1}}$ if $\delta \approx 0$.

$$\beta_n^* = M^{-1}b$$

where,

$$M = (1 + \lambda_{mem}) \sum_{k=1}^d M_{10}^T M_{10} + \lambda N M_{00}$$

$$b = M_{00} \tilde{\varsigma} + \lambda_{mem} \sum_{k=1}^d M_{10}^T K_{n-1,k}$$

Applications to Nonlinear filtering

∇ -LSTD-RKHS-memory

Gain updates are done at $t = n\delta$, where δ is the inter-sampling time.

Continuity : $K_n = K_{t_n} \approx K_{t_{n-1}}$ if $\delta \approx 0$.

$$\beta_n^* = M^{-1}b \quad \text{ok, but no details.}$$

where, $M = (1 + \lambda_{mem}) \sum_{k=1}^d M_{10}^T M_{10} + \lambda N M_{00}$

$$b = M_{00} \tilde{s} + \lambda_{mem} \sum_{k=1}^d M_{10}^T K_{n-1,k}$$

Both these improvements can be applied independently or simultaneously.

Applications to Nonlinear filtering

Markov kernel approximation [Taghvaei 16]

No time

Approximates the transition kernel of the Langevin diffusion:

$$h = P_\epsilon h + \int_0^\epsilon P_s(c - \hat{c})ds,$$

Give the expression for kernel approx. Empirical approximation on particle locations x^i :

$$h_i = \sum_{j=1}^N T_{ij} h_j + \epsilon(c - \hat{c}), \text{ for } i = 1 \text{ to } N$$

Gain K is then obtained by approximating the gradient ∇h_i .

Application to nonlinear filtering

Numerical example - Gain approximation for a fixed t

Example: For a fixed t , ρ_t a Gaussian mixture

$$c(x) \equiv x$$

$$T = 10^4, 10^5, 10^6 \text{ with } \delta = 0.01, N = 1000$$


Application to nonlinear filtering

Numerical example - Gain approximation for a fixed t

Example: For a fixed t , ρ_t a Gaussian mixture

$$c(x) \equiv x$$

$$T = 10^4, 10^5, 10^6 \text{ with } \delta = 0.01, N = 1000$$



∇ -LSTD with $\psi_i = x^i \rho_1(x)$, $\psi_{i+1} = x^i \rho_2(x)$ with $1 \leq i \leq 5$.


Application to nonlinear filtering

Numerical example - Gain approximation for a fixed t

Example: For a fixed t , ρ_t a Gaussian mixture

$$c(x) \equiv x$$

$$N = 100, 500, 1000$$



∇ -LSTD-L with $\psi_i = x^i \rho_1(x)$, $\psi_{i+1} = x^i \rho_2(x)$ with $1 \leq i \leq 5$.

Application to nonlinear filtering


Numerical example - Gain approximation for a fixed t

Example: For a fixed t , ρ_t a Gaussian mixture

$$c(x) \equiv x$$

$$N = 100, 500, 1000$$

kill



Grid-search using $K_\varepsilon := e^{-\frac{\|x-x'\|^2}{4\varepsilon}}$, yields and $\varepsilon = 0.1$ and $\lambda = 10^{-2}$.


Application to nonlinear filtering

Numerical example - Gain approximation for a fixed t

Example: For a fixed t , ρ_t a Gaussian mixture

$$c(x) \equiv x$$

$$N = 100, 500, 1000$$



∇ -LSTD-RKHS-N with Gaussian kernel, $\varepsilon = 0.1$ and $\lambda = 10^{-2}$ (best!)


Application to nonlinear filtering

Numerical example - Gain approximation for a fixed t

Example: For a fixed t , ρ_t a Gaussian mixture

$$c(x) \equiv x$$

$$N = 100, 500, 1000$$



∇ -LSTD-RKHS-2N with Gaussian kernel, $\varepsilon = 0.1$ and $\lambda = 10^{-2}$


Application to nonlinear filtering

Numerical example - Gain approximation for a fixed t

Example: For a fixed t , ρ_t a Gaussian mixture

$$c(x) \equiv x$$

$$N = 100, 500, 1000$$



$\nabla\text{-LSTD-RKHS-OM}$ with Gaussian kernel, $\varepsilon = 0.1$ and $\lambda = 10^{-2}$


Application to nonlinear filtering

Numerical example - Gain approximation for a fixed t

Example: For a fixed t , ρ_t a Gaussian mixture

$$c(x) \equiv x$$

$$N = 100, 500, 1000$$



Markov kernel approximation with $\epsilon = 0.1$

Application to nonlinear filtering

Numerical example - Gain approximation for a fixed t


Example: For a fixed t , ρ_t a Gaussian mixture

$$c(x) \equiv x$$

$$N = 100, 500, 1000$$

Kill

Parameters $\{\beta_i\}$ from RKHS simplified, optimal and OM methods




Application to nonlinear filtering

Numerical example - Gain approximation for a fixed t

Example: For a fixed t , ρ_t a Gaussian mixture

$$c(x) \equiv x$$

$$N = 100, 500, 1000$$




Histogram of MSEs obtained from 400 independent trials ▶



Application to nonlinear filtering

Numerical example - Gain approximation for a fixed t

Example: $\rho(x) = \prod_{k=1}^d \rho_k(x_k)$
 $c(x) = C^T x$, where $C = \mathbb{I}_d$
 $d = 2, 5, 10$, $N = 1000$




∇ -LSTD-RKHS-OM

Application to nonlinear filtering

Numerical example - Gain approximation for a fixed t

Example: $\rho(x) = \prod_{k=1}^d \rho_k(x_k)$
 $c(x) = C^T x$, where $C = \mathbb{I}_d$
 $d = 2, 5, 10$, $N = 1000$



Markov kernel approximation

Application to nonlinear filtering

Numerical example - Gain for a nonlinear oscillator model

Example: $\rho(x)$ is a mixture of von Mises densities on a circle

$$d\vartheta = \omega dt + \sigma_B dB_t \mod 2\pi,$$

$$dZ_t = c(\vartheta)dt + \sigma_W dW_t, \quad c(\vartheta) = \frac{1}{2}[1 + \cos(\vartheta)]$$

Kill?


Application to nonlinear filtering

Numerical example - Gain for a nonlinear oscillator model

Example: $\rho(x)$ is a mixture of von Mises densities on a circle

$$d\vartheta = \omega dt + \sigma_B dB_t \mod 2\pi,$$

$$dZ_t = c(\vartheta)dt + \sigma_W dW_t, \quad c(\vartheta) = \frac{1}{2}[1 + \cos(\vartheta)]$$




∇ -LSTD-L with $\psi_i = \sin(ix)$, $\psi_{i+1} = \cos(ix)$ with $1 \leq i \leq \ell/2$,
 $\ell = 4, 6, 8$.

Applications to Nonlinear filtering

Numerical example - Parameter estimation

Example: Parameter estimation with bimodal prior

Observations: parameter plus additive noise with $\sigma_W = 1$.




Applications to Nonlinear filtering

Numerical example - Parameter estimation

Example: Parameter estimation with bimodal prior

Observations: parameter plus additive noise with $\sigma_W = 1$.




Applications to Nonlinear filtering

Numerical example - Parameter estimation

Example: Parameter estimation with bimodal prior

Observations: parameter plus additive noise with $\sigma_W = 1$.




Applications to Nonlinear filtering

Numerical example - Parameter estimation

Example: Parameter estimation with bimodal prior

Observations: parameter plus additive noise with $\sigma_W = 1$.




Applications to Nonlinear filtering

Numerical example - Parameter estimation

?

Example: Parameter estimation with bimodal prior

Observations: parameter plus additive noise with $\sigma_W = 1$.




Applications to Nonlinear filtering

Numerical example - Nonlinear 2d ship dynamics model

Example: Nonlinear ship dynamics model in 2d.

Observations: $c(x) = \arctan(x_1/x_2)$ with std. deviation $\approx 18^\circ$.

10 secs




Applications to Nonlinear filtering

Numerical example - Nonlinear 2d ship dynamics model

Example: Nonlinear ship dynamics model in 2d.

Observations: $c(x) = \arctan(x_1/x_2)$ with std. deviation $\approx 18^\circ$.




Applications to Nonlinear filtering

Numerical example - Nonlinear 2d ship dynamics model

Example: Nonlinear ship dynamics model in 2d.

Observations: $c(x) = \arctan(x_1/x_2)$ with std. deviation $\approx 18^\circ$.




Applications to Nonlinear filtering

Numerical example - Nonlinear 2d ship dynamics model

Example: Nonlinear ship dynamics model in 2d.

Observations: $c(x) = \arctan(x_1/x_2)$ with std. deviation $\approx 18^\circ$.



Applications to Nonlinear filtering

Numerical example - Nonlinear 2d ship dynamics model

Summary table of RMSEs from 100 trials

Type of filter	$\Sigma_1 = I_{2 \times 2}$	$\Sigma_2 = 5I_{2 \times 2}$	Lost track (Σ_2)
FPF RKHS-OM	0.9023	1.6254	4 times
FPF RKHS mem.	0.9162	1.9408	7 times
FPF const. gain	1.3060	2.3231	14 times
SIR PF	3.1481	4.2648	57 times
EKF	6.5203	18.441	93 times

Applications to MCMC

Introduction to MCMC

In many applications, we need to compute

$$\eta = \int c(x)\rho(x) \, dx$$

- $c: \mathbb{R}^\ell \rightarrow \mathbb{R}$ is a measurable function.
- ρ is a target probability density in \mathbb{R}^ℓ .

Markov-Chain Monte Carlo (MCMC) methods provide numerical algorithms to obtain estimates:

$$\eta_t = \frac{1}{t} \int_0^t c(\Phi(s)) \, ds$$

Φ is a Markov process with steady state distribution ρ .

Applications to MCMC

Asymptotic Variance

Estimates η_t obey a Central Limit Theorem,

$$\sqrt{t}(\eta_t - \eta) \xrightarrow{d} N(0, \gamma^2)$$

γ^2 2. Variance

Rate of convergence captured by **asymptotic variance**

$$\gamma^2 = \lim_{t \rightarrow \infty} E \left[\left(\frac{1}{\sqrt{t}} \int_0^t (c(\Phi(s)) - \eta) ds \right)^2 \right]$$

Alternate representation in terms of covariance

$$\gamma^2 := \int_{-\infty}^{\infty} R(s) ds, \quad R(s) = E[\tilde{c}(\Phi_0)\tilde{c}(\Phi_s)]$$

Applications to MCMC

Asymptotic Variance

Slide I

+ n

Representation in terms of h (Glynn 96):

$$\gamma^2 = 2\langle h, \tilde{c} \rangle$$

$$D \zeta = -\tilde{c}$$

Applications to MCMC

Asymptotic Variance

slide I

Representation in terms of h [Glynn 96]:

$$\begin{aligned}\gamma^2 &= 2\langle h, \tilde{c} \rangle \\ &= 2\|\nabla h\|_{L^2} \\ (\text{For Langevin diffusion})\end{aligned}$$

Applications to MCMC

Control Variates

Goal: To minimize asymptotic variance.

Applications to MCMC

Control Variates

Goal: To minimize asymptotic variance.

Idea: Modify the estimator using control variates [Henderson 01, CTCN]

$$c_g = c + \underbrace{\mathcal{D}g}_{\text{Control variate}}, \quad \text{where } g \in \mathcal{H}$$

$$\eta_t^g = \frac{1}{t} \int_0^t c_g(\Phi_s) ds$$

For asymptotically unbiased estimates, control variate needs to have zero-mean with respect to ρ .

Too many words

Applications to MCMC

Control Variates

Goal: To minimize asymptotic variance.

Idea: Modify the estimator using control variates [Henderson 01, CTCN]

$$c_g = c + \underbrace{\mathcal{D}g}_{\text{Control variate}}, \quad \text{where } g \in \mathcal{H}$$

$$\eta_t^g = \frac{1}{t} \int_0^t c_g(\Phi_s) ds$$

For asymptotically unbiased estimates, control variate needs to have zero-mean with respect to ρ .

For any $g \in C^2$, ~~P_ρ is invariant with respect to L_2~~ $\langle \mathcal{D}g, 1 \rangle_{L^2} = 0$.

1 ~~P_ρ is invariant with respect to L_2~~
 L_2

Applications to MCMC

Optimal control variates

Let $\tilde{h}_g = h - g$,

$$\begin{aligned}\mathcal{D}\tilde{h}_g &= \mathcal{D}h - \mathcal{D}g \\ &= -c_g + \eta\end{aligned}$$

Thus \tilde{h}_g is the solution to Poisson's equation with forcing function c_g .

Applications to MCMC

Optimal control variates

Let $\tilde{h}_g = h - g$,

$$\begin{aligned}\mathcal{D}\tilde{h}_g &= \mathcal{D}h - \mathcal{D}g \\ &= -c_g + \eta\end{aligned}$$

Thus \tilde{h}_g is the solution to Poisson's equation with forcing function c_g .

Asymptotic variance of the new estimator:

$$\gamma_g^2 = 2\langle \tilde{h}_g, \tilde{c}_g \rangle_{L^2}$$

Applications to MCMC

Optimal control variates

Slide III

Let $\tilde{h}_g = h - g$,

$$\begin{aligned}\mathcal{D}\tilde{h}_g &= \mathcal{D}h - \mathcal{D}g \\ &= -c_g + \eta\end{aligned}$$

Thus \tilde{h}_g is the solution to Poisson's equation with forcing function c_g .

Asymptotic variance of the new estimator:

$$\gamma_g^2 = 2\langle \tilde{h}_g, \tilde{c}_g \rangle_{L^2}$$

$$= 2\|\nabla h - \nabla g\|_{L^2}^2$$

no line break

(For Langevin diffusion)

Applications to MCMC

Optimal control variates

Let $\tilde{h}_g = h - g$,

$$\begin{aligned}\mathcal{D}\tilde{h}_g &= \mathcal{D}h - \mathcal{D}g \\ &= -c_g + \eta\end{aligned}$$

Thus \tilde{h}_g is the solution to Poisson's equation with forcing function c_g .

Asymptotic variance of the new estimator:

$$\begin{aligned}\gamma_g^2 &= 2\langle \tilde{h}_g, \tilde{c}_g \rangle_{L^2} \\ &= 2\|\nabla h - \nabla g\|_{L^2}^2\end{aligned}$$

(For Langevin diffusion)

Can be minimized using ∇ -LSTD algorithms.

Applications to MCMC

Numerical Examples - Sample variance v Asymptotic variance

Sample variance v Asymptotic variance

$$\sigma^2 = \langle \tilde{c}, \tilde{c} \rangle_{L^2} = R(0)$$

- Sample variance

Applications to MCMC

Numerical Examples - Sample variance v Asymptotic variance

$$R(t) = \langle \tilde{c}, P^t \tilde{c} \rangle$$

Sample variance v Asymptotic variance

$$\sigma^2 = \langle \tilde{c}, \tilde{c} \rangle_{L^2} = R(0) \quad - \text{Sample Variance}$$

$$\gamma^2 = 2\langle h, \tilde{c} \rangle_{L^2} = \int_{-\infty}^{\infty} R(s)ds \quad - \text{Asymptotic variance}$$

Applications to MCMC

Numerical Examples - Sample variance v Asymptotic variance

Sample variance v Asymptotic variance

$$\sigma^2 = \langle \tilde{c}, \tilde{c} \rangle_{L^2} = R(0) \quad - \text{Sample Variance}$$

$$\gamma^2 = 2\langle h, \tilde{c} \rangle_{L^2} = \int_{-\infty}^{\infty} R(s)ds \quad - \text{Asymptotic variance}$$

Minimizing σ^2 is easier than minimizing γ^2 [Oates 14, Papamarkou 14]

~~Appropriate only if samples are i.i.d.~~

Minimizing σ^2 also minimizes γ^2 ?

Applications to MCMC

Numerical Examples - Sample variance v Asymptotic variance

Sample variance v Asymptotic variance

$$\sigma^2 = \langle \tilde{c}, \tilde{c} \rangle_{L^2} = R(0) \quad \text{- Sample variance}$$

$$\gamma^2 = 2\langle h, \tilde{c} \rangle_{L^2} = \int_{-\infty}^{\infty} R(s)ds \quad \text{- Asymptotic variance}$$

Minimizing σ^2 is easier than minimizing γ^2 [Oates 14, Papamarkou 14]



Appropriate only if samples are i.i.d.

Minimizing σ^2 also minimizes γ^2 ? NO !

Applications to MCMC

Numerical Examples - Sample variance vs Asymptotic variance

Example: Unadjusted Langevin algorithm (ULA)
 $c(x) \equiv x$



Applications to MCMC



Numerical Examples - ULA

Example: Unadjusted Langevin algorithm (ULA)

$$c(x) \equiv x, \rho \sim 0.5\mathcal{N}(-1, 0.4472) + 0.5\mathcal{N}(1, 0.4472)$$

pbt will
be better.

Same density as prev.
page?



Applications to MCMC



Numerical Examples - RWM

kill?

Example: Random walk Metropolis (RWM)

$$c(x) \equiv x, \rho \sim 0.5\mathcal{N}(-1, 0.4472) + 0.5\mathcal{N}(1, 0.4472)$$


Theoretical justification based on [Brosse et al. 19]



Applications to MCMC

Numerical Examples - ULA v RWM

Unadjusted Langevin algorithm (ULA) vs Random walk Metropolis (RWM)



?

Applications to MCMC

Numerical Examples - Logistic regression with RWM sampling

(other stuff list)

~~Example:~~ Logistic Regression for Swiss bank notes

$X \in \mathbb{R}^{200 \times 4}$ - Covariates measurements of four features of 200 bank notes.

$\{Y_i \in \{0, 1\}, 1 \leq i \leq 200\}$ - Binary response variables.

$\Theta \in \mathbb{R}^4$ - Regression coefficients for classification.



$$\rho(\Theta | \{X_i, Y_i\}_1^N) \propto$$

$$\exp \left(\underbrace{\sum_{i=1}^N \{Y_i \Theta^\top X_i - \log(1 + e^{\Theta^\top X_i})\}}_{\text{Likelihood}} - \underbrace{\frac{\Theta^\top \Sigma^{-1} \Theta}{2}}_{\text{Prior}} \right)$$

Applications to MCMC

Numerical Examples - Logistic Regression with RWM sampling

Example: Logistic Regression for Swiss bank notes
RWM sampling



Box plots of estimates of Θ_4 .

Conclusions

- Differential LSTD learning based approaches to approximate solution to Poisson's equation for the Langevin diffusion.
 - Finite dimensional basis.
 - RKHS.
- Two interesting applications
 - Asymptotic variance reduction in MCMC algorithms.
 - Gain function approximation in Feedback particle filter.
- Extended the approach to include reversible Markov chains.

References

-  A. Radhakrishnan, A. Devraj and S. Meyn, "Learning techniques for feedback particle filter design," *2016 IEEE 55th Conference on Decision and Control (CDC)*, Las Vegas, NV, 2016.
-  A. Radhakrishnan, S. Meyn, "Feedback particle filter design using a differential-loss reproducing kernel Hilbert space," *2018 American Control Conference (ACC)*, Milwaukee, WI, 2018.
-  S.P.Meyn, "Control Techniques for Complex Networks", *Cambridge University Press*, Dec 2007.
-  S. Henderson. *Variance Reduction Via an Approximating Markov Process*. PhD thesis, Stanford University, Stanford, California, 1997.
-  T. Yang, P. G. Mehta and S. P. Meyn, "Feedback Particle Filter," in *IEEE Transactions on Automatic Control*, vol. 58, no. 10, pp. 2465-2480, Oct. 2013.
-  A. M. Devraj and S. P. Meyn, "Differential LSTD learning for value function approximation," *2016 IEEE 55th Conference on Decision and Control (CDC)*, Las Vegas, NV, 2016.
-  D.X. Zhou, "Derivative reproducing properties for kernel methods in learning theory," *Journal of Computational and Applied Mathematics*, Vol. 220, Issues 1?2, 2008.

Thank You!

Questions?