

SS08_pemb5635

December 4, 2022

```
[1]: import pandas as pd
import numpy as np
import matplotlib.pyplot as plt # read in the plotting library matplotlib and
    ↪ call it plt
import statsmodels.api as sm # import stats package
```

Can you come up with a qualitative explanation for why superconductivity might break down under

First we must understand how temperature breaks up superconductivity. Based on the fact that nature favours lower energy states. There is a band gap energy (Δ) between coupled paired electrons and normal state electrons. If temperature (kT) is comparable to Δ , essentially exciting the paired electrons, super conductivity breaks. In terms of fermi surface in k space, to illustrate band gap energy Δ , one can draw another surface surrounding the original fermi surface, having a gap corresponding to Δ . There are no states available in between those surfaces. Increasing T will increase the surface and at a certain T , two surfaces will meet, corresponding to breaking of symmetry. To explain why superconductivity breaks at high current, we use the fermi surface logic again. When current is turned on, fermi surface, as a sphere, gains a net momentum shifting it. For high current densities, the sphere will be shifted until it contacts the other surface at Δ gap energy, i.e. breaking superconductivity.

1 4.1 Observation of the normal-superconducting transition using the resistivity of tin wire

1.1 4.1.1 Preliminary measurements

1. Why is the four wire method preferable to using just two wires? To avoid contact messing up resistance measurements
2. Why is it necessary to reverse the current through the sample? Applying current on the wire creates a Temperature gradient. By reversing current direction, one can cancel out that effect. Temperature is an important factor for this experiment
3. How can one check that the current used for the resistance measurement is not heating the sample? Just leave it and see if heats up. What is important is the rate of temperature difference.

1.2 4.1.1 The normal-superconducting transition

```
[2]: #Pressure to Temperature Converter
def pressure2temp(p_mbar): ## converts pressure of helium in mbar to
    ↪temperature in K
    x = np.log10(p_mbar) ##note that log10() is used as opposed to log()
    T = 1.24177 + 0.23793*(x) + 0.36207*(x**2) - 0.33188*(x**3) + 0.
    ↪20738*(x**4) - 0.05294*(x**5) + 0.00552*(x**6)
    return T

[3]: #Read File
ResistivityDataRemote = pd.read_csv("../data/resistivity/
    ↪ResistivityDataRemote.txt", "\t", names=["Helium Pressure", "V+", "V-"])
#Drop lines
ResistivityDataRemote = ResistivityDataRemote.drop(ResistivityDataRemote.
    ↪index[range(0,6)])

#Define Variables
current = 1.002; #Value given in data file. Used in Lower regime

helium_pressure = ResistivityDataRemote['Helium Pressure'].values.astype(float)
v_plus = ResistivityDataRemote['V+'].values.astype(float)
v_minus = ResistivityDataRemote['V-'].values.astype(float)

Note that -V just indicates that current direction is flipped


[4]: #Calculation
helium_temp = pressure2temp(helium_pressure)
v_average = (v_plus - v_minus)/2 #Note the - in front of v_minus



resistance_plus = v_plus/current
resistance_minus = v_minus/current
resistance_average = v_average/current

[5]: #Plot
plt.plot(helium_temp, resistance_plus)
## plt.title('Pressure against V')
plt.xlabel('Helium Temperature (K)') # Plot a label on x axis of Xlabel on graph
plt.ylabel('Resistance Plus (Ohms)') # Plot a label on x axis of Xlabel on graph
plt.show()

plt.plot(helium_temp, resistance_minus)
## plt.title('Pressure against V')
plt.xlabel('Helium Temperature (K)') # Plot a label on x axis of Xlabel on graph
plt.ylabel('Resistance Minus (Ohms)') # Plot a label on x axis of Xlabel on
    ↪graph
plt.show()
```

```
plt.plot(helium_temp, resistance_average)
## plt.title('Pressure against V')
plt.xlabel('Helium Temperature (K)') # Plot a label on x axis of Xlabel on graph
plt.ylabel('Resistance Average (Ohms)') # Plot a label on y axis of Ylabel on graph
plt.show()
```





[6]: *#Calculate Linear Fit for the transition of Resistance*

```

##Selecting transition range
helium_temp_trans = []
resistance_plus_trans = []
for i in range(len(resistance_plus)):
    if resistance_plus[i] > 0.02 and resistance_plus[i] < 0.08:
        helium_temp_trans.append(helium_temp[i])
        resistance_plus_trans.append(resistance_plus[i])
helium_temp_trans = np.array(helium_temp_trans)
resistance_plus_trans = np.array(resistance_plus_trans)

## add linear fit
X = sm.add_constant(helium_temp_trans) # add a constant to fit
results = sm.OLS(resistance_plus_trans, X).fit() # save results of fit


```

```

[7]: #Plot with Fit & Summary
plt.plot(helium_temp, resistance_plus)
plt.plot(helium_temp_trans, results.params[0]+results.
    ↪params[1]*helium_temp_trans, 'r' , label='fitted_line')
plt.title('Resistance against Temperature')
plt.xlabel('Temperature (K)') # Plot a label on x axis of Xlabel on graph
plt.ylabel('Resistance (Ohms)') # Plot a label on x axis of Xlabel on graph
plt.show()

print(results.summary()) # print results out to screen

```



OLS Regression Results

Dep. Variable:	y	R-squared:	0.984
Model:	OLS	Adj. R-squared:	0.981
Method:	Least Squares	F-statistic:	477.1
Date:	Wed, 20 Jan 2021	Prob (F-statistic):	2.04e-08
Time:	18:19:58	Log-Likelihood:	46.697
No. Observations:	10	AIC:	-89.39
Df Residuals:	8	BIC:	-88.79
Df Model:	1		
Covariance Type:	nonrobust		

	coef	std err	t	P> t	[0.025	0.975]
const	-3.4412	0.160	-21.520	0.000	-3.810	-3.072
x1	0.9450	0.043	21.842	0.000	0.845	1.045

Omnibus:	2.211	Durbin-Watson:	1.873
Prob(Omnibus):	0.331	Jarque-Bera (JB):	0.893
Skew:	-0.181	Prob(JB):	0.640
Kurtosis:	1.581	Cond. No.	791.

Notes:

[1] Standard Errors assume that the covariance matrix of the errors is correctly specified.

/usr/local/anaconda3/lib/python3.7/site-packages/scipy/stats/stats.py:1604:

UserWarning: kurtosistest only valid for n>=20 ... continuing anyway, n=10
 "anyway, n=%i" % int(n))

```
[8]: #Calculating the critical temperature
R_min = min(resistance_plus)
R_max = max(resistance_plus)
Rc = (R_min + R_max)/2 #Selecting around the middle point of the transition
    ↪range

Tc = (Rc- results.params[0])/results.params[1]
print(Tc)
```

3.693474566320118

Note that Critical(Transition) Temperature isn't really defined precisely. A bit arbitrary. We chose the middle point Why is the width of the transition finite? This phenomenon is a 2nd order Gibbs phase transition. Finite width comes from impurities of the material used.

1.3 4.1.3 The critical magnetic field

```
[9]: file=pd.read_csv("../data/resistivity/ProbeA3mB", "\t")
file
pressure2temp(3)
```

```
[9]: 1.411170942905004
```

From script, we know that shunt giving 0.5V/A provides 0.018T/A at the center. (Conversion of Volt to Tesla)

```
[10]: #Grand scheme
pressureList = [3 ,21, 43, 58, 74, 92, 130, 170, 210, 250, 290, 330, 370, 400,
↪430, 450, 473, 500, 520, 540, 560, 570, 595, 610, 630, 890 ] #1010 not using

ratio = 0.018/0.5


Bc = []

for num in pressureList:
    filename = "../data/resistivity/ProbeA"+str(num)+"mB"
    df = df = pd.read_csv(filename, "\t",names=["Acquisition Time (ms)", "Shunt_
↪Voltage (V)", "Sample Voltage (V)","Integrated Sample Voltage (V)","N/A","N/
↪A2"])
    df = df.drop(df.index[range(0,2)])
    B = df["Shunt Voltage (V)"].values.astype(float)*ratio
    V = df["Sample Voltage (V)"].values.astype(float)
    B_trans = []
    V_trans = []
    for i in range(len(V)):
        if V[i] > 0.00001 and V[i] < 0.00008:
            B_trans.append(B[i])
            V_trans.append(V[i])
    B_trans = np.array(B_trans)
    #B_midpoint = ( B_trans[0]+ B_trans[len(B_trans)-1] )*0.5
    if len(B_trans) != 0:
        Bc.append(( B_trans[0]+ B_trans[len(B_trans)-1] )*0.5 )
```

```
[11]: #Plot Bc and T
temp = []
for i in range(len(pressureList)):
    temp.append(pressure2temp(pressureList[i]))
temp_c = temp[:len(Bc)]

temp_square = [element**2 for element in temp_c]
temp_square = np.array(temp_square)
```

```
plt.plot(temp_square,Bc)
plt.xlabel('Temperature^2 (K^2)') # Plot a label on x axis of Xlabel on graph
plt.ylabel('Magnetic Field (T)') # Plot a label on x axis of Xlabel on graph
plt.show()
```



```
[12]: ## add linear fit
Lfit = sm.add_constant(temp_square) # add a constant to fit
results = sm.OLS(Bc, Lfit).fit() # save results of fit
print(results.summary())

#Add fit to graph
plt.plot(temp_square, results.params[0]+results.params[1]*temp_square, 'r', label='fitted_line')
plt.plot(temp_square,Bc)
plt.xlabel('Temperature^2 (K^2)') # Plot a label on x axis of Xlabel on graph
plt.ylabel('Magnetic Field (T)') # Plot a label on x axis of Xlabel on graph
plt.legend()
plt.show()
```

OLS Regression Results

```
=====
Dep. Variable:          y      R-squared:                0.998
Model:                  OLS    Adj. R-squared:            0.998
Method:                 Least Squares    F-statistic:          1.153e+04
Date:                   Wed, 20 Jan 2021    Prob (F-statistic):    2.01e-31
```



Time: 18:19:59 Log-Likelihood: 159.30
 No. Observations: 24 AIC: -314.6
 Df Residuals: 22 BIC: -312.2
 Df Model: 1
 Covariance Type: nonrobust

	coef	std err	t	P> t	[0.025	0.975]
const	0.0284	0.000	146.089	0.000	0.028	0.029
x1	-0.0020	1.9e-05	-107.393	0.000	-0.002	-0.002

Omnibus: 4.972 Durbin-Watson: 1.512
 Prob(Omnibus): 0.083 Jarque-Bera (JB): 3.020
 Skew: -0.766 Prob(JB): 0.221
 Kurtosis: 3.822 Cond. No. 29.7

Notes:

[1] Standard Errors assume that the covariance matrix of the errors is correctly specified.



```
[13]: ##Procedure for one pressure
df = pd.read_csv("../data/resistivity/ProbeA58mB", "\t", names=["Acquisition_
↳Time (ms)", "Shunt Voltage (V)", "Sample Voltage (V)", "Integrated Sample_
↳Voltage (V)", "N/A", "N/A2"])
```

```

df = df.drop(df.index[range(0,2)])
df

#plot B versus time
ratio = 0.018/0.5

t = df["Acquisition Time (ms)"].values.astype(float)
B = df["Shunt Voltage (V)"].values.astype(float)*ratio
V = df["Sample Voltage (V)"].values.astype(float)
iV = df["Integrated Sample Voltage (V)"].values.astype(float)

plt.plot(B, V)
## plt.title('Pressure against V')
plt.xlabel('Magnetic Field (T)') # Plot a label on x axis of Xlabel on graph
plt.ylabel('Sample Voltage (V)') # Plot a label on x axis of Xlabel on graph
plt.show()



#Extract Bc
B_trans = []
V_trans = []
for i in range(len(V)):
    if V[i] > 0.00001 and V[i] < 0.00008:
        B_trans.append(B[i])
        V_trans.append(V[i])

B_trans = np.array(B_trans)

B_midpoint = ( B_trans[0]+ B_trans[len(B_trans)-1] )*0.5

plt.plot(B,V)
plt.plot(B_trans,V_trans)
plt.xlabel('Magnetic Field (T)') # Plot a label on x axis of Xlabel on graph
plt.ylabel('Sample Voltage (V)') # Plot a label on x axis of Xlabel on graph
plt.show()

```



```
[14]: results.params[0]
```

```
[14]: 0.028401566892467554
```

```
[15]: #Calculate Tc
Tc = np.sqrt(results.params[0]/abs(results.params[1]))
print("Critical Temperature: "+str(Tc)+"K")
```

Critical Temperature: 3.7305713495770574K

Questions: Can you explain the shape of the transition between superconducting and normal behaviour? Until $B=0.014\text{T}$, for all temperature, material is superconducting. After that point, there is a small increase in resistance (sample voltage, they are similar as current is 1), and superconductivity is broken around critical magnetic field. How can we explain the small increment before critical magnetic field? Think about a sphere in uniform magnetic field. Due to boundary conditions, magnetic flux density is squeezed together at the edges of the sphere, increasing the density. So, flux density at edges is actually larger than applied field. For cylinder, it gives a factor of 2. Now, the probe for this experiment is made of a coil of wire wrapped around a rectangular material. The wires on the sides that are perpendicular to applied field will experience the above effect. So, at $B=0.014\text{T}$, the two sides will experience “early” as if there is 0.028T . This tendency can be seen in graphs. Does the transition look different for upsweeps and downsweeps of the magnetic field? This is a fundamental property of superconductors. When downsweeping, around critical magnetic field, a loop of current is produced around the superconducting path of the material, as there is no resistance. Then, the decrement of magnetic field is countered by the increase of current in superconductors because induced current is the rate of flux change. So after critical magnetic field, magnetic field inside the superconductor will not decrease. There will still be large flux density inside. Won’t see the tendency of the left side of the above graph.

1.4 4.2 The Meissner Effect in Tin

```
[16]: ddf = pd.read_csv("../data/meissner_effect/ProbeB80mbar","\t", names = [
    ↪ ["Acquisition Time (ms)", "Shunt Voltage (V)", "Sample Voltage (V)",
    ↪ ("Integrated Sample Voltage (V)", "N/A")])
ddf = ddf.drop(ddf.index[range(0,2)])
ddf
```

```
[16]: Acquisition Time (ms) Shunt Voltage (V) Sample Voltage (V) \
2          9933750      0.0113017242      5.7455022E-6
3          9934000      0.0232247586      2.1404448E-6
4          9934250      0.036030491      4.1537169E-6
5          9934500      0.0477310741      3.7277462E-6
6          9934750      0.060276027      4.0416193E-6
..          ...          ...          ...
238         9992750      0.0516333573     -2.11892674E-5
239         9993000      0.0393370663     -2.11533963E-5
240         9993250      0.0268615244     -1.68712794E-5
241         9993500      0.0145754424     -1.47638501E-5
242         9993750      0.0024378459     -1.18358686E-5
```

Integrated Sample Voltage (V) N/A

```

2          4.5455022E-6 NaN
3          5.4859469E-6 NaN
4          8.4396638E-6 NaN
5          1.096741E-5 NaN
6          1.38090293E-5 NaN
..          ... ..
238         1.299903031E-4 NaN
239         1.076369068E-4 NaN
240         8.95656274E-5 NaN
241         7.36017773E-5 NaN
242         6.05659088E-5 NaN

```


[241 rows x 5 columns]

```

[17]: B = ddf["Shunt Voltage (V)"].values.astype(float)*ratio
      iV = ddf["Integrated Sample Voltage (V)"].values.astype(float)
      V = ddf["Sample Voltage (V)"].values.astype(float)

      plt.plot(B,iV)
      plt.xlabel('Magnetic Field (T)') # Plot a label on x axis of Xlabel on graph
      plt.ylabel('Integrated Sample Voltage (V)') # Plot a label on x axis of Xlabel_
      ↪on graph
      plt.show()


```



```
[18]: #Sweeping current at 4.2K
print(pressure2temp(1010))
ddf_4K = pd.read_csv("../data/meissner_effect/ProbeB1010mbar","\t", names = [
    ↪ "Acquisition Time (ms)", "Shunt Voltage (V)", "Sample Voltage (V)", "Integrated Sample Voltage (V)", "N/A"])
ddf_4K = ddf_4K.drop(ddf_4K.index[range(0,2)])
B_4K = ddf_4K["Shunt Voltage (V)"].values.astype(float)*ratio
iV_4K = ddf_4K["Integrated Sample Voltage (V)"].values.astype(float)
V_4K = ddf_4K["Sample Voltage (V)"].values.astype(float)

plt.plot(B_4K,iV_4K)
plt.xlabel('Magnetic Field (T)') # Plot a label on x axis of Xlabel on graph
plt.ylabel('Integrated Sample Voltage (V)') # Plot a label on x axis of Xlabel
    ↪ on graph
plt.show()
```

4.221493495675776



Note that it is a straight line. It is because the temperature is above critical temperature, so no superconductivity can be seen.


```
[19]: #Seeing where transition part is
ddf = pd.read_csv("../data/meissner_effect/ProbeB80mbar",
    ↪ "\t", names=["Acquisition Time (ms)", "Shunt Voltage (V)", "Sample Voltage (V)", "Integrated Sample Voltage (V)", "N/A"])
```

```

ddf = ddf.drop(ddf.index[range(0,2)])
B = ddf["Shunt Voltage (V)"].values.astype(float)*ratio
iV = ddf["Integrated Sample Voltage (V)"].values.astype(float)
B_trans = []
iV_trans = []
before = iV[0]
for i in range(int(len(iV)/2)):
    after = iV[i]
    if after-before >= 0.0001:
        B_trans.append(B[i])
        iV_trans.append(iV[i])
    before = after
B_trans = np.array(B_trans)

plt.plot(B,iV)
plt.plot(B_trans,iV_trans,label='transition region')
plt.xlabel('Magnetic Field (T)') # Plot a label on x axis of Xlabel on graph
plt.ylabel('Integrated Sample Voltage (V)') # Plot a label on x axis of Xlabel_
↳ on graph
plt.legend()
plt.show()

```



[20]: #Grand Scheme

```

pressure_list_m = [
    ↪[2,20,41,58,80,98,132,165,190,220,238,270,320,360,390,410,440,460,470,490,525,540,590,690,7
Bc_m = []



for num in pressure_list_m:
    filename = "../data/meissner_effect/ProbeB"+str(num)+"mbar"
    ddf = pd.read_csv(filename, "\t",names=["Acquisition Time (ms)", "Shunt_
    ↪Voltage (V)", "Sample Voltage (V)","Integrated Sample Voltage (V)","N/A"])
    ddf = ddf.drop(ddf.index[range(0,2)])
    B = ddf["Shunt Voltage (V)"].values.astype(float)*ratio
    iV = ddf["Integrated Sample Voltage (V)"].values.astype(float)
    plt.plot(B,iV)
    plt.xlabel('Magnetic Field (T)') # Plot a label on x axis of Xlabel on graph
    plt.ylabel('Integrated Sample Voltage (V)') # Plot a label on x axis of
    ↪Xlabel on graph
    B_trans = []
    iV_trans = []
    before = iV[0]
    for i in range(int(len(iV)/2)):
        after = iV[i]
        if after-before >= 0.0001:
            B_trans.append(B[i])
            iV_trans.append(iV[i])
        before = after
    B_trans = np.array(B_trans)
    #B_midpoint = ( B_trans[0]+ B_trans[len(B_trans)-1] )*0.5
    if len(B_trans) != 0:
        Bc_m.append(( B_trans[0]+ B_trans[len(B_trans)-1] )*0.5 )
Bc_m = np.array(Bc_m)

plt.show()

temp_m = temp[:len(Bc_m)]

temp_square_m = [elements**2 for elements in temp_m]
temp_square_m = np.array(temp_square_m)
plt.plot(temp_square_m,Bc_m)
plt.ylabel('Magnetic Field (T)') # Plot a label on x axis of Xlabel on graph
plt.xlabel('Temperature^2(K^2)') # Plot a label on x axis of Xlabel on graph
plt.show()

```

How to explain the shape of the transition between superconducting and normal behaviour? note that integrated sample voltage represents flux inside the material. we can see a very small increment of flux around 0.014T as explained previously. Another interesting part of this graph is

the downsweep of magnetic field. One would expect flux to be constant despite the decrease of B because, as explained previously, as soon as the material hits critical field, a loop of current is induced and the current will cancel out the effect of decreasing magnetic field. However that is not the case in here. The reason lies in Tin. Tin has an abnormal tendency to show semiconductor characteristics at low temperature. (Tin's phase transition). The tiny dip during downsweep is due to the small portion of Tin acting metallic but soon to be dominated by semi-conducting attributes.

```
[21]: #Add fit

## add linear fit
Lfit_m = sm.add_constant(temp_square_m) # add a constant to fit
results_m = sm.OLS(Bc_m, Lfit_m).fit() # save results of fit
print(results_m.summary())

#Add fit to graph
plt.plot(temp_square_m, results_m.params[0]+results_m.params[1]*temp_square_m,
         color='r', label='fitted_line')
plt.plot(temp_square_m, Bc_m)
plt.xlabel('Temperature^2 (K^2)') # Plot a label on x axis of Xlabel on graph
plt.ylabel('Magnetic Field (T)') # Plot a label on y axis of Ylabel on graph
plt.legend()
plt.show()
```

OLS Regression Results

```
=====
Dep. Variable:          y      R-squared:          0.993
Model:                  OLS    Adj. R-squared:       0.993
Method:                 Least Squares    F-statistic:      2777.
Date:                   Wed, 20 Jan 2021    Prob (F-statistic): 4.62e-22
Time:                   18:20:00    Log-Likelihood:    126.36
No. Observations:      21    AIC:              -248.7
Df Residuals:          19    BIC:              -246.6
Df Model:               1
Covariance Type:       nonrobust
=====
```


	coef	std err	t	P> t	[0.025	0.975]
const	0.0307	0.000	80.790	0.000	0.030	0.032
x1	-0.0021	3.95e-05	-52.693	0.000	-0.002	-0.002

```
=====
Omnibus:                0.648    Durbin-Watson:          0.521
Prob(Omnibus):          0.723    Jarque-Bera (JB):        0.289
Skew:                   0.284    Prob(JB):                0.865
Kurtosis:               2.910    Cond. No.:               27.4
=====
```

Notes:

[1] Standard Errors assume that the covariance matrix of the errors is correctly

specified.



```
[22]: #Compare with previous part
      #print(results.summary())
      #print(results_m.summary())

      B_0_error = np.absolute((results.params[0]-results_m.params[0])*100/results_m.
        ↪params[0])
      print("B_0 percentage difference: "+str(B_0_error)+"%")
      T_c_m = np.sqrt(-results_m.params[0]/results_m.params[1])
      print("Critical Temperature: "+str(T_c_m)+"K")
      T_c_diff = np.absolute((Tc-T_c_m)/T_c_m)
      print(T_c_diff)
```

```
B_0 percentage difference: 7.607613149575059%
Critical Temperature: 3.8440181250919583K
0.02951254958304521
```

```
[23]: #gamma
      gamma = (2/(4*np.pi*10**(-7)))*(results_m.params[0]/T_c_m)**2
      print(gamma)
      T_c_m
```

```
101.77977944186577
```

[23] : 3.8440181250919583

[]:

[]:

[]: