1. The probability of a leap year selected at random contain 53						
Sunday is:						
	(b) 1/7					
2. A bag contains	s 3 red and 2 bl	ue marbles. A r	narble is drawn at			
random. The prol	bability of draw	ing a black ball	is:			
(a) 3/5	(b) 2/5	(c) 0/5	(d) 1/5			
3. The probabilit	y that it will raii	n tomorrow is 0	.85. What is the			
probability that it	will not rain to	morrow				
(a) 0.25	(b) 0.145	(c) 3/20	(d) none of these			
4. What is the pr	obability that a	number selecte	ed from the numbers			
(1, 2, 3,,15) is a multiple o	f 4?				
(a) 1/5	(b) 4/5	(c) 2/15	(d) 1/3			
5. What are the	total outcomes	when we throw	three coins?			
(a) 4	(b) 5	(c) 8)	(d) 7			
6. The probability	ty that a prime	number selecte	d at random from the			
numbers (1,2,3, .	35) is :					
(a) 12/35	(b) 11/35	(c) 13/35	(d) none of these			
7. The sum of th						
(a) 2	(b) 1 (c) (0 (d) none	e of these.			
			e the correct answer			
for that which is	not possible.					
(a) 0.15	(b) 2/7	(c) 7/5	(d) none of these.			
			n the probability of			
getting at least tv	wo heads, is:					
(a) 1/4	(b) 3/8	(C) $\frac{1}{2}$	(d) 1/8			
10. A letter is ch						
♦ ASSASSINATION	ON�. The prob	ability that the	letter chosen has:			
(a) 6/13	(b) 7/13	(c) 1	letter chosen has: (d) none of these.			
11. A dice is thro	wn. Find the pro	obability of gett	ting an even number.			
(A) 2/3	(B) 1	(C) 5/6	D) 1/2			
12. Two coins are thrown at the same time. Find the probability of						
getting both heads.						
(A) 3/4 (B) 1/4	(C) 1/2	(D) 0				
			1 1 110.			
13. Two dice are thrown simultaneously. The probability of getting a						

1

sum of 9 is:

(A) 1/10	(B) 3/10	(C) 1/9	(D) 4	./9			
14. 100 cards are numbered from 1 to 100. Find the probability of getting a prime number.							
(A) 3/4		(C) 1/4	(D) 29/100			
_	a blue ball is do n a bag is:			If the probability en the number of			
		this box. The					
17. Cards marked with numbers 2 to 101 are placed in a box and mixed thoroughly. One card is drawn from this box randomly, then the probability that the number on card is a perfect square. (A) 9/100 (B) 1/10 (C) 3/10 (D) 19/100							
18. What is the probability of getting 53 Mondays in a leap year? (A) 1/7 (B) 53/366 (C) 2/7 (D) 7/366							
 19. A card is drawn from a well shuffled deck of 52 cards. Find the probability of getting a king of red suit. (A) 1/26 (B) 3/26 (C) 7/52 (D) 1/13 							
20. A game of chance consists of spinning an arrow which is equally likely to come to rest pointing to one of the number 1,2,312, then the probability that it will point to an odd number is: (A) 1/6 (B) 1/12 (C) 7/12 (D) 5/12							
21. A game consists of tossing a one rupee coin 3 times and noting its outcome each time. Aryan wins if all the tosses give the same result i.e. three heads or three tails and loses otherwise. Then the probability that Aryan will lose the game. (A) 3/4 (B) 1/2 (C) 1 (D) 1/4							

same birthda	Kajal are friends. ay is the same birt (B) 31/365	hday is:				
2. Then the	er x is chosen at raprobability that x ² (C) 3/5	< 2 is?	numbers -2, -1, 0 , 1,			
a marble is ored is 2/3, th		rom the jar, the p white marbles in	d others are white. If robability that it is the jar is:			
Then the pro	er is selected at rai bbability that it is a B) 4/25 (C) 1/25	multiple of 3 and	50 natural numbers. d 4 is:			
26. Consider a dice with the property that that probability of a face with n dots showing up is proportional to n. The probability of face showing 4 dots is?						
a) $\frac{1}{7}$	b) $\frac{5}{42}$	c) $\frac{1}{21}$	d) 21			
	_		hes are 50, 70, 82,			
	Fhe standard devia b) 25.49		d) 25.69			
28. Find median and mode of the messages received on 9 consecutive days 15, 11, 9, 5, 18, 4, 18, 13, 17.						
a) 13, 15	b) 13, 18)	c) 18, 15	d) 13, 16			
29. A coin is tossed up 4 times. The probability that tails turn up in						
3 cases is a) $\frac{1}{2}$ 30. X is a value a) 8	b) $^1\!/_3$ riate between 0 an	c) 1/4 ad 3. The value of c) 27	d) $\frac{1}{6}$ E(X ²) is			
31. The random variables X and Y have variances 0.2 and 0.5 respectively. Let Z= 5X-2Y. The variance of Z is?						

32.Out of the following values, which one is not possible in probability? a) $P(x) = 1$ b) $\sum x P(x) = 3$ c) $P(x) = 0.5$ d) $P(x) = -0.5$						
33.If E(x) =	2 and E(z) = 4, b) 6		•	Insufficient	data	
34.The cov	ariance of two i	ndepende	nt random va	riable is		
a) 1	b) 0	c) - 1	d) Undefined		
) = k² – 8 then, t b) 1) Insufficient	data	
	0.5 and x = 4, t b) 0.5	hen E(x) = c) 4) 2		
37.In a discrete probability distribution, the sum of all probabilities is always? a) 0 b) Infinite c) 1 d) Undefined						
ŕ	robability of hitt	•	ŕ			
a) 0.4, 0.24	b) 0.6, 0	.24	c) 0.4, 0.1	6 d) 0.6,	0.16	
39.If the probability that a bomb dropped from a place will strike the target is 60% and if 10 bombs are dropped, find mean and variance? a) 0.6, 0.24 b) 6, 2.4 c) 0.4, 0.16 d) 4, 1.6						
 40. Find the mean of tossing 8 coins. a) 2 b) 4 c) 8 d) 1 41. What is the mean and variance for standard normal distribution? 						

c) 5

d) 7

a) 3

b) 4

a) Mean is 0 and variance is 1 b) Mean is 1 and variance is 0 c) Mean is 0 and variance is ∞ d) Mean is ∞ and variance is 0								
42. Variance of a random variable X is given by a) $E(X)$ b) $E(X2)$ c) $E(X2)$ – $(E(X))2$ d) $E(X)$ 2								
	43.Mean of a random variable X is given by a) E(X)							
44.Mean of a constant 'a' is a) 0								
45. Variance of a constant 'a' is a) 0								
46.Find the mean and variance of X?								
x 0 1 2 3 4								
	f(x)	1/9	2/9	3/9	2/9	1/9		
a) 2	, 4/ 3	b) 3	, 4/3		c) 2, 2/3		d) 3, 2/3	

47. Find the expectation of a random variable X?

	X	0	1	2	3	
	f(x)	1/6	2/6	2/6	1/6	
a) ().5		b) 1.5		c) 2.5	d) 3.5

48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by

b) npq c) np2q

d) npq2

- 49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use **Binomial Distribution.**
- a) P(X = x) = nCx px qx
- b) P(X = x) = nCx px q(n-x)
- c) P(X = x) = xCn qx p(n-x)
- d) P(x = x) = xCn pn qx
- 50. If 'p', 'q' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard **Deviation?**

- a) \sqrt{np} b) \sqrt{pq} c) (np)2 d) \sqrt{npq}