

Cetamol Reva

Tablets



Composition:

Each Cetamol Reva 500 Tablet contains Paracetamol (Acetaminophen) 500 mg.

Each Cetamol Reva 1000 Tablet contains Paracetamol (Acetaminophen) 1000 mg.

Mechanism of Action:

Paracetamol is an effective, analgesic and antipyretic. It works by inhibiting the synthesis of prostaglandins in the central nervous system and to a lesser extent through a peripheral effect by blocking the generation of pain impulses.

It is possible that paracetamol's antipyretic effect is due to its central effect on the hypothalamic thermoregulatory center.

Pharmacokinetics:

Paracetamol is rapidly absorbed from the gastrointestinal tract. The concentration in plasma reaches a peak in 30 minutes to 2 hours after ingestion. It is metabolized in the liver and excreted in the urine mainly as the glucuronide and sulphate conjugates. Less than 5% of the administered dose is removed unchanged. The elimination half-life varies from about 1 to 4 hours.

Indications:

Cetamol Reva is a mild pain reliever and antipyretic and is recommended for most pain and fever conditions such as, headache and migraine, toothache, neuralgia, colds and flu, sore throats, back pain, rheumatic pain and dysmenorrhea.

Contraindications:

- Hypersensitivity to Paracetamol, or to any other components of the product.
- Patients with liver and kidney damage.

Warnings & Precautions:

Caution should be exercised when paracetamol is administered to patients with hepatic or renal impairment.

Pregnancy & Lactation:

Pregnancy: Available data on the use of paracetamol in pregnant women indicate that there are no toxic (deform) effects in the newborn. Paracetamol can be used during pregnancy if necessary, but it should be used at the lowest effective dose, for the shortest possible time and as frequently as possible.

Lactation: Paracetamol is excreted in breast milk, but not in large quantities. The available data show that it is not contraindicated in this case.

Drug Interactions:

-Cholestyramine: The speed of absorption of paracetamol is reduced by cholestyramine, therefore, cholestyramine should not be taken within an hour of taking paracetamol.

-Warfarin: The risk of bleeding may be increased

when it uses with warfarin and other coumarins in prolonged regular daily.

- Chloramphenicol: increased plasma concentration of chloramphenicol.

- Metoclopramide and domperidone: These drugs may increase the absorption of paracetamol when one of them is used concomitantly with paracetamol.

Side Effects:

Adverse effects of paracetamol are rare but hypersensitivity including skin rash may occur. There have been very rare reports of blood dyscrasias including thrombocytopenia.

Driving & Using Machines:

Paracetamol has no influence on the ability to drive and use machines.

Dosage & Administration:

Children 10-15 years: 1 tablet of Cetamol Reva 500, every 4 hours as needed (with a maximum of 4 tablets in 24 hours). It is not recommended for children under 10 years of age.

Adults and children over 16 years: 1-2 tablets of Cetamol Reva 500, up to four times daily as needed (up to a maximum of 8 tablets in 24 hours). Or half tablet to one tablet Cetamol Reva 1000, orally 1-3 times a day.

- These doses should not be given frequently and no more than 4 doses should be given in 24 hours.

- The dose should not exceed 4g in 24 hours.

Overdose:

Hepatic damage can occur in adults (if they took 6g or more of paracetamol) and in children (if they took more than 150 mg/kg of paracetamol).

Treatment: Prompt treatment is essential in managing paracetamol overdose.

The patient should be referred to hospital urgently to obtain immediate medical attention.

The (N-acetylcysteine) antidote is given intravenously within 24 hours of taking paracetamol. If vomiting is not present, methionine may be given orally as a suitable alternative (out of hospital).

Storage:

Keep out of reach of children. Store below 25°C.

Packaging:

Each Cetamol Reva 500 carton contains 20 tablets in two blister strips.

Each Cetamol Reva 500 gross contains 500 tablets in 50 blister strips.

Each Cetamol Reva 1000 carton contains 20 tablets in two blister strips.

-A medication is a product which affects your health, and its consumption contrary to instructions is dangerous for you.

- Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medication.

- The doctor and the pharmacist experts in medicine, its benefits and risks.

- Do not repeat the same prescription without consulting your doctor.

KEEP THE MEDICAMENTS OUT OF REACH OF CHILDREN

Council of Arab Health Ministers- Union of Arab Pharmacies

Reva Pharmaceutical Industry - Syria