

DIMESTIL Oral Drops



Composition:

Each 1 ml DIMESTIL Oral Drops contains:
Dimetindene Maleate 1 mg.

Pharmacological Properties & Mechanism of Action:

Dimetindene Maleate is a H_1 -receptors antagonist with high affinity for these receptors.

In addition, Dimetindene Maleate has shown a weak anticholinergic activity. It considerably reduces the hyper permeability of the capillaries which are associated with immediate hypersensitivity reactions.

Pharmacokinetics:

The bioavailability of Dimetindene Maleate taken as oral drops is approximately 70%. Peak serum concentrations of Dimetindene Maleate are reached within two hours. Approximately 90% of Dimetindene Maleate is bound to human plasma proteins. Metabolic reactions involve the addition of hydroxyl and methoxyl to the compound. The half-life of elimination is about six hours. Dimetindene Maleate and its metabolites are eliminated by both the biliary and urinary routes.

Indications:

DIMESTIL drops are used in:

- Symptomatic treatment of drug and food allergies.
- Reducing itching caused by various factors: Endogenous pruritus, eczema, skin itching, insect bites, angioedema, chicken pox, urticaria.
- Reducing seasonal fever (hay fever) and perennial allergic rhinitis.

Contraindications:

- Hypersensitivity to Dimetindene Maleate or any components of the product.
- Infants less than one month of age, particularly premature.

Warnings & Precautions:

DIMESTIL is recommended to be used with caution in:

- Patients with glaucoma, urinary retention associated with prostatic hyperplasia, bronchial asthma, chronic obstructive pulmonary diseases, epilepsy.
- Infants aged less than 1 year: The sedative effect may be associated with episodes of sleep apnoea.
- DIMESTIL are ineffective in case of itching associated with cholestasis.

Pregnancy & Lactation:

Pregnancy: Pregnancy category B. DIMESTIL can be administration during pregnancy only according to the doctor's recommendations.

Lactation: The use of DIMESTIL is not recommended during breast feeding.

Drug Interactions:

- The sedative effect of central nervous system depressants, such as sedatives, hypnotics and alcohol may be enhanced when used concomitantly with DIMESTIL.

- Coadministration of MAOIs may increase the anticholinergic and CNS depressant effects of antihistamines; therefore, simultaneous use is not recommended.

- Concomitant administration of tricyclic antidepressants and anticholinergic drugs may enhance the anticholinergic effect of antihistamines, increasing the risk of urinary retention.

Side Effects:

The side effects include: Somnolence, nausea, dry mouth/dry throat, headache, dizziness, excitability, oedema, skin rash, muscle spasms and dyspnea.

Driving & Using Machines:

Demestil may decrease alertness in some patients. Therefore, it is not recommended for use while driving, operating machinery, or performing other tasks requiring alertness.

Dosage & Administration:

Adults and children over 12 years of age: 20-40 drops in the mouth, three times a day (1-2 ml) (20 drops = 1 ml contains 1 mg Dimetindene maleate).

Children:

The usual dose is about 0.1 mg/kg of body weight/day.

- Infants aged one month to one year: 3-10 drops in the mouth, three times daily.
- Children aged 1 to 3 years: 10-15 drops in the mouth, three times daily.
- Children aged 3 to 12 years: 15-20 drops in the mouth, three times daily.

Overdose:

Symptoms: Symptoms and signs of overdose with H_1 -antihistamines may include: CNS depression with drowsiness (mainly in adults), CNS stimulation and antimuscarinic effects (especially in children), including excitement, ataxia, hallucinations, tremors, convulsions, urinary retention and fever. Hypotension, coma and cardiorespiratory collapse may follow.

Treatment: There is no specific antidote for overdose with antihistamines. The usual emergency assistance must be undertaken, including activated charcoal, saline laxative and usual cardiorespiratory supportive measures. Stimulants should not be used; vasopressors may be used to treat hypotension.

Storage:

Keep out of reach of children. Store below 25 °C.

Packaging:

Each DIMESTIL Oral Drops carton box contains amber glass bottle of 20 ml.

*A medication is a product which affects your health, and it's consumption contrary to instructions is dangerous for you.
- Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medication.
*The doctor and the pharmacist experts in medicine, it's benefits and risks.
*Do not repeat the same prescription without consulting your doctor.
KEEP THE MEDICATIONS OUT OF REACH OF CHILDREN
Council of Arab Health Ministers, Union of Arab Pharmacies

Reva Pharmaceutical Industry - Syria

