

Dentyl Film coated Tablets



Composition:

Each DENTYL Film Coated Tablet contains:

Spiramycin 750.000 IU and Metronidazole 125 mg.

Each DENTYL Fort Film Coated Tablet contains:

Spiramycin 1.500.000 IU and Metronidazole 250 mg.

Mechanism of Action:

Spiramycin is a macrolide antibiotic and its bacteriostatic effect is due to the inhibition of protein synthesis by binding to the 50S subunit of bacterial ribosomes.

Metronidazole belongs to the imidazole group. The nitrate group in metronidazole acts as an electron acceptor, forming cytotoxic reducing complexes that bind with proteins and DNA, causing cell death. Metronidazole is active against various anaerobic bacteria and protozoa. The combination of spiramycin and metronidazole is indicated to provide a synergistic action in the treatment of dental oral infections.

Pharmacokinetics:

Spiramycin is rapidly but incompletely absorbed after oral administration, with an oral bioavailability of 33% to 39%. Plasma protein binding is low. It is metabolized in the liver with the formation of active metabolites. Over 80% of an administered dose excreted in the bile. Urinary excretion accounts for only 4% to 14% of an administered dose.

Metronidazole is readily absorbed after oral administration and bioavailability approaches 100%, and has a large volume of distribution in the body and it is metabolized in the liver. The elimination half life is about 8 hours and is longer in neonates and in patients with severe liver disease. The majority of a dose of Metronidazole is excreted in the urine mainly as metabolites; a small amount appears in the feces.

Indications:

DENTYL is highly effective in treating acute, chronic and relapsing oral infections such as:

- Oral infections caused by aerobic and anaerobic bacteria sensitive to spiramycin and metronidazole.
- Dental infections such as tooth abscesses, stomatitis, aphthous ulcers, pericoronitis and gingivitis.
- Parotitis and submandibular inflammation.
- Preventive treatment before and after surgical operations (oral and dental surgery).

Contraindications:

- Hypersensitivity to Spiramycin and/or Metronidazole.
- It is generally not recommended to use DENTYL in combination with disulfiram, alcohol, and medicinal products containing alcohol.
- DENTYL is not recommended for use in patients with glucose-6-phosphate-dehydrogenase deficiency due to the risk of hemolytic anemia.

Warnings & Precautions:

- The treatment should be discontinued in case of ataxia and vertigo.
- DENTYL should be used with caution in patients with blood dyscrasias or with active diseases of the central nervous system.
- DENTYL should be used with caution in patients with hepatic dysfunction.

Pregnancy & Lactation:

Pregnancy: DENTYL should not be used in the first trimester of pregnancy.

Lactation: Breast-feeding is not recommended to women taking DENTYL.

Drug Interactions:

- Spiramycin has been reported to inhibit the absorption of carbidopa and decrease Levodopa plasma levels. When necessary, patients should be closely monitored and the levodopa dosage levels adjusted.
- Drugs that may be affected by metronidazole include anticoagulants, disulfiram, hydantoins and lithium.
- Spiramycin may increase serum theophylline concentration.
- DENTYL can increase the effects of oral anticoagulants and digoxin.

Side Effects:

Common: Mild gastrointestinal disturbances as nausea, vomiting, diarrhea, unpleasant taste and darkening of urine may occur.

Rare: skin rash, urticaria, vertigo, ataxia and leukopenia.

Very rare: cases of pseudo-membranous colitis, liver function abnormalities, acute angioedema, hemolysis and anaphylactic shocks.

Driving & Using Machines:

Patients should be warned of the potential risk of dizziness or confusion, and advised not to drive vehicles or use machinery in the event of these disturbances.

Dosage & Administration:

Adults: 4-6 tablets of DENTYL or 2-3 tablets of DENTYL Fort daily in two or three doses during meals. for treatment in severe cases, the dose may be increased to 8 tablets of DENTYL or 4 tablets of DENTYL Fort.

Children 10-15 years: One DENTYL Tablet three times daily.

Children 6-10 years: One DENTYL Tablet twice daily.

Overdose:

Symptoms:

- Spiramycin: No case of accidental overdosage has been reported. In oral doses over 4g per day, abdominal discomfort, nausea or diarrhea may occur.
- Metronidazole: In accidental overdose poisoning, the symptoms were limited to vomiting, ataxia, and disorientation.

Treatment:

No specific treatment has been proposed. Management should be symptomatic.

Storage:

Keep out of reach of children. Store below 30 °C.

Packaging:

Each DENTYL carton box contains 20 Film Coated Tablets in two blister strips.

Each DENTYL Fort carton box contains 10 Film Coated Tablets in one blister strip.

*A medication is a product which affects your health, and it's consumption contrary to instructions is dangerous for you.
- Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medication.
- The doctor and the pharmacist experts in medicine, it's benefits and risks.
- Do not repeat the same prescription without consulting your doctor.
KEEP THE MEDICATIONS OUT OF REACH OF CHILDREN
Council of Arab Health Ministries - Union of Arab Pharmacists

Reva Pharmaceutical Industry - Syria

