

Composition:

Each FURMED 40 Tablet contains: Furosemide 40 mg.
Each FURMED 80 Tablet contains: Furosemide 80 mg.
Each FURMED 500 Tablet contains: Furosemide 500 mg.

Mechanism of Action:

Furosemide is a loop diuretic that inhibits the $\text{Na}^+/\text{K}^+/\text{2Cl}^-$ cotransporter in the ascending thick loop of Henle. So it decreases the sodium, chloride, and potassium reabsorption from the tubule. It is more effective than other diuretics.

Pharmacokinetics:

FURMED (Furosemide) is rapidly absorbed after oral administration but incompletely (60-70)% and its effect increases significantly within 4 hours. Furosemide binds to plasma proteins up to 99%. And subject to a simple metabolism. Furosemide is excreted mainly by the kidneys (80-90)%; A small portion of the ingested dose is subject to bile elimination and (10-15)% of it can appear in stool.

Indications:

- FURMED is recommended for use in all cases that require rapid and effective diuresis.
- Treatment of edema associated with congestive heart failure, cirrhosis and kidney disease including nephrotic syndrome and pulmonary edema.
- Treatment of peripheral edema due to mechanical obstruction and venous insufficiency.
- Treatment of mild to moderate hypertension when a rapid diuretic response is required, alone or in combination with other antihypertensive agents in the treatment of more severe cases.
- Furosemide 500 mg Tablets should not be given to patients with normal kidney function.

Contraindications:

- Hypersensitivity to furosemide, amiloride, sulfonamides or sulfonamide derivatives, or any of the ingredients of the product.
- Renal failure associated with anuria that does not respond to furosemide, and kidney failure due to poisoning with toxic agents for kidney or liver.
- Coma and pre-existing hepatic tumor associated with hepatic encephalopathy.
- Severe hypokalemia, severe hyponatremia, hypovolemia and dehydration (with or without hypotension accompanying blood).
- Addison's disease.

- Concomitant potassium supplements or potassium-sparing diuretics.

- Furosemide Tablets 500mg should not be given to patients with normal renal function.

Warning & Precaution:

Particularly careful monitoring is necessary in:

- Patients with hypotension, or who are at risk from a pronounced fall in blood pressure.
- Patients where latent diabetes may become manifest or the insulin requirements of diabetic patients may increase.
- Patients with gout (regular monitoring of uric acid in serum).
- Patients with hypoproteinaemia, e.g. associated with nephrotic syndrome.
- Premature infants (renal function must be monitored and renal ultrasonography performed).

Pregnancy & Lactation:

Pregnancy: Pregnancy category C. FURMED should be used during pregnancy only if the desired benefits justify the potential risk to the fetus.

Lactation: It is transmitted to the fetus through breast milk. Therefore, caution should be exercised when FURMED is administered to a nursing mother.

Driving & Using Machines:

FURMED has minor or moderate influence on the ability to drive and use machines.

Drug Interactions:

- FURMED may increase the ototoxic and nephrotoxic potential of certain antibiotics (eg. aminoglycosides) and certain cephalosporins (e.g. cephaloridine) and other ototoxic drugs.
- Sucralfate decreases the absorption of FURMED from the intestine and hence, reduces its effect.
- FURMED decreases the excretion of lithium salts and may cause increased serum lithium levels.
- It is recommended that thyroid hormones be monitored when high doses of FURMED is used.
- Probenecid: effects of furosemide may be reduced by probenecid.
- Alcohol, anaesthetic agents, alprostadil, levodopa, TCAs, MAOI, ACE inhibitors, and theophylline: enhanced hypotensive effect.
- Potassium salts (contraindicated), drospiridone, ciclosporin and tacrolimus: increased risk of hyperkalaemia.
- NSAIDs – increased risk of nephrotoxicity. Indometacin and ketorolac may antagonise the effects of furosemide
- Antihistamines, reboxetine, glycyrrizin, corticosteroids and amphoterecin: increased risk of hypokalaemia.

Side Effects:

Very common: Electrolyte disturbances, dehydration and hypovolemia, blood triglyceride increased. Hypotension and orthostatic intolerance, creatinine increased in blood.

Common: Haemoconcentration, hyponatremia, hypochloremia, hypokalemia, cholesterol increased, blood uric acid increased and gout. Hepatic encephalopathy in patients with hepatic insufficiency. Urinary volume increased.

Uncommon: Thrombocytopenia, allergic skin and mucosal reactions. Glucose tolerance decreased and hyperglycaemia, hearing disorders, usually transitory, nausea, dermatitis bullosus, erythema multiforme, pemphigoid, dermatitis exfoliative, purpura, photosensitivity.

Rare: Fever, vomiting, diarrhoea, vasculitis, paraesthesia, eosinophilia, leukopenia, anaphylactic shock.

Dosage & Administration:

Dosage in adults and children over 12 years:

Oedema: Initially 40mg daily in the morning. Maintenance dose is 20mg daily or 40mg on alternate days, increased in resistant oedema to 80mg daily.

Hypertension: 20-40mg twice daily; if 40mg twice daily does not lead to a clinically satisfactory response, the addition of other antihypertensive agents should be considered.

Elderly: Furosemide is generally eliminated more slowly. The dosage should be titrated until the required response is achieved.

Overdose:

Symptoms: Overdose may lead to hypotension, orthostatic intolerance, electrolyte imbalance or alkalosis. Excessive fluid loss may cause pronounced hypovolaemia, dehydration, circulatory collapse and haemoconcentration with increased risk for thrombosis.

Treatment: should be aimed at fluid replacement and correction of the electrolyte imbalance. Together with the prevention and treatment of serious complications resulting from such disturbances and of other effects on the body.

Storage:

Keep this medicine out of the sight and reach of children. Store below 25°C. Store in the original package.

Packaging:

FURMED (40-80-500): Each carton box contains 20 Tablets in two blisters.

آيَةُ التَّائِبِ:

الاستطابات:

مضادات الاستطباب:

- مرض أديسون.
- مكملات البوتاسيوم المصاحبة أو مدرات البول التي تحافظ على البوتاسيوم.

التحذيرات والاحتياطات:

زيادة خطر الإصابة بنقص بوتاسيوم الدم.

إنتاج شركة ريفال للصناعات الدوائية - سوريا