

**Composition:**

Each INDOMETHACIN-REVA Suppository contains:
Indomethacin 100 mg.

Pharmacological Properties:

Indomethacin belongs to non-steroidal anti-inflammatory drugs, it competitively inhibits COX-1 and COX-2 cyclooxygenase, it is a potent inhibitor of prostaglandin synthesis, with high anti-inflammatory activity and has analgesic and antipyretic properties.

Indications:

INDOMETHACIN-REVA Suppositories are indicated for the treatment of:

- Rheumatoid arthritis and moderate to severe juvenile rheumatoid arthritis.
- Ankylosing spondylitis.
- Degenerative joint disease of the hip.
- Acute gouty arthritis.
- Pains and symptoms associated with primary dysmenorrhea.
- Low back pains (lumbar pain).
- Inflammation, pain and edema after orthopedic procedures.
- Acute musculoskeletal disorders: such as bursitis, synovitis, tendinitis and tenosynovitis.

Contraindications:

INDOMETHACIN-REVA should not be used in the following cases:

- Hypersensitivity to the active substance or to any of the components of the product.
- People with a peptic ulcer or a previous history of gastro-intestinal ulceration.
- Patients with severe asthma attacks, allergic rhinitis caused by the use of non-steroidal anti-inflammatory drugs.
- Patients with inflammation or bleeding in the rectum.
- During the third trimester of pregnancy.

Warnings & Precautions:

Caution should be exercised when INDOMETHACIN-REVA is used in patients with:

- Psychiatric problems, Parkinson's disease or epilepsy.
- Heart disorders or high blood pressure.
- Impairment in renal or hepatic function.
- Elderly patients.

Pregnancy & Lactation:

Pregnancy: Indomethacin should not be used during the first and second trimesters of pregnancy unless the potential benefits outweigh the risks. Indomethacin

is contraindicated during the last trimester of pregnancy.

Lactation: Indomethacin is excreted in breast milk. Administration of indomethacin is not recommended in breast-feeding mothers.

Drug Interactions:

- **Diuretics:** Concomitant use with indomethacin may lead to increased plasma levels of indomethacin and decreased renal clearance.
- **Aspirin:** The concomitant use of aspirin and other salicylates with indomethacin should be avoided due to its significant gastro-intestinal effects.
- **Probenecid and methotrexate:** Concomitant use may increase plasma levels of indomethacin.
- **Antihypertensives and diuretics:** The concomitant use of indomethacin may reduce the effect of antihypertensive drugs (beta-blockers, ACEs, ARBs, alpha-blockers) and diuretics.
- **Corticosteroids:** The risk of gastrointestinal bleeding and ulcers with NSAIDs may be increased when used concomitantly with corticosteroids.
- **Vancomycin:** Concomitant administration with indomethacin should be used with caution due to the possibility of nephrotoxic effects.
- **Digoxin:** Increased plasma concentrations and a prolongation of the half-life of digoxin have been reported with co-administration of indomethacin. Therefore, serum digoxin levels should be monitored when they are used concomitantly.

Driving & Using Machines:

Patients may experience dizziness, drowsiness, visual disturbances, or headaches, so caution should be exercised when driving or operating machinery.

Side Effects:

Headache, dizziness, depression, nausea, vomiting, abdominal pain, single or multiple ulcers, increased blood pressure, fluid retention, edema, skin rash, urticaria, itching and irritation of the rectal mucosa, diarrhea or constipation.

Dosage & Administration:

In Adults: One suppository once or twice a day.
The daily dose should not exceed 200 mg.

Storage:

Store below 25 °C.

Packaging:

Each INDOMETHACIN-REVA Carton Box contains 10 Suppositories in two PVC/PE strips.

*A medication is a product which affects your health, and its consumption contrary to instructions is dangerous for you.
*Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medication.
*Do not expect the same prescription without consulting your doctor.
*KEEP THE MEDICATIONS OUT OF REACH OF CHILDREN
Council of Arab Health Ministries - Union of Arab Pharmacists

