

IVA

Syrup

**Composition:**

Each 5 ml IVA Syrup contains:

Dried Ivy Leaf Extract 35 mg.

Pharmacological Properties & Mechanism of Action:

The Anti-bronchospasm, bronchodilator, expectorant activity, of the extract and/or the isolated compounds such as α -Hederin has been demonstrated in several in vitro and in vivo studies. The mechanism of the mucolytic effect observed in experimental clinical studies has not yet been determined. Also, anti-inflammatory effects have been shown in various in vivo models using alcoholic extracts of ivy leaves taken orally.

Pharmacokinetics:

There are no data available on the pharmacokinetics of ivy leaf extracts. There are no pharmacokinetic studies of ivy Leaf and extracts prepared from them in humans.

In animal studies, following oral administration of ivy leaf extract, Hederacoside C was found in plasma 5 minutes after an oral dose. Peak plasma concentrations are reached after approximately 15 - 60 minutes. This indicates that Hederacoside C is rapidly absorbed from the gastrointestinal tract of mice and is mainly excreted via urine.

Indications:

IVA is a herbal medicinal product used to improve the symptoms of chronic bronchitis and acute respiratory infections with cough, through its mucolytic, bronchodilating and cough sedative effects.

Contraindications:

Hypersensitivity to the active substance or to any of the components of the product.

Warnings & Precautions:

- Persistent or recurrent coughing in children aged 2-4 years requires a medical diagnosis prior to treatment.
- When shortness of breath, fever, or pu-

lulent sputum occurs, you should consult a doctor.

- Caution is advised when using IVA in patients with gastritis or stomach ulcers.

Pregnancy & Lactation:

Generally consult a doctor or pharmacist before taking any drug during pregnancy or lactation.

Drug Interactions:

No drug interactions have been known far.

Side Effects:

Side effects resulting from the use of ivy leaf extract are rare. Gastrointestinal symptoms such as nausea, vomiting and diarrhea may appear.

Driving & Using Machines:

IVA syrup has no effect on the ability to drive or use machines.

Dosage & Administration:

Unless otherwise prescribed by a doctor, the usual dose is:

- Children under two years of age: according to the doctor's advice.
- Children between 2-5 years of age: 2.5 ml (half a teaspoonful) two times daily.
- Children aged 6-12 years: 5 ml (teaspoonful) two times daily.
- Children over 12 years old and adults: 5 ml (teaspoonful) three times daily.
- * If the symptoms persist for more than a week after using the medicine, a doctor or a pharmacist should be consulted.

Overdose:

In previous single and repeated dose toxicity studies using dried extract of ivy leaf, the results did not raise any safety concerns.

Storage:

Keep out of reach of children. Store below 30 °C.

Packaging:

Each IVA syrup carton box contains amber glass bottle of 100 ml.

- A medication is a product which affects your health, and its consumption contrary to instructions is dangerous for you.
- Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medication.
- The doctor and the pharmacist experts in medicine, it's benefits and risks
- Do not repeat the same prescription without consulting your doctor.
KEEP THE MEDICAMENTS OUT OF REACH OF CHILDREN
Council of Arab Health Ministers, Union of Arab Pharmacists

Reva Pharmaceutical Industry - Syria

نتاج شركة ريفال للصناعات الدوائية - سوريا