

INDOMETHACIN-B Capsules



Composition:

Each INDOMETHACIN-B Capsule contains:

Indomethacin 25 mg.

Thiamine hydrochloride (vitamin B1) 50 mg.

Mechanism of Action:

Indomethacin is a non-steroidal anti-inflammatory, and it exhibits antipyretic and analgesic properties. Indomethacin is a potent inhibitor of prostaglandin synthesis, which is produced through inhibition of the cyclooxygenase (COX1, COX2). Thiamine (Vitamin B1) works to inhibit the transmission of nerve impulses and thus has an analgesic effect on neuropathic pains.

Pharmacokinetics:

Indomethacin is rapidly and almost completely absorbed from the gastrointestinal tract following oral ingestion. Peak plasma levels occur in 0.5 - 2 hours in fasting subjects, longer if taken with or after food. More than 90% is bound to plasma proteins. It is distributed in the synovial fluid and crosses the blood-brain barrier and the placenta. It is metabolised in the liver and converted to inactive metabolites. Half-life in plasma is variable from 2-11 hours. It is mainly excreted in the urine, lesser amounts in the feces.

Thiamine is well absorbed from the gastrointestinal tract following oral administration. It is widely distributed to most body tissues. Thiamine is not stored to any appreciable extent in the body; amounts in excess of the body's requirements are excreted in the urine as unchanged thiamine or metabolites.

Indications:

INDOMETHACIN-B is indicated in the following cases:

- Moderate to severe rheumatoid arthritis including acute flares of chronic disease.
- Moderate to severe ankylosing spondylitis.
- Moderate to severe osteoarthritis.
- Acute painful shoulder (bursitis and/or tendinitis).
- Acute gouty arthritis.

Contraindications:

INDOMETHACIN-B is not indicated in patients who:

- Are hypersensitive to the active material or the excipients.
- Have experienced asthma, urticaria or allergic reactions after taking other NSAIDs.
- In the setting of coronary artery bypass graft (CABG) surgery.
- Have an active or pre-bleeding peptic ulcer or recurrent peptic ulcer.

Warnings & Precautions:

INDOMETHACIN-B should use with caution in people with:

- History of high blood pressure, congestive heart failure, peripheral arterial disease or renal disease.
- Mental health problems e.g. mood changes or epilepsy or parkinson's diseases.
- Severe infection.
- Previous complaint related to gastrointestinal diseases including ulcerative colitis, Crohn's disease, ulceration, bleeding and perforation.
- Asthma.

Indomethacin may be associated with a small increased risk of heart attack (myocardial infarction) or stroke.

Thiamine could give false positive results for urobilinogen determination by the Ehrlich's reaction.

Pregnancy & Lactation:

Pregnancy: Pregnancy category C, indomethacin should be avoided during pregnancy because it may cause premature closure of the ductus arteriosus.

Lactation: Indomethacin is excreted in the milk of lactating mothers. Indomethacin is not recommended for use during lactation.

Drug Interactions:

Antihypertensive drugs (e.g., Angiotensin II receptors antagonists and Beta-blockers): NSAIDs including indomethacin may diminish the antihypertensive effect of these drugs.

Aspirin: The use of indomethacin in conjunction with aspirin or other salicylates is not recommended, the incidence of gastrointestinal side effects was significantly increased with combined therapy.

Digoxin: Indomethacin increases the serum concentration and prolong the half-life of digoxin.

NSAIDs: The concomitant use of indomethacin with other NSAIDs is not recommended

Oral Anticoagulants: The patients should be observed for alterations of the prothrombin time.

Probenecid: When indomethacin is given to patients receiving probenecid, the plasma levels of indomethacin are likely to be increased.

Diuretics: NSAIDs can reduce natriuretic effect of furosemide and thiazide diuretics.

Side Effects:

Most common adverse reactions are headache, dizziness, dyspepsia (heartburn and epigastric pain), nausea, diarrhea, abdominal pain, constipation, vertigo, somnolence, depression and fatigue including malaise and listlessness, tinnitus.

Driving & Using Machines:

INDOMETHACIN-B generally has no adverse effects on the ability to drive and use machinery.

Dosage & Administration:

Adults:

- The dosage for rheumatoid arthritis, ankylosing spondylitis and osteoarthritis is one capsule 2 or 3 times a day.
- The dosage for acute painful shoulder (bursitis and/or tendinitis) is one capsule 3 or 4 times a day.
- The dosage for acute gouty arthritis is two capsules three times a day

Children:

Indomethacin is contraindicated in children under 14 years old as its safety has not been established.

Overdose:

Indomethacin:

Symptoms: Similar to side effects, sometimes more severe. Gastrointestinal bleeding may also occur. In cases of serious poisoning, acute renal failure and liver damage may occur. Treatment: Treatment is symptomatic and as needed. Gastric emptying or oral administration of activated charcoal within 1 hour of administration of the drug is indicated. If indomethacin has already been absorbed, alkalis should be given to enhance urinary excretion of indomethacin.

Thiamine:

No overdose has been reported.

Storage:

Keep out of reach of children. Store in a dry place at a temperature below 25 °C.

Packaging:

Each INDOMETHACIN-B carton box contains 30 Capsules in three blister strips.

*A medication is a product which affects your health, and it's consumption contrary to instructions is dangerous for you.
- Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medication.
- The doctor and the pharmacist experts in medicine, it's benefits and risks.
- Do not repeat the same prescription without consulting your doctor.
- KEEP THE MEDICAMENTS OUT OF REACH OF CHILDREN
Council of Arab Health Ministries, Union of Arab Pharmacists

