

Levox

Film Coated Tablets



Composition:

Each Levox (500 - 750) Film Coated Tablet contains:

Levofloxacin (as hemihydrate) 500 mg or 750 mg.

Mechanism of Action:

Levofloxacin is a fluoroquinolone antibiotic. Its bactericidal effect is produced by inhibiting the second and fourth types of the enzyme topoisomerase (DNA-gyrase), which is necessary for the copy, repair and replication of bacterial DNA.

Pharmacokinetics:

Levofloxacin is rapidly absorbed after oral administration, reaching peak blood concentrations after 1-2 hours. Absolute bioavailability is approximately (99-100)%. The plasma proteins binding of levofloxacin is about 40%. It is widely distributed in various tissues of the body, and is excreted largely unchanged through the urine (about 85%) and, to a lesser extent, through the feces. The elimination half-life is approximately 6-8 hours.

Indications:

Levox is used to treat different types of bacterial infections: Pneumonia, community-acquired pneumonia, acute bacterial sinusitis, acute bacterial exacerbation of chronic bronchitis, inhalation anthrax, skin and skin structure infections, chronic bacterial prostatitis, complicated urinary tract infections, acute pyelonephritis, urinary tract infections not complicated.

Contraindications:

- Patients with hypersensitivity to quinolones or any of the components of the product.
- Patients with a history of tendon disorders related to the administration of fluoroquinolones.
- Epilepsy patients.
- Children and adolescents.

Warnings & Precautions:

- Caution should be taken when using Levox in patients with known risk factors for prolongation of the QT interval .
- As with all quinolones, disturbances in blood glucose, including both hypoglycaemia and hyperglycaemia have been reported.
- It should be used with caution in patients with renal or hepatic disorders.
- Levox should be used with caution in patients with myasthenia gravis.
- Levox should be used with caution in patients with glucose-6-phosphate dehydrogenase deficiency due to the risk of hemolysis.
- Levox should be used with extreme caution in patients who are prone to seizures, or are being treated with drugs that lower the seizure threshold.
- Levox use has rarely been associated with Hypersensitivity reactions.

- Tendonitis and/or rupture of tendons are observed during treatment with quinolones. After the first signs of pain or inflammation, the treatment should be discontinued.

- You should consult a doctor before continuing treatment with Levox in the event that symptoms of peripheral neuropathy appear, such as pain, tingling or numbness, to prevent progression of the condition.

Pregnancy & Lactation:

Levox is contraindicated in both pregnancy and lactation.

Driving & using machines:

Some side effects (Such as dizziness and visual disturbances) may impair the patient's ability to drive and using machines.

Drug Interactions:

- Levofloxacin absorption is significantly reduced when iron salts, or magnesium or aluminium-containing antacids, or didanosine, or sucralfate are administered concomitantly with Levox tablets.
- The renal clearance of levofloxacin was reduced by cimetidine (24%) and probenecid (34%).
- The half-life of ciclosporin was increased by 33% when coadministered with levofloxacin.
- Concomitant administration of Levox with vitamin K antagonists may increase its anticoagulant effect.

Side Effects:

Common: Insomnia, headache, dizziness, diarrhoea, vomiting, nausea and hepatic enzymes increased (e.g. ALT/AST).

Uncommon: Fungal infection including candida infection, pathogen resistance, leukopenia, eosinophilia, anorexia, anxiety, confusion, nervousness, somnolence, tremor, vertigo, dyspnoea, dyspepsia, constipation, skin rash and urticaria.

Dosage & Administration:

Levox tablets are administered once or twice daily. The dosage depends on the type and severity of the infection and the sensitivity of the presumed causative pathogen. The duration of therapy varies according to the course of the disease (see table below).

Dosage in patients with normal renal function:

Type of Infection	Daily dose regimen	Duration of treatment
Community-acquired pneumonia	500 mg once daily	7 - 14 days
	or 750 mg once daily	5 days
Nosocomial Pneumonia	750 mg once daily	7 - 14 days
Acute bacterial sinusitis	500 mg once daily	10 - 14 days
	or 750 mg once daily	5 days
Acute Bacterial Exacerbation of Chronic Bronchitis	500 mg once daily	≥ 7 days
	500 mg once daily	60 days
Inhalation Anthrax	500 mg once daily	60 days
	500 mg once daily	60 days
Skin/Skin Structure Infections	Uncomplicated: 500 mg once daily	7-10 days
	Complicated: 750 mg once daily	7-14 days
Chronic bacterial prostatitis	500 mg once daily	28 days

Complicated Urinary Tract Infections & Acute Pyelonephritis	250 mg once daily	10 days
	or 750 mg once daily	5 days
Uncomplicated Urinary Tract Infections	250 mg once daily	3 days

Dosage in Adult Patients with Renal Impairment (Creatinine Clearance less than 50 ml/minute):

Creatinine Clearance greater than or equal to 50 ml/minute	Creatinine Clearance 20 to 49 ml/minute	Creatinine Clearance 10 to 19 ml/minute	Haemodialysis or Chronic Ambulatory Peritoneal Dialysis (CAPD)
750 mg every 24 hours	750 mg every 48 hours	Initial dose 750 mg, then 500 mg every 48 hours	Initial dose 750 mg, then 500 mg every 48 hours
500 mg every 24 hours	Initial dose 500 mg, then 250 mg every 24 hours	Initial dose 500 mg, then 250 mg every 48 hours	Initial dose 500 mg, then 250 mg every 48 hours
250 mg every 24 hours	No dosage adjustment required	250 mg every 48 hours. If treating uncomplicated UTI, then no dosage adjustment is required	No information on dosing adjustment is available

Overdose:

Symptoms: central nervous system symptoms such as confusion, dizziness, impairment of consciousness, convulsive seizures, increases in QT interval as well as gastro-intestinal reactions such as nausea and mucosal erosions.

Treatment: Patients should be referred to hospital urgently for immediate medical attention.

Storage: Keep out of reach of children. Store below 25°C.

Packaging: Each Levox (500-750) carton box contains: 10 Film Coated Tablets in a blister strip.

*A medication is a product which affects your health, and it's consumption contrary to instructions is dangerous for you.
 - Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medication.
 -The doctor and the pharmacist experts in medicine, its benefits and risks.
 -Do not repeat the same prescription without consulting your doctor.
 KEEP THE MEDICATION OUT OF REACH OF CHILDREN.
 Council of Arab Health Ministers, Union of Arab Pharmacists
 Revva Pharmaceutical Industry - Syria

نوع الانتان	نظام الجرعة اليومية	مدة العلاج
الانتان الرئوي المكتسب من المجتمع	٥٠٠ ملغ مرة واحدة يومياً	٧- ١٤ يوماً
الانتان الرئوي	أو ٧٥٠ ملغ مرة واحدة يومياً	٥ أيام
	٧٥٠ ملغ مرة واحدة يومياً	٧- ١٤ يوماً
انتان الجيوب الأنفية الجرثومي الحاد	٥٠٠ ملغ مرة واحدة يومياً	١٠- ١٤ يوماً
	أو ٧٥٠ ملغ مرة واحدة يومياً	٥ أيام
التفاقم الجرثومي الحاد لالتهاب الشعب الهوائية المزمن	٥٠٠ ملغ مرة واحدة يومياً	≤ ٧ أيام
استنشاق الجمرة الخبيثة	٥٠٠ ملغ مرة واحدة يومياً	٦٠ يوماً
انتانات الجلد وبنية الجلد	غير معقد: ٥٠٠ ملغ مرة واحدة يومياً	٧- ١٠ يوماً
	معقد: ٧٥٠ ملغ مرة واحدة يومياً	٧- ١٤ يوماً
التهاب البروستات الجرثومي المزمن	٥٠٠ ملغ مرة واحدة يومياً	٢٨ يوماً
انتانات المسالك البولية المعقدة & التهاب الحويضة والكلية الحاد	٢٥٠ ملغ مرة واحدة يومياً	١٠ أيام
	أو ٧٥٠ ملغ مرة واحدة يومياً	٥ أيام

- يجب استشارة الطبيب قبل مواصلة العلاج بليفوكس في حال ظهور أعراض الإعتلال العصبي المحيطي مثل الألم أو الوخز أو التنميل لمنع تطور الحالة.

التحذيرات والاحتياطات:

الدواء مستحضر يوزن على مسبك واستعماله خلافاً لتعليمات بمرحلتي التحضير
وبناعة وصفية للطبيب بطريقة الاستعمال المتخصص عن غلبته وتعليمات الصيدلي
التي صرنا لك ، فالطبيب هو الصيدلي هما الخبيران في الدواء ويقنعوا بضرره
لاكثر صرف الدواء بدون وصفة طبية.
لا تترك الأدوية بمقتول ايدي الأطفال
مجلس وزراء الصحة العرب واتحاد الصيدلة العرب