

Losartan Reva

Film Coated Tablets



Composition: Each Losartan Reva Film Coated Tablet contains: Losartan Potassium (25-50-100) mg.

Mechanism of Action:

Losartan Reva belongs to a group of medicines known as angiotensin II receptors antagonists, which help control high blood pressure. Angiotensin II binds to the AT1 receptor and causes blood vessels to constrict, which leads to high blood pressure. Losartan selectively blocks the AT1 receptor. As a result, blood vessels relax and blood pressure decreases.

Pharmacokinetics:

Following oral administration, losartan is well absorbed and undergoes first-pass hepatic metabolism, forming an active carboxylic acid metabolite and other inactive metabolites. The systemic bioavailability of losartan is approximately 33%. Mean peak plasma concentrations of losartan and its active metabolite are reached in 1 hour and in 3-4 hours, respectively. Both losartan and its active metabolite are $\geq 99\%$ bound to plasma proteins, primarily albumin. Both biliary and urinary excretion contribute to the elimination of losartan and its metabolites.

Indications: Losartan Reva is indicated in the following cases:

- Essential hypertension in adults and children greater than 6 years old.
- Diabetic nephropathy with an elevated serum creatinine and proteinuria in patients with type II diabetes and a history of hypertension.
- Reduction in the risks of stroke in adults hypertensive patients with left ventricular hypertrophy.

Contraindications:

Losartan Reva is contraindicated in the following cases:

- Allergy to Losartan or any of the other ingredients in the product.
- In patients with severe hepatic impairment.
- The concomitant use of Losartan Reva with aliskiren-containing products is contraindicated in patients with diabetes mellitus or renal impairment.

Warnings & Precautions:

- Symptomatic hypotension may occur in some patients after the first dose or after an increase in the dose. In this case hypovolemia or electrolyte depletion must be corrected before Losartan Reva is administered.
- Losartan Reva should use with caution in patients with bilateral renal artery stenosis or stenosis of the artery to a solitary kidney.
- Patients with a history of angioedema should be closely monitored.
- In patients with heart failure, with or without renal insufficiency, there is a risk of developing severe arterial hypotension and (often severe) renal insufficiency.
- Losartan Reva is not recommended in patients with primary hyperaldosteronism.
- The dose should be reduced in patients with hepatic insufficiency.

Pregnancy & Lactation:

Pregnancy: Pregnancy Category D, Losartan Reva is contraindicated in pregnancy. When pregnancy is diagnosed, treatment with losartan should be stopped immediately.

Lactation: Losartan is not recommended and alternative treatments with better established safety profiles during breast-feeding are preferable, especially while nursing a newborn or premature.

Drug Interactions:

- Other antihypertensive agents: may increase the hypotensive action of Losartan.
- Fluconazole: reduces the systemic concentration of the active metabolite losartan by approximately 50%.
- Rifampicin: Concomitant administration with losartan results in a 40% decrease in the plasma concentration of the active metabolite.
- Caution and frequent monitoring of serum potassium levels should be exercised when concomitantly used with potassium supplements, potassium-sparing diuretics or other drugs that may increase potassium levels (such as heparin).
- NSAIDs: Simultaneous use leads to increased risk of renal impairment and reduced diuretic, natriuretic, and antihypertensive

effects.

- Co-administration of lithium and losartan should be undertaken with caution. If this combination proves essential, serum lithium level monitoring is recommended during concomitant use.

Side Effects:

- Common side effects: dizziness, vertigo, hyperkalemia, fatigue, hypotension. Anemia and increase in blood urea were common in patient with chronic heart failure.
- Uncommon side effects: somnolence, headache, sleep disorders, palpitations, abdominal pain, constipation, fatigue, dyspnoea, cough, pruritus, skin rash and oedema.

Driving & using machines:

When driving vehicles or operating machinery, it must be borne in mind that dizziness or drowsiness may occasionally occur when taking antihypertensive therapy, in particular during initiation of treatment or when the dose is increased.

Dosage & Administration:

Patients with type II diabetes mellitus with Hypertensive with proteinuria > 0.5 g/day:

The usual starting dose and maintenance dose is 50 mg once daily for most patients. The maximum antihypertensive effect is attained 3-6 weeks after initiation of therapy. Some patients may receive an additional benefit by increasing the dose to 100 mg once daily (taken in the morning). Losartan Reva Tablets may be administered with other antihypertensive drugs.

Heart failure: The starting dose is 12.5 mg once daily. Dosage should be adjusted at weekly intervals (12.5 mg daily, 25 mg daily and 50 mg daily) according to patient tolerance.

Reduction in the risks of stroke in adults hypertensive patients with left ventricular hypertrophy:

The usual starting dose is 50 mg of Losartan Reva once daily. A low dose of hydrochlorothiazide should be added and/ or the dose of Losartan Reva should be increased to 100 mg once daily based on the observed response blood pressure.

Patients with intravascular volume depletion (e.g. those treated with high-dose diuretics): Starting dose 25 mg once daily should.

Patient with hepatic impairment: Low doses should be started in patients with a history of hepatic impairment. Losartan is also contraindicated in patients with severe hepatic impairment.

Children from 6 to 18 years old: For patients weighing 20 to 50 kg who can take the tablets, the recommended dose is 25 mg once daily. In exceptional cases the dose can be increased to a maximum of 50 mg once daily. Dosage should be adjusted according to blood pressure response.

In patients > 50 kg, the usual dose is 50 mg once daily. In exceptional cases the dose can be adjusted to a maximum of 100 mg once daily.

Losartan Reva is not recommended for use in children under 6 years old and in children with glomerular filtration rate < 30 ml/min / 1.73 m².

Use in Elderly: Treatment may be started with 25 mg daily in patients over 75 years of age, dosage adjustment is not usually necessary for elderly.

Overdose:

Symptoms: The most common symptoms of an overdose are hypotension and tachycardia. Bradycardia could occur as a result of stimulation of the vagus nerve.

Treatment: If symptomatic hypotension should occur, supportive treatment should be instituted.

Stabilisation of the circulatory system should be given priority. If the drug is taken orally, sufficient dose of activated charcoal should be indicated. Afterwards, close monitoring of the vital parameters should be performed and it should be corrected if necessary. Neither Losartan nor the active metabolite can be removed by haemodialysis.

Storage: Store at a temperature below 30°C. Keep out of reach of children.

Packaging:

Each Losartan Reva (25-50-100) carton box contains 20 or 30

Film Coated Tablets in two or three blister strips.

- A medication is a product which affects your health, and it's consumption contrary to instructions is dangerous for you.
- Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sells the medication.
- The doctor and the pharmacist experts in medicine, it's benefits and risks.
- Do not repeat the same prescription without consulting your doctor.
- KEEP THE MEDICATIONS OUT OF REACH OF CHILDREN
Council of Arab Health Ministers, Union of Arab Pharmacies

