

New Pect D.C Syrup



Composition:

Each 5 ml of New Pect D.C Syrup contains:

Dextromethorphan (HBr)	10 mg.
Oxomemazine (HCl)	1.5 mg.
Sodium Benzoate	60 mg.

Pharmacological Properties:

New Pect D.C contains an effective combination in the treatment of dry and allergic cough. Dextromethorphan is a cough inhibitor which has a central action on the cough center in the medulla.

Oxomemazine is an antihistamine from the family of phenothiazines. It has antitussive and sedative properties.

Sodium benzoate dissolves phlegm and increases the fluidity of bronchial secretions.

Pharmacokinetics:

Pharmacokinetic data with oxomemazine are lacking. Phenothiazines group, including oxomemazine, can cross the blood-brain barrier, leading to sedative effects. A small percentage of active substance found unchanged in the urine. Dextromethorphan is well absorbed from the gastrointestinal tract. It undergoes rapid and extensive first-pass metabolism in the liver after oral administration. Dextromethorphan, which also has antitussive action, is the main metabolite. In some individuals metabolism proceeds more slowly and unchanged dextromethorphan predominates in the blood and urine.

Sodium benzoate is absorbed from the gastrointestinal tract and is combined with glycine in the liver, forming hyperuric acid, which is rapidly excreted urine route.

Indications:

New Pect D.C is indicated in the treatment of dry and allergic cough.

Contraindications:

- Hypersensitivity to any of the components of the product.
- Concurrent use with MAO inhibitors or within 14 days of stopping MAO inhibitors therapy.
- Severe hepatic impairment.
- Infants (less than 2 years of age).

Warnings & Precautions:

- Caution should be exercised in patients with asthma and New Pect D.C should not be taken during an acute attack.
- New Pect D.C should not be used in patients with respiratory failure.
- Prolonged use of New Pect D.C may cause habituation, even at therapeutic doses.
- Caution should be exercised in patients with narrow-angle glaucoma, urinary retention, prostatic hypertrophy, hepatic insufficiency and epilepsy.
- You should see a doctor if the cough persists for more than several days.

Pregnancy & Lactation:

Pregnancy: The effect of this medication during pregnancy is not well known, this product should not be used during pregnancy without medical advice.

Lactation: No data are currently available on the excretion of this drug in breast milk. As a precaution, it is not recommended for use during breastfeeding.

Drug Interactions:

Avoid concurrent use of New Pect D.C with the following medicines:

- Alcohol: Avoid use with alcohol, due to the increased risks of hepatotoxicity and hypnotic effect.
- Tricyclic antidepressants, MAO inhibitors, Anti-Parkinson's disease drugs, disopyramide, neuroleptic phenothiazines: Additional atropine (anti muscarinic) side effects and serotonergic effects, including the development of a serotonin syndrome.
- Antidepressants with sedative effects, barbiturates and benzodiazepines: Increase their inhibitory effects.

Side Effects:

Drowsiness, dizziness, nausea, vomiting. Anti-cholinergic effects such as drying out of mucous, constipation, mydriasis, cardiac palpitations, urinary retention. Very rare cases of serious skin reactions have been reported.

Driving & Using Machines:

This medicine can impair cognitive function and can affect a patient's ability to drive and use machines safely.

Dosage & Administration:

New Pect D.C is well tolerated in adults and children.

Children 2-6 years: 2.5-5 ml, two or three times daily.

Children 6-12 years: 2.5-5 ml, three or four times daily.

Adults: 5-10 ml, three or four times daily.

Overdose:

Symptoms:

Signs of oxomemazine overdose are convulsion (especially for infants and children), disorientation, and coma. Dextromethorphan overdose may be associated with nausea, vomiting, agitation, confusion, somnolence, nystagmus, cardiotoxicity (tachycardia, abnormal ECG including QTc prolongation), ataxia, psychosis with visual hallucinations, hyperexcitability. In the event of massive overdose, the following symptoms may be observed: coma, respiratory depression, convulsions.

Treatment:

- Symptomatic treatment is applied.

- Activated charcoal can be administered to asymptomatic patients who have ingested overdoses within the preceding hours.

- For patients who have ingested dextromethorphan and are sedated or comatose, naloxone in the usual doses for treatment of opioid overdose, can be considered. Benzodiazepines can be used to treat convulsions and lower the temperature by cooling the body from the outside to lower the temperature associated with serotonin syndrome.

Storage:

Keep out of reach of children. Store below 25 °C.

Packaging:

Each New Pect D.C carton box contains amber glass bottle of 100 ml.

-A medication is a product which affects your health, and it's consumption contrary to instructions is dangerous for you.
- Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medication.
- Do not repeat the same prescription without consulting your doctor.
- The doctor and the pharmacist experts in medicine, it's benefits and risks.
- KEEP THE MEDICATIONS OUT OF REACH OF CHILDREN
Council of Arab Health Ministries - Union of Arab Pharmacists

Reva Pharmaceutical Industry - Syria

التركيب:

كل ٥ مل نيو بكت ديسي شراب تحتوي على:

- ديکستروميثورفان (هيدروبروميد) ۱۰ ملغ.
 اوكسو ميمازين (هيدروکلوريد) ۱,۵ ملغ.
 بنزوات الصوديوم ۶۰ ملغ.

يحتوي نيو بكت د.سي على تركيبة فعالة في معالجة السعال الجاف والتحسسي.

أوكسوميمازين هو مضاد هستامين من عائلة الفينوثيازين. له خصائص مضادة للسعال ومهدئة. يذيب بنزوات الصوديوم البلغم ويزيد من سيولة إفرازات الشعب الهوائية.

إن المعلومات حول الحرائك الدوائية للأوكسيميازين قليلة. يمكن لمجموعة الفينوثيازين، بما في ذلك الأوكسيميازين، عبور الحاجز الدماعي الشوكي مما يعطي آثاراً مهدئة. كما يلاحظ العبور إلى نسبة صغيرة من المادة الفعالة دون تغيير في البلوغ. يقص ديكستروميثورفان بشكل جيد من القناة الهضمية. يخضع للاستقلاب عبر المرور الكبدي الأول بشكل سريع وواسع النطاق بعد تناوله عن طريق الفم. ديكستروميثورفان وهو المستقلب الرئيسي ويمتلك أيضاً تأثيراً مضاداً للسعال. لدى بعض الأفراد يكون الاستقلاب أبطأ ويتواجد ديكستروميثورفان دون تغيير في الدم والبول.

يتم امتصاص بنزوات الصوديوم من القناة الهضمية ويتم ربطه بالغليسين في الكبد، مكوناً حمض الهيپوريك، والذي يتم إطرأه بسرعة عن طريق البول.

يُستطب نيو بكت د.سي في علاج السعال الجاف والتحسسي.

[illegible]

يحفظ بعيداً عن متناول أيدي الأطفال. يحفظ في درجة حرارة أقل من ٢٥°م.

كل عبوة كرتونية نيو بكت. د. سي. تحتوي على عبوة زجاجية عاتمة
سعة ١٠٠ مل.

الحمل والارضاع:

الحمل: إن تأثير هذا الدواء أثناء الحمل غير معروف بشكل جيد، يجب عدم استخدام هذا الدواء أثناء الحمل دون استشارة طبية.

الإرضاع: لا يوجد بيانات متوفرة حالياً حول إفراز هذا الدواء في حليب الثدي. كإجراء احترازي، لا يُنصح باستعماله أثناء الرضاعة الطبيعية.

الدواء مستحضر بوتر علي صحتك واستعماله خلافاً للتعليمات يهتكك للخطر
 يقع بصفة الطبيب وطريقة الاستعمال المنصوص عليها والتعليمات الصيدلي
 الذي صرفها لك ، فاطمئني والصديقي هما الخياران في الدواء وينتفع وضرره
 لاكثر صرف الدواء بدون وصفة طبية.
 لاترك الأدوية بمقتول أيدي الأطفال.
 مجلة ١١٠٤ الصحة العامة، العدد الثالث المجلد الثاني