



Composition:

Each Olen (2.5, 5, 7.5, 10, 15) Film Coated Tablet contains:
Olanzapine: 2.5, 5, 7.5, 10, or 15 mg.

Pharmacological Properties & Mechanism of Action:

Olen efficacy in schizophrenia is mediated through a combination of dopamine and serotonin type 2 (5HT₂) antagonism. The mechanism of action of olanzapine in the treatment of acute manic or mixed episodes associated with bipolar I disorder is unknown.

Pharmacokinetic:

Olanzapine is well absorbed after oral administration, reaching peak plasma concentrations within 5 to 8 hours. Olanzapine binds to plasma proteins (mainly albumin and α 1-acid-glycoprotein) by about 93%. It is metabolized in the liver by conjugative and oxidative pathways. Following a single oral dose of olanzapine, 7% of the dose of olanzapine was recovered in the urine as unchanged drug. Approximately 57% and 30% of the dose was recovered in the urine and feces, respectively.

Indications:

- Treatment of schizophrenia in adults.
- Treatment of manic or mixed episodes associated with bipolar I disorder and maintenance treatment of bipolar I disorder.
- Adjunct to valproate or lithium in the treatment of manic or mixed episodes associated with bipolar I disorder.
- Depressive episodes associated with bipolar I disorder.

Contraindications:

- Hypersensitivity to olanzapine or to any of the excipients.
- Patients with known risk of narrow-angle glaucoma.

Warnings & Precautions:

- Olen is not recommended for use in patients with dementia-related psychosis and/or behavioral disturbances because of an increase in the risk of cerebrovascular accident.
- The level of glucose and lipids in the blood should be monitored regularly.
- Acute symptoms such as sweating, insomnia, tremor, anxiety, nausea, or vomiting have been reported rarely when olanzapine is stopped abruptly.
- If signs or symptoms of tardive dyskinesia appear in a patient on olanzapine, a dose reduction or discontinuation should be considered.
- Postural hypotension was infrequently observed in the elderly over 65 years of age.
- Caution should be exercised in patients:
 - With prostatic hypertrophy, or paralytic ileus.
 - With elevated ALT and/or AST, or with signs and symptoms of

hepatic impairment.

- With low leukocyte and/or neutrophil counts for any reason.
- Who have a history of nervous seizures.
- Worsening of Parkinson's disease symptoms and hallucinations when using olanzapine in the treatment of psychosis associated with dopamine agonists.

Pregnancy & Lactation:

Pregnancy: Pregnancy category C. Olen should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Lactation: Olanzapine is excreted in breast milk. Mothers should be advised not to breast-feed an infant if they are taking olanzapine.

Driving & Using Machines:

Because olanzapine may cause somnolence and dizziness, patients should be cautioned about operating machinery, including motor vehicles.

Drug Interactions:

- Diazepam and alcohol: May potentiate orthostatic hypotension.
- Carbamazepine and smoking: The metabolism of olanzapine is induced by smoking and carbamazepine, which may lead to reduced olanzapine concentrations in the blood.
- Fluvoxamine: May increase olanzapine blood levels.
- Activated charcoal reduces the bioavailability of oral olanzapine by 50% to 60%.
- Antihypertensive Agents: Enhanced antihypertensive effect.
- Levodopa and Dopamine Agonists: May antagonize levodopa/dopamine agonists.

Side Effects:

The most frequently reported adverse reactions were somnolence, weight gain, eosinophilia, elevated prolactin, cholesterol, glucose and triglyceride levels, glucosuria, increased appetite, dizziness, akathisia, parkinsonism, leukopenia, neutropenia, dyskinesia, orthostatic hypotension, anticholinergic effects (constipation, dry mouth), transient asymptomatic elevations of hepatic aminotransferases, rash, fatigue, pyrexia, arthralgia, increased alkaline phosphatase, high uric acid, high creatine phosphokinase and oedema, erectile dysfunction in males, decreased libido in both males and females.

Dosage & Administration:

Olen may be given without regard to meals.

Indication	Adults	Adolescents (13-17 Years)
Schizophrenia	Start at 5-10 mg once daily. Target: 10 mg/day within several days.	Start at 2.5-5 mg once daily. Target: 10 mg/day within several days.

Bipolar I Disorder (manic or mixed episodes)	Start at 10 or 15 mg once daily.	Start at 2.5-5 mg once daily. Target: 10 mg/day within several days.
Depressive Episodes associated with Bipolar I Disorder	In combination with fluoxetine: Start at 5 mg of oral olanzapine and 20 mg of fluoxetine once daily.	In combination with fluoxetine: Start at 2.5 mg of oral olanzapine and 20 mg of fluoxetine once daily.

Renal and/or Hepatic Impairment: Consideration should be given to starting at a dose lower than 5 mg for such patients. In cases of moderate hepatic impairment, the starting dose should be 5 mg and increased only with caution.

Children: Olanzapine is not recommended for use in children and adolescents below 13 years of age due to a lack of data on safety and efficacy.

Overdose:

Symptoms: in overdose include, agitation/aggressiveness, dysarthria, various extrapyramidal symptoms, delirium, convulsion, coma, possible neuroleptic malignant syndrome, respiratory depression, hypertension or hypotension, cardiac arrhythmias and cardiopulmonary arrest.

Treatment: There is no specific antidote for olanzapine. Induction of emesis is not recommended. Standard procedures for management of overdose may be indicated (i.e., gastric lavage, administration of activated charcoal). The concomitant administration of activated charcoal was shown to reduce the oral bioavailability of olanzapine by 50% to 60%.

Symptomatic treatment and monitoring of vital organ function should be instituted according to clinical presentation, including treatment of hypotension and circulatory collapse and support of respiratory function.

Storage:

Keep out of reach of children. Store below 25 °C.

Packaging:

Each Olen carton box contains 20 or 30 Film Coated Tablets in two or three Alu/Alu blisters.

