

# REVACINOLON

Suppositories & Rectal Cream & Ointment



## Composition:

Each REVACINOLON Suppository contains:

|                        |          |
|------------------------|----------|
| Fluocinolone Acetonide | 100 mcg. |
| Lidocaine HCl          | 40 mg.   |
| Bismuth Subgallate     | 100 mg.  |
| Menthol                | 5 mg.    |

Each 1 g of REVACINOLON Rectal Cream or Ointment contains:

|                        |          |
|------------------------|----------|
| Fluocinolone Acetonide | 100 mcg. |
| Lidocaine HCl          | 20 mg.   |
| Bismuth Subgallate     | 50 mg.   |
| Menthol                | 2.5 mg.  |

## Pharmacological Classification:

Antihemorrhoidal Drugs.

## Intended Use (Common Use of the Drug):

Treatment of hemorrhoids and anal purities and proctitis.

## Mechanism of Action:

- Fluocinolone acetonide is a corticosteroid that inhibits the inflammatory response to a variety of stimuli including multiple inflammatory cytokines. It prevents edema, fibrin deposition, capillary dilation, and scar formation associated with inflammation.

- Lidocaine hydrochloride has a local anesthetic effect, acting on sodium channels on the inner surface of nerve cell membranes, relieving pain and discomfort in affected areas.

- Bismuth Subgallate exert a protective action on mucous membranes. They have a mild astringent effect and have antiseptic and antihemorrhagic properties.

- Menthol provides a cooling sensation when applied to the skin, which helps to relieve pain in the subcutaneous tissues.

## Pharmacokinetics:

REVACINOLON has a local action with minimal of systemic effects. Fluocinolone can be absorbed from normal intact skin, and systemic absorption is less than 0.1 ng/ml. It is metabolized primarily in the liver and excreted by the kidneys. Lidocaine is partially absorbed but plasma levels will be low. It is metabolized in the liver and about 90% of a given dose undergoes de-ethylation. The rest of the active ingredients in REVACINOLON product exert their therapeutic effect without systemic absorption.

## Indications:

- Use this medicine exactly as described in this leaflet or as directed by your doctor or pharmacist, consult your doctor if you are not sure.

- REVACINOLON are used to relieve symptoms of pain, swelling and irritation associated with hemorrhoid, anal fissures and anal pruritus.

## Contraindications:

- This medicine should not be used if you are hypersensitive to the active substance or any of the excipients.

- Severe viral or bacterial infections for which appropriate antimicrobial treatment is not provided.

- Children under 2 years of age.

## Side Effects:

- Like all medicines, this one can cause side effects, although not everyone experiences them.

- Very rarely increased irritation may occur at the site of application

- Inform your doctor or pharmacist if you experience any side effects not listed in this leaflet.

## Warnings & Precautions:

- REVACINOLON should be used with caution and under the supervision of a physician in the event of:

- Use for more than 7 days due to the possibility of systemic absorption.

- Persistence or worsening of symptoms.

- People with severe rectal bleeding.

- Use in children under 12 years of age.

- It is recommended to separate topical preparations by two hours.

## Drug and Food Interactions:

- Tell your doctor or pharmacist if you are using, have recently used or might use any other drugs, and for how long you will be using the drug.

- Concurrent use with other corticosteroid preparations, either topical or oral, may increase the potential for systemic effects.

## Pregnancy & Lactation & Fertility:

**Pregnancy:** There is no conclusive evidence of the safety of REVACINOLON during pregnancy.

**Lactation:** There is no conclusive evidence of the safety of REVACINOLON during breastfeeding.

**Fertility:** Its effect on fertility after topical application in humans has not been studied.

## Effects on ability to drive and use machines:

REVACINOLON topical has no or negligible influence on the ability to drive and use machines.

## Method of use:

- Use this drug exactly as your doctor or pharmacist has prescribed it. Ask your doctor if you are not sure.

Adults and children 12 years and over:

**Suppositories:** One rectal suppository taken in the morning and evening and after each bowel movement, not exceeding three suppositories per day as a maximum.

**Cream or Ointment:** may be intrarectally applied by using the special applicator provided.

The application of cream or ointment is 2 - 3 times daily.

Cream or Ointment should be gently applied on the affected area after using warm anal bath.

- Inform your doctor or pharmacist if you forget to apply the medicine and before stopping its use.

## Overdoses/Toxicity/methods of treatment:

No cases of overdose have been reported.

## Physical Properties:

- Yellow, smooth, uniform, free from foreign matter Cream or Ointment.

- Yellow suppositories.

## Storage Conditions:

Store at a temperature below 30° C.

Keep out of reach of children.

Keep tube tightly closed.

## Packaging:

Each REVACINOLON suppositories carton box contents 10 suppositories in two PVC/PE strips.

Each REVACINOLON cream or ointment carton box contains Aluminum Tube of 20 g, special applicator attached.

\*A medication is a product which affects your health, and it's consumption contrary to instructions is dangerous for you.  
- Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medication.  
- The doctor and the pharmacist experts in medicine, it's benefits and risks.  
- Do not repeat the same prescription without consulting your doctor.  
**KEEP THE MEDICATIONS OUT OF REACH OF CHILDREN**  
Council of Arab Health Ministries - Union of Arab Pharmacies

