

REVAFURYL

Oral Suspension - Capsule



Composition:

Oral Suspension: Each 5 ml of REVAFURYL Oral Suspension contains Nifuroxazide 218 mg equivalent to 200 mg / 5 gr of suspension.

Capsules: Each REVAFURYL Capsule contains Nifuroxazide 200 mg.

Pharmacological Properties:

Nifuroxazide is an antibacterial belonging to the nitrofur group. It has an Intestinal antibacterial effect by direct contact with the inner lumen of the intestine, as it is poorly absorbed from the gastro-intestinal tract.

Indications:

REVAFURYL is indicated in the following cases:

- Acute diarrhea of bacterial origin.
- Inflammatory bowel.
- Acute and chronic colitis.
- As an intestinal disinfectant in all other types of diarrhea and intestinal infections.
- Prevention of infection of the intestine with opportunistic germs.

Contraindications:

- Hypersensitivity to nitrofur, nifuroxazide, or any of the components of the product.
- Premature infants and newborns under the age of one month.

Warnings & Precautions:

- If there is blood or mucus associated with the stool.
- If there is a fever, vomiting, or other associated digestive disturbances.
- You should see a doctor if the diarrhea continues after three days of starting treatment.
- Nifuroxazide can cause acute pancreatitis, so if you see the following symptoms: (nausea, Vomiting accompanied by abdominal pain) treatment should be stopped immediately and a doctor should be consulted.
- Nifuroxazide does not improve rehydration, so its use must be accompanied by a specific dietary schedule to prevent dehydration and loss of minerals and vitamins, it is recommended to eat rice, bananas, grilled meat and mint tea in the event of diarrhea, and it is preferable to stay away from milk and its derivatives.

Pregnancy & Lactation:

Pregnancy: Although studies on experimental animals exclude the distorting effect of fetuses when using nifuroxazide, there are no adequate studies to prove this, it is preferable not to use nifuroxazide during pregnancy without consulting a doctor, and treating diarrhea in pregnant women by modifying the food intake only.

Lactation: A nursing woman can continue breastfeeding her child in the event that her use of nifuroxazide is for a short period (not exceeding three days).

Drug Interactions:

- In order to avoid interactions with other medicines, the doctor or pharmacist must be informed of any other treatment the patient takes it.
- This medicine should not be used concomitantly with the use of sedatives (benzodiazepines: Calpromazepam, alprazolam, lorazepam, midazolam, and others).

Side Effects:

Nifuroxazide can cause the following side effects:

- Quincke's edema: This is angioedema of the tissues under the skin and manifesting an allergic reaction.
- Urticaria.
- Anaphylactic shock.

Dosage & Administration:

Children over two years: 5 ml of the suspension 3 times a day, for 3 days.

Adults: one capsule 4 times a day.

- The maximum daily dose of nifuroxazide is 800 mg.
- The treatment period should not last more than 7 days.
- It is not recommended for use in children under 2 years of age.
- The use of the medicine is not related to eating (it can be used before or after meals).

Storage:

Store in a dry place at a temperature below 25 °C and protect from light.

Packaging:

Oral Suspension: Each carton box REVAFURYL Oral Suspension contains amber glass bottle of 100 ml.

Capsules: Each carton box REVAFURYL Capsules contains 20 Capsules in two blister strips.

*A medication is a product which affects your health, and it's consumption contrary to instructions is dangerous for you.
- Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medication.
- The doctor and the pharmacist experts in medicine, it's benefits and risks.
- Do not repeat the same prescription without consulting your doctor.
KEEP THE MEDICATIONS OUT OF REACH OF CHILDREN
Council of Arab Health Ministers, Union of Arab Pharmacists

Reva Pharmaceutical Industry - Syria

التداخلات الدوائية:

التأثيرات الجانبية:

- الصدمة التأقية.

البالغون: كبسولة واحدة

- يجب ألا تدوم فترات

الحفظ:

التَّعَبُّة:

الكبسول: كل عبوة

0.0000

الكبسول: كل كبسولة ريفافوريل تحتوي على
نيفوروكسازيد ٢٠٠ ملغ.

يُستطب ريفافوريل في الحالات التالية:

- الإسهال الحاد ذو المنشأ الجرثومي.

- التّهاتات القولون الحاد و المزمن.

وأخماج الأمعاء.

مضادات الاستطباب:

أو أي من مكونات المستحضر.

التحذيرات والاحتياطات:

- في حال وجود حمى أو إقباءات أو اضطرابات

- بحسب مراجعة الطبيب في حال استمرار الإسهال

- يمكن أن يسبب النيفوروكسازيد التهاب البنكرياس

(غُثَّانٌ، اقْبَاءٌ مِثْرَ افِقٍ مَعَ أَلَمٍ فِي الْبَطْنِ) حَبٌّ فَوْراً

- نفور و کسازند لایحسین من الاماهة، لذلك حب أن

و فقدان المعادن و الفيتامينات، نصح بتناول الأرز و الموز

بالإسهال، وبفضل الابتعاد عن الحليب ومشتقاته.

الحمل: على الرغم من أن الدراسات على حيوانات

10 9 8 7 6 5 4 3 2 1

الدواء مستحضر يؤثر على صحتك واستعماله خلافاً للتعليمات هو من خطا
بيع دواء وصفه الطبيب بطريقة الاستعمال المتضمن في تعليمات الصيدلي
التي ترافقها ، فاقطعوا وصيدلي ما يبيعان في الدواء وينتفعه وضرره .
لا تكرر صرف الدواء بدون وصفة طبية .
لا تترك الأدوية بمسالك أيدي الأطفال .
مجلس وزراء الصحة العرب اتحاد الصيدلة العرب

إنتاج شركة ريفال للصناعات الدوائية - سوريا