

Reva-Flu

Film Coated Tablets - Syrup



Composition:

Tablets: Each Reva-Flu Film Coated Tablet contains:

Paracetamol 650 mg.

Chlorpheniramine Maleate 4 mg.

Phenylephrine Hydrochloride 10 mg.

Syrup: Each 5 ml of Reva-Flu contains:

Paracetamol 160 mg.

Chlorpheniramine Maleate 1 mg.

Phenylephrine Hydrochloride 2.5 mg.

Pharmacological Properties:

Reva-Flu has a special combination between paracetamol with its both analgesic and antipyretic activity due to inhibition of the synthesis of prostaglandins in the central nervous system. Chlorpheniramine maleate with its anti-allergic effect due to its blocking of histamine receptors H₁ (which makes it useful in cases of rhinorrhea and allergic cough). Phenylephrine which is a recognised decongestant and acts by vasoconstriction to reduce oedema and nasal swelling.

Pharmacokinetics:

Paracetamol is rapidly and almost completely absorbed from the gastrointestinal tract. It is metabolized in the liver and excreted in the urine mainly as glucuronide and sulphate conjugates.

Chlorpheniramine is well absorbed from the gastro-intestinal tract, following oral administration. Its effects begin to appear within 30 minutes, reach a maximum within 1-2 hours and last 4-6 hours. The half-life of chlorpheniramine is approximately 12 - 15 hours. Chlorpheniramine is metabolized to the monodesmethyl and didesmethyl derivatives. About 22% of an oral dose is excreted unchanged in the urine.

Phenylephrine HCl is irregularly absorbed from the gastrointestinal tract and undergoes first-pass metabolism by monoamine oxidase in the gut and liver; orally administered phenylephrine has reduced bioavailability. It is excreted in the urine almost entirely as the sulphate conjugate.

Indications:

Reva-Flu is indicated for the relief of the symptoms of colds and influenza, including the relief of aches and pains and nasal congestion, runny nose, sore throat and lowering of temperature.

Contraindications:

- Hypersensitivity to Paracetamol, Phenylephrine, Chlorpheniramine or to any of the excipients.
- Severe coronary heart disease, cardiovascular disorders and hypertension.
- It should not be used in patients with: Prostate enlargement, hyperthyroidism, glucose-6-phosphate dehydrogenase deficiency, epilepsy, severe renal or hepatic impairment.

Warnings & Precautions:

- Reva-Flu should be used with caution in patient with: Raynaud's phenomenon, diabetes mellitus or closed angle glaucoma.
- Asthma, chronic obstructive pulmonary disease and pyloroduodenal obstruction.
- Hemolytic anemia, constipation, vertigo and urinary problems.

Pregnancy & Lactations:

Due to the phenylephrine content of Reva-Flu, it should not be used during pregnancy or breastfeeding without consulting a doctor.

Drug Interaction:

- The anticoagulant effect of warfarin and other coumarins may be enhanced by prolonged regular daily use of paracetamol.
- Monoamine oxidase inhibitors: Hypertensive interactions occur between phenylephrine and MAOIs.
- Phenylephrine may reduce the efficacy of beta-blockers and antihypertensives.
- Tricyclic antidepressants: May increase the risk of cardiovascular side effects with phenylephrine.
- Concurrent use of Reva-Flu and hypnotics (e.g., barbiturates) or anxiolytics may cause an increase in sedative effects.
- Chlorpheniramine inhibits phenytoin metabolism and can lead to phenytoin toxicity.

Side Effects: Common side effects may include: Dizziness, drowsiness, blurred vision, dry mouth and nose, constipation, feeling nervous.

Driving & Using Machines:

This medicine has major influence on the ability to drive and use machines due to possible side effect drowsiness, particularly in the beginning of the treatment.

Dosage & Administration:

- **Tablets:** Adults and children above 12 years: The usual recommended dose is 1 tablet 3-4 times daily.

- **Syrup:**

Children 6-12 years: 10 ml, 3-4 times daily.

Children 2-6 years: 5 ml, 3-4 times daily.

Overdose:

Symptoms: Chlorpheniramine overdose can result in seizures, loss of consciousness, dystonic reactions, cardiovascular collapse including arrhythmias and maybe causes coma.

paracetamol overdose can result to poisoning, especially in the elderly, young children, and patients with liver disease. Phenylephrine overdosage is likely to result in effects similar to those listed under adverse reactions. Additional symptoms may include hypertension and possibly reflex bradycardia. In severe cases confusion and seizures may occur.

Treatment: Patients should be referred to hospital urgently for immediate medical attention. Treatment with activated charcoal should be considered if the overdose has been taken within 1 hour. Treatment with N-acetylcysteine may be used up to 24 hours after ingestion of paracetamol.

Storage:

Keep out of reach of children, store below 25°C and in the original package.

Packaging:

Tablets: Each Reva-Flu carton box contains 20 Film Coated Tablets in two blister Strips.

Syrup: Each Reva-Flu carton box contains a 100 ml opaque bottle.

- A medication is a product which affects your health, and it's consumption contrary to instructions is dangerous for you.
- Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medication.
- The doctor and the pharmacist experts in medicine, it's benefits and risks.
- Do not repeat the same prescription without consulting your doctor.
KEEP THE MEDICATIONS OUT OF REACH OF CHILDREN
Council of Arab Health Ministers - Union of Arab Pharmacists

Reva Pharmaceutical Industry - Syria

