

Spiramycin Reva

Film Coated Tablets



Composition:

- Each Spiramycin Reva 1.5 Film Coated Tablet contains: Spiramycin 1.500.000 IU.
- Each Spiramycin Reva 3 Film Coated Tablet contains: Spiramycin 3.000.000 IU.

Mechanism of Action:

Spiramycin is a macrolide antibiotic and its antibacterial effect is due to the inhibition of protein synthesis by binding to the 50S subunit of bacterial ribosomes. It is primarily bacteriostatic, but may be bactericidal against more sensitive strains when used in high concentrations.

Pharmacokinetics:

Spiramycin is rapidly, but incompletely, absorbed after oral administration, with a bioavailability of 30 to 39%. Its half-life is approximately 8 hours. It binds to plasma proteins by 10-25%. It is metabolized in the liver to active metabolites. More than 80% of the administered dose is excreted in the bile, while only about 4 to 14% is excreted in the urine.

Indications:

Spiramycin Reva is used in the treatment of the following diseases:

- Infections due to susceptible bacteria and upper respiratory tract infections, bronchopulmonary, cutaneous, stomatological (oral) and non-gonococcal genital infections.
- Toxoplasmosis in pregnant women.
- It can be used to prevent meningococcal infection, in cases where rifampicin cannot be used.
- In the prevention of rheumatic fever attacks in case of penicillin allergy.

Contraindications:

Hypersensitivity to the active ingredient, to any of the other macrolides, or to any of the components of the product.

Warning & Precaution:

- QT interval prolongation has been reported in patients taking macrolides, including spiramycin.
- Very rare cases of hemolytic anemia have been reported in patients with G-6 PD deficiency. Spiramycin is therefore not recommended for such patients.
- Cases of severe skin reactions including Stevens-Johnson syndrome, Toxic epidermal necrolysis (Lyell's syndrome), acute generalized exanthematous pustulosis have been reported with spiramycin. Patients should be informed of the signs and symptoms and close skin monitoring should be performed.

Pregnancy & Lactation:

Pregnancy: Spiramycin can be used during pregnancy if necessary. Extensive use of spiramycin during pregnancy has shown no evidence of malformation or fetotoxicity to date.

Lactation: breast-feeding is not recommended to women taking spiramycin.

Drug Interactions:

- Combination of spiramycin with carbidopa and levodopa causes inhibition of carbidopa reuptake and a decrease in levodopa plasma levels. When needed, patients should be monitored closely and the dose of Carbidoa adjusted.
- Although some ischemia cases have been reported in the concomitant use of macrolides with ergot derivatives, there is no report on spiramycin. When needed, it should be used with extreme caution.
- Class IA Antiarrhythmics (e.g., quinidine, hydroquinidine), class III Antiarrhythmics (e.g., amiodarone, sotalol), sultopride, cisapride, levofloxacin, moxifloxacin: There is an increased risk of ventricular arrhythmia disorders, particularly Torsades de Pointes.

Side Effects:

- Gastrointestinal effects: nausea, vomiting, diarrhea and very rare cases of pseudo-membranous colitis.
- Liver: very rare cases of liver tests function abnormalities.
- Hypersensitivity reactions: skin rash, urticaria, pruritus, very rarely angioedema and anaphylactic shocks. Isolated cases of vasculitis, including Henoch-Schönlein purpura.
- Peripheral and central nervous system: occasional cases of transient paresthesia.
- Hematology: very rare cases of acute hemolysis have been reported.

Driving & Using Machines:

Spiramycin Reva is not expected to affect the ability to drive or use machines.

Dosage & Administration:

Usual adult and adolescent dose:

- Antibacterial: Oral, 3 – 4.5 MIU two times a day; or 1,5 - 3 MIU three times a day. For severe infections, the dose may be increased to 6- 7,5 MIU two times a day.
- Gonorrhea: 12 to 13,5 MIU in a single dose.

Usual pediatric dose:

- Antibacterial: Children 20 kg of body weight and over: Oral, 75,000 IU per kg of body weight two times a day, or 50,000 IU per kg of body weight three times a day.

Overdose:

Symptoms: No case of accidental overdosage has been reported. In oral doses over 4g per day, abdominal discomfort, nausea or diarrhea may occur.

Treatment: No specific treatment has been proposed.

Management should be symptomatic.

Storage:

Keep out of reach of children. Store below 25° C.

Packaging:

Each Spiramycin Reva (1.5- 3) carton box contains 10 Film Coated Tablets in a blister strip.

-A medication is a product which affects your health, and it's consumption contrary to instructions is dangerous for you.
-Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medication.
-Do not repeat the same prescription without consulting your doctor.
-KEEP THE MEDICATIONS OUT OF REACH OF CHILDREN
Council of Arab Health Ministries, Union of Arab Pharmacists

Reva Pharmaceutical Industry - Syria

