

TEMPRO

Oral Drops - Syrup - Oral Suspension - Capsules



Composition:

Oral Drops: Each 1 ml TEMPRO Oral Drops contains:

Paracetamol (Acetaminophen) 100 mg.

Syrup: Each 5 ml TEMPRO Syrup contains:

Paracetamol (Acetaminophen) 160 mg.

Oral Suspension: Each 5 ml TEMPRO 200 Oral Suspension contains: Paracetamol (Acetaminophen) 200 mg.

Capsules: Each TEMPRO 500 Capsule contains:

Paracetamol (Acetaminophen) 500 mg.

Mechanism of Action:

Paracetamol is an effective analgesic and antipyretic. It works by inhibiting the synthesis of prostaglandins in the central nervous system and to a lesser extent through a peripheral effect by blocking the generation of pain impulses.

It is possible that paracetamol's antipyretic effect is due to its central effect on the hypothalamic thermoregulatory center.

Pharmacokinetics:

Paracetamol is rapidly absorbed from the gastrointestinal tract. The concentration in plasma reaches a peak in 30 minutes to 2 hours after taking the dose orally. It is metabolized in the liver and excreted in the urine mainly as the glucuronide and sulphate conjugates. Less than 5% of the administered dose is removed unchanged. The elimination half-life varies from about 1 to 4 hours.

Indications:

TEMPRO is an analgesic used for the relieve of mild to moderate pain including: Headache, sore throat, musculoskeletal pains, fever and pain after vaccinations, pain after dental procedures/tooth extraction, toothache. TEMPRO is also used for relief of fever.

Contraindications:

- Hypersensitivity to paracetamol or to any of the excipients.
- Patients with severe hepatic dysfunction.

Warnings & Precautions:

- If fever persists for more than three days (72 hours) or the pain persists for more than 5 days, consult a physician.
- Do not give this medicine to your child for more than 3 days without consult doctor or pharmacist.
- Care is advised in the administration of paracetamol to patients with severe renal or severe hepatic impairment.

Pregnancy & Lactation:

Pregnancy: Available data on the use of paracetamol in pregnant women indicate that there are no toxic (deform) effects in the newborn. Paracetamol can be used during pregnancy, if necessary, but it should be used at the lowest effective dose, for the shortest possible time and as frequently as possible.

Lactation: Paracetamol is excreted in breast milk, but not in large quantities. The available data show that it is not contraindicated in this case.

Drug Interactions:

- The hepatotoxicity of paracetamol, particularly after over-dosage, may be increased by drugs which induce liver enzymes such as barbiturates, tricyclic antidepressants, and alcohol.
- The speed of absorption of paracetamol may be increased by metoclopramide or domperidone and absorption reduced by colestyramine.

Side Effects:

Adverse effects of paracetamol are rare but hypersensitivity including skin rash may occur. There have been reports of

blood dyscrasias rare including thrombocytopenia and agranulocytosis, but the occurrence of such cases was not necessarily related to taking paracetamol.

Dosage & Administration:

Children under 10 years old:

No more than 4 doses should be given in a 24 hour period. Do not give this medicine to your child for more than 3 days without consulting doctor or pharmacist.

TEMPRO Oral Drops:

- 3-5 months: Half a dropper, 3-4 times a day after feeding.

- 6-12 months: One dropper, 3-4 times a day after feeding.

- 1-2 years: 1-1.5 dropper, 3-4 times a day after food.

The dropper capacity is 1ml. TEMPRO drops can be given directly into the mouth with a measured dropper or can be mixed with fruit juice or water, while TEMPRO syrup is given as a teaspoon.

TEMPRO Syrup 160 mg/5ml:

- 6-12 months: 3 ml, 3-4 times daily.

- 1-4 years: 3-5 ml, 3-4 times daily.

- 4-6 years: 5-7.5 ml, 3-4 times daily.

- 6-10 years: 10-15 ml, 3-4 times daily.

TEMPRO 200 Oral Suspension 200 mg/5ml:

- 6-12 months: 2.5 ml, 3-4 times daily.

- 1-5 years: 2.5-5 ml, 3-4 times daily.

- 6-12 years: 5-10 ml, 3-4 times daily.

Children 10-15 years: One capsule of TEMPRO 500 mg every 4 hours as needed, with a maximum of 4 capsules in 24 hours, it is not recommended for children under 10 years of age.

Adults and children over 16 years: 1-2 capsules of TEMPRO 500 mg up to four times daily as needed, up to a maximum of 8 capsules in 24 hours, the dose should not exceed 4g in 24 hours.

Overdose:

Hepatic damage can occur in adults (if they took 6g or more of paracetamol) and in children (if they took more than 150 mg/kg of paracetamol).

Treatment: Prompt treatment is essential in managing paracetamol overdose. The patient should be referred to hospital urgently to obtain immediate medical attention. The N-acetylcysteine antidote is given intravenously within 24 hours of taking paracetamol. If vomiting is not present, methionine may be given orally as a suitable alternative (out of hospital).

Storage:

Keep out of reach of children.

Store TEMPRO 200 Oral Suspension below 25 °C.

Store TEMPRO (Oral Drops - Syrup - Capsules) at room temperature between (15-30) °C.

Packaging:

Oral Drops: Each TEMPRO Oral Drops carton box contains amber glass bottle of 30 ml attached to a 1 ml dropper.

Syrup: Each TEMPRO Syrup carton box contains amber glass bottle of 100 ml.

Oral Suspension: Each TEMPRO 200 Oral Suspension carton box contains amber glass bottle of 100 ml.

Capsules: Each TEMPRO 500 carton box contains 20 Capsules in two blister strips.

Or Each TEMPRO 500 Cross box contains 500 Capsules in 50 blister strips.

-A medication is a product which affects your health, and it's consumption contrary to instructions is dangerous for you.
-Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medication.
-The doctor and the pharmacist experts in medicine, it's benefits and risks.
-Do not repeat the same prescription without consulting your doctor.
-KEEP THE MEDICAMENTS OUT OF REACH OF CHILDREN
Council of Arab Health Ministries, Union of Arab Pharmacies

Reva Pharmaceutical Industry - Syria

