

VOLTAC 100 S.R

Capsules



Composition:

Each VOLTAC 100 S.R Capsule contains:
Diclofenac sodium as sustained release pellets 100 mg.

Mechanism of Action:

Diclofenac sodium is a nonsteroidal anti-inflammatory drug that possesses anti-inflammatory, analgesic and antipyretic activity. The therapeutic effects are due to its cyclooxygenase inhibitory effect, which leads to a significant decrease in the synthesis of prostaglandins.

Pharmacokinetics:

Diclofenac is 100% absorbed after oral administration. However, due to metabolism during the first hepatic passage, only about 50% of the absorbed dose reaches the systemic circulation.

Diclofenac is more than 99% bound to human serum proteins, primarily to albumin. Diclofenac diffuses into and out of the synovial fluid. Biotransformation of diclofenac takes place mainly by single and multiple hydroxylation and methoxylation, resulting in several phenolic metabolites, most of which are converted to glucuronide conjugates. The metabolites are excreted in the urine (65%) and in the bile (35%).

Indications:

VOLTAC 100 S.R is used to relieve pain and inflammation of all degrees in a wide range of cases, including:

- Arthritic: rheumatoid arthritis, osteoarthritis, ankylosing spondylitis, acute gout,
- Acute musculo-skeletal disorders: such as periarthritis (for example frozen shoulder), tendinitis, tenosynovitis and bursitis.
- Other painful conditions resulting from trauma, including fracture, low back pain, sprains, strains, dislocations, orthopaedic, dental and other minor surgery.

Contraindications:

VOLTAC 100 S.R is contraindicated in patients with:

- Known hypersensitivity to the active substance or to any of the excipients.
- History of asthma, urticaria, or allergic-type reactions after taking aspirin or other NSAIDs.
- Active gastric or intestinal ulcer, bleeding or perforation of the digestive tract.
- Hepatic failure and renal failure.
- Congestive heart failure (NYHA-II-IV), ischemic heart diseases, peripheral arterial diseases and/or cerebrovascular diseases.

Warnings & Precautions:

- Like other NSAIDs, diclofenac may mask the signs and symptoms of infection.
- Allergic reactions, including anaphylactic/anaphylactoid reactions can occur without earlier exposure to diclofenac.
- Caution is recommended when using diclofenac after gastro-intestinal surgery and in patients with ulcerative colitis, as their condition may be exacerbated.
- To reduce gastrointestinal risk in patients with a history of ulcers, treatment should be initiated and maintained at the lowest effective dose.
- During prolonged treatment with diclofenac, regular monitoring of hepatic and renal function is recommended.
- Patients with congestive heart failure (NYHA-I) or patients with significant risk factors for cardiovascular disease (e.g., hypertension, hyperlipidaemia, diabetes mellitus, smoking) should only be treated with diclofenac after careful consideration.
- Diclofenac may reversibly inhibit platelet aggregation. Patients with coagulation defects, haematological abnormalities, or a predisposition to bleeding should be carefully monitored.
- Diclofenac can precipitate bronchospasm if administered to patients suffering from, or with a previous history of bronchial asthma.

Pregnancy & Lactation:

Pregnancy: Pregnancy Category C. During the 1st and 2nd trimester of pregnancy, VOLTAC 100 S.R should not be

given unless clearly necessary. During the last trimester of pregnancy VOLTAC 100 S.R is contraindicated.

Lactation: Diclofenac passes into the breast milk in small amounts. Therefore, diclofenac should not be administered during breast feeding.

Drug Interactions:

- Lithium, Phenytoin, Methotrexate and Digoxin: If used concomitantly, diclofenac may raise plasma concentrations of these drugs.
- Diuretics and Anti-hypertensive drugs: concomitant use of diclofenac with diuretics or antihypertensive drugs may cause a decrease in their antihypertensive effect.
- Drugs known to cause hyperkalemia: Concomitant treatment with potassium-sparing diuretics, ciclosporin, tacrolimus or trimethoprim may be associated with increased serum potassium levels.
- Anticoagulants and anti-platelet drugs: Caution is recommended since concomitant administration could increase the risk of bleeding.
- Other NSAIDs, selective serotonin reuptake inhibitors (SSRIs) and corticosteroids: Co-administration with these drugs may increase the risk of gastrointestinal bleeding or ulceration.
- Mifepristone: NSAIDs should not be used for 8-12 days after mifepristone administration as NSAIDs can reduce the effect of mifepristone.
- Cardiac glycosides: Concomitant use of cardiac glycosides and NSAIDs in patients may exacerbate cardiac failure, reduce GFR and increase plasma glycoside levels.
- Colestipol and cholestyramine: These drugs can decrease the absorption of diclofenac.

Side Effects:

Common side effects include: Headache, dizziness, vertigo, nausea, vomiting, diarrhoea, dyspepsia, abdominal pain, flatulence, decreased appetite, rash and transaminases increased.

Uncommon and Rare side effects include: Somnolence, tiredness, cardiac failure, palpitations, chest pain, gastritis, gastrointestinal ulcer with or without bleeding or perforation and hypersensitivity.

Driving & Using Machines:

Patients who experience side effects like visual disturbances, dizziness, vertigo and somnolence while taking NSAIDs should refrain from driving or operate machinery.

Dosage & Administration:

The capsule should be swallowed whole with liquid, preferably with or after food, and must not be divided or chewed.

Adults: One capsule VOLTAC 100 S.R daily. If necessary, the daily dosage can be increased to maximum (150 mg) by supplementation with the conventional dosage forms containing diclofenac sodium 25 mg or 50 mg.

Elderly: It is recommended that the lowest effective dosage be used in frail elderly patients or those with a low body weight.

Children: This medicine is not suitable for children.

Overdose:

Symptoms: headache, nausea, vomiting, epigastric pain, gastrointestinal haemorrhage, diarrhoea, dizziness, excitation, coma, drowsiness, fainting and convulsions. In the case of significant poisoning acute renal failure and liver damage are possible.

Treatment: Patients should be managed by symptomatic and supportive care following a NSAID overdose. There are no specific antidotes. Emesis and/or the use of activated charcoal and/or osmotic cathartic may be used in patients whose did not take the overdose more than 4 hours old.

Storage

Keep out of reach of children. Store below 30° C.

Packaging: Each VOLTAC 100 S.R capsule contains 20 capsules in two blister strips.

-A medication is a product which affects your health, and its consumption contrary to instructions is dangerous for you.
-Follow only the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medication.
-The doctor and the pharmacist experts in medicine, it's benefits and risks.
-Do not repeat the same prescription without consulting your doctor.
-KEEP THE MEDICATIONS OUT OF REACH OF CHILDREN.
Council of Arab Health Ministers' Union of Arab Pharmacies

Revva Pharmaceutical Industry - Syria

 ريفا فارما

الدواء مستحضر يؤخذ على صمغته واستعماله خلافاً لتعليمات بعض أطباء الخطر تتبع بنقطة وصفة الطبيب وطريقة الاستعمال المنصوص عليها وتعليمات الصيدلي التي صر بها لك ، فالطبيب والصيدلي هما الخبيران في الدواء ونفعه وضرره .
لا تذكر صرف الدواء بدون وصفة طبية .
لا تتناول الأدوية بمنقول أيدي الأطفال .
مجلس وزراء الصحة العرب والحاد للصحة العامة العرب .