

α -Lipovit

Enteric Coated Tablets



Composition:

Each α -Lipovit Enteric Coated Tablet contains:

Thiamine HCl (Vitamin B1)	250 mg
Pyridoxine HCl (Vitamin B6)	250 mg
Cyanocobalamin (Vitamin B12)	1 mg
Alpha-lipoic acid	300 mg

Pharmacological Properties:

α -Lipovit is a combination of three important and necessary neuroprotective vitamins (B1, B6 and B12) plus alpha lipoic acid, which are important for metabolism in the peripheral and central nervous system. Alpha lipoic acid has an effect on the body's metabolic functions. It also has neuroprotective (antioxidant) properties against harmful metabolic products. Therapeutic efficacy is greatly enhanced when vitamins B1, B6, B12 and alpha lipoic acid are combined, compared to the effect seen when these vitamins are given alone.

Pharmacokinetics: Thiamine (B1) is absorbed from the gastrointestinal tract and is widely distributed in most tissues of the body. It is not stored to an appreciable extent in the body and amounts in excess of requirements are excreted in the urine as thiamine or unchanged metabolites.

Pyridoxine (B6) is absorbed from the gastrointestinal tract and converted to the active form « pyridoxal phosphate ». It is excreted in the urine as 4-pyridoxic acid.

Cyanocobalamin (B12) is primarily stored in the liver, with a daily requirement of approximately 1 mcg. Its turnover is 2.5 mcg per day. Vitamin (B12) is excreted primarily in the bile and is largely reabsorbed through the enterohepatic circulation.

After oral administration, alpha lipoic acid is rapidly absorbed and distributed to tissues. Its metabolism mainly occurs by oxidative exchange of the side chain (beta-oxidation) and/or S-methylation of the corresponding thiols. It has been shown to be excreted mainly as metabolites by the kidney.

Indications: α -Lipovit is used for:

- Patients with risk factors for deficiency of vitamins B1, B6, B12 and/or patients with preclinical general complaints such as mental and physical fatigue, forgetfulness, irritability, weakness and tremor.
- Neuritis, nerve pain, diabetic neuropathy, shingles.
- Arthritis, periartthritis, sciatica, back pain, joint pain, muscle pain, spasticity and rheumatic pain.

- As an aid in the treatment of cardiomyopathy, vomiting after surgery, radiation sickness, rheumatic fever and chronic intoxication, especially in alcoholism.

Contraindications:

- α -Lipovit is contraindicated in patients known to be hypersensitive to any of the ingredients of the drug.
- Patients with renal or hepatic insufficiency.

Warnings & Precautions:

- Vitamin (B12) is not recommended in patients with « Leber's disease » due to the increased risk of optic atrophy.

- Due to the high vitamin (B6) content, α -Lipovit should not be taken in higher doses or for a longer period than recommended.

- The use of α -Lipovit is not recommended in patients with megaloblastic anemia.

- Because α -Lipovit contains alpha-lipoic acid, a change in urine odor without clinical significance may be observed.

Pregnancy & Lactation:

Vitamins B: There are no risks associated with using Vitamins B1, B6, and B12 during pregnancy at the recommended dose. Vitamins B1, B6, and B12 are excreted in breast milk, but no risk of overdosage has been reported in children.

Alpha lipoic acid: Alpha lipoic acid is mostly safe when taken orally. It has been used safely during pregnancy in doses of up to 600 mg daily for up to 4 weeks. In general, information regarding safety and effectiveness during pregnancy and lactation is not precisely available.

Driving & using machines:

α -Lipovit has no effect on the ability to drive and use machines.

Drug Interactions:

• Thiamine hydrochloride (vitamin B1): Both thiosemicarbazone and 5-fluorouracil inhibit the activity of thiamine. Antacids also block thiamine absorption.

• Pyridoxine Hydrochloride (Vitamin B6): High doses of Vitamin (B6) inhibit the effect of levodopa. Use of cycloserine, hydralazine, isoniazid, desoxypyridoxine, dipenicillamine, oral contraceptives, and alcohol may decrease blood levels of pyridoxine.

• Cyanocobalamin (Vitamin B12): High doses of folic acid can lower blood levels of vitamin (B12).

You should avoid taking vitamin C within one hour of taking vitamin (B12) by mouth, because blood levels may be too low. Alcohol, aminosulcylates, colchicine, aminoglycosides, antihistamines H₂, metformin, oral contraceptives, and proton pump inhibitors may reduce vitamin (B12) absorption.

• Alpha Lipoic Acid: May increase the hypoglycemic effect of insulin and/or medications used to treat diabetes mellitus. It is a metal chelator and therefore should not be used with products containing minerals (such as iron preparations).

Side Effects:

Common: dizziness.

Not known: allergic reactions including (urticaria, facial edema, erythema, rash and blisters), headache, peripheral neuropathy, somnolence, paresthesia. Increase in aspartate aminotransferase, decrease in blood folic acid levels, diarrhoea, indigestion, nausea and abdominal pain.

Very rare: disturbance of taste and sense of taste.

Dosage & Administration:

For prevention: one tablet daily.

For treatment: 2-4 tablets daily. The dose may be reduced when symptoms subside.

α -Lipovit should be swallowed with a sufficient amount of water on an empty stomach.

Taking it with food may decrease the absorption of alpha lipoic acid.

Overdose: There is no evidence that α -Lipovit can cause an overdose if it is used as recommended.

Treatment: No specific treatment has been suggested. Treatment should be occasional.

Storage: Keep out of reach of children.

Store below 30° C.

Packaging: Each α -Lipovit carton box contains 20 Enteric Coated Tablets in two blister strips.

« A medication is a product which affects your health, and its consumption contrary to instructions is dangerous for you. »
- Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medication.
- The doctor and the pharmacist experts in medicine, it's benefits and risks.
- Do not repeat the same prescription without consulting your doctor.
KEEP THE MEDICATIONS OUT OF REACH OF CHILDREN
Council of Arab Health Ministers & Union of Arab Pharmacists

Reva Pharmaceutical Industry - Syria



إنتاج شركة ريفال للصناعات الدوائية - سوريا

[illegible]

انتاج شركة ريفيا للصناعات الدوائية - سوريا