

Draft: thermographs, Newton polygons, and the missing residue

a9lim

June 2026

Abstract

This note pursues `docs/OPEN.md` problem `under*(e_g^e_s)`: whether the two tropical consumers in `ogdoad`—game thermography and valuation/Newton polygons—are one tropical object or merely a convention mirror. The result of this pass is a negative theorem at the thermograph level and a more precise positive target.

The negative theorem is small but decisive: the thermograph is not a congruence for disjunctive sum. There is no binary operation, single- or multi-step, that takes only $\text{Th}(G)$ and $\text{Th}(H)$ and returns $\text{Th}(G + H)$ for all short games. The witness is already in the temperature-zero layer: $\text{Th}(*) = \text{Th}(\uparrow)$, but $* + * = 0$ while $\uparrow + *$ is still a non-number of temperature 0.

The useful replacement is not “no tropical structure”. Temperature itself has exactly the expected tropical-valuation shape: $\text{temp}(G + H) \leq \max(\text{temp } G, \text{temp } H)$, with equal-temperature pairs forming the game-side vanishing locus. What fails is the proposed angular component $(\text{mean}(G), \text{temp}(G))$: it forgets the leading thermic residue. A single object, if it exists, should be an associated graded object of the game group by temperature, enriched at least by the all-small residue data (atomic weight plus its kernel), not the ordinary thermograph alone. I hit the wall at making that graded object multiplicative: the repo has no Norton/overheating product, and the game group itself is not a ring.

Claim levels. **[PROVED]** – proof from standard short-game identities plus the live implementation’s definitions; **[TESTED]** – verified by the commands in Appendix A; **[STANDARD MATH]** – standard thermography/Newton-polygon math used by the repo; **[CONJECTURAL]** – plausible but not proved here; **[OPEN]** – this pass did not solve it.

1 The two tropical objects currently in the code

[STANDARD MATH] The place axis is the usual non-Archimedean tropicalization. In `src/scalar/valued.rs`, a valued scalar satisfies

$$v(xy) = v(x) + v(y), \quad v(x + y) \geq \min(v(x), v(y)),$$

with equality when $v(x) \neq v(y)$. The adaptor `tropicalize` lands in the min-plus semiring. In `src/scalar/newton.rs`, $\text{NP}(f)$ is the lower convex hull of the points $(i, v(a_i))$, and its side slopes are the negatives of root valuations. The Springer tests in `src/forms/springer/local.rs` check that the Newton polygon forgets only the residue square-class data: its slopes are exactly the valuation layers.

[STANDARD MATH] The game axis is also tropical, but internally. A short game’s thermograph consists of a left wall $L_G(t)$, a right wall $R_G(t)$, the mast $\text{mean}(G)$, and the temperature $\text{temp}(G)$; the implementation lives in `src/games/thermography.rs`. The tropical naming layer makes the left wall the max-plus fold over Left options’ right walls, the right wall the min-plus

fold over Right options' left walls, and cooling ordinary addition of the temperature parameter, i.e. tropical multiplication. The test `via_tropical_matches_thermograph` pins this naming equal to the ordinary thermograph.

Remark 1. This is already enough to justify the slogan “thermography is tropical arithmetic”. It is not enough to make thermographs compose under disjunctive sum. Newton polygons have Dumas additivity under polynomial multiplication. Thermographs do not have an analogous strict additivity under game sum.

2 The thermograph is not a sum invariant

Definition 1. For a short game G , write

$$\text{Th}(G) = (L_G, R_G, \text{mean}(G), \text{temp}(G))$$

for the exact thermograph computed by `thermograph`. A thermograph-level sum law would be any operation Φ such that

$$\text{Th}(G + H) = \Phi(\text{Th}(G), \text{Th}(H))$$

for all games in the domain of ordinary thermography.

Proposition 1 (No thermograph-level composition law). **[PROVED]** *The map $G \mapsto \text{Th}(G)$ is not a congruence for disjunctive sum. Consequently, no operation whose inputs are only $\text{Th}(G)$ and $\text{Th}(H)$ can determine $\text{Th}(G + H)$ for all short games.*

Proof. Use the standard games $*$ = $\{0 \mid 0\}$ and $\uparrow$ = $\{0 \mid *\}$. The live thermography engine, and standard temperature theory, give

$$\text{Th}(*) = \text{Th}(\uparrow) = (0, 0, \text{mean} = 0, \text{temp} = 0),$$

where both walls are the constant zero wall.

Now add the same partner $*$. Since $*$ is the nimber of order two,

$$* + * = 0,$$

so the result is a number and the repo reports temperature -1 for the cold numeric case. But $\uparrow + *$ is still all-small and non-numeric; its thermograph has temperature 0 and constant zero walls. Thus the two input pairs

$$(\text{Th}(*), \text{Th}(*)) \quad \text{and} \quad (\text{Th}(\uparrow), \text{Th}(*))$$

are identical, while the output thermographs differ. This contradicts the existence of Φ . $\square$

Corollary 1. **[PROVED]** *The pair $(\text{mean}(G), \text{temp}(G))$ cannot be the game-side analogue of $(\text{ac}(x), v(x))$ for a valued field element. It forgets the leading thermic residue that decides cancellation.*

Remark 2. The witness is deliberately low-tech. It lives at temperature zero, before any subtle sidling or nested hot-game pathology appears. The obstruction is not an implementation artifact of complicated thermographs; it is present in the first infinitesimal layer.

3 What survives: temperature as a tropical valuation

[STANDARD MATH] Temperature does satisfy the valuation-shaped inequality

$$\text{temp}(G + H) \leq \max(\text{temp}(G), \text{temp}(H)),$$

with numbers treated as colder than all infinitesimal/hot games. Reversing the order by $v_T(G) = -\text{temp}(G)$ gives the familiar valuation form

$$v_T(G + H) \geq \min(v_T(G), v_T(H)).$$

The equal-temperature case is the game-side vanishing locus. It can cancel downward, or it can retain the same leading temperature.

[TESTED] In this checkout I probed a small catalogue consisting of integers, $*$, $\uparrow$, $\downarrow$, $*2$, switches $\{a \mid b\}$ for small integer a, b , and one-option nested hot games. Across 324 pairwise sums there were no violations of $\text{temp}(G + H) \leq \max(\text{temp } G, \text{temp } H)$. Every unequal-temperature pair hit equality. Equal-temperature pairs produced the expected downward spread:

$$\begin{aligned} \text{temp}(*) &= \text{temp}(*) = 0, & \text{temp}(* + *) &= -1, \\ \text{temp}(*) &= \text{temp}(\uparrow) = 0, & \text{temp}(* + \uparrow) &= 0, \\ \text{temp}(\{1 \mid -1\}) &= \text{temp}(\{2 \mid 0\}) = 1, & \text{temp}(\{1 \mid -1\} + \{2 \mid 0\}) &= -1. \end{aligned}$$

Other equal-temperature probes at input temperature 2 produced output temperatures $-1, 0, 1$, and 2 .

Remark 3. This is exactly the tropical-hyperfield shape if one keeps only the scalar temperature: unequal leading values add strictly by max-plus; equal leading values may cancel to any lower layer. The problem is that this scalar law is too coarse to recover walls, masts below the leading layer, or the resulting game value.

4 The one-object probe: switches versus one-side Newton polygons

[TESTED] There is a clean but shallow dictionary for switches. For $m \in \mathbb{Q}$ and $\tau > 0$, the switch

$$S_{m,\tau} = \{m + \tau \mid m - \tau\}$$

has

$$L_S(t) = m + \tau - t, \quad R_S(t) = m - \tau + t \quad (0 \leq t \leq \tau),$$

and both walls freeze to m at $t = \tau$.

The Newton polygon of a binomial $x^n - \pi^k$ has one side from $(0, k)$ to $(n, 0)$, hence one slope and one root-valuation layer. After an affine change of axes, one wall of $S_{m,\tau}$ is exactly such a one-side polygon; the other wall is its dual reflection. This realizes the requested “one-object probe” only in the trivial one-parameter family.

Proposition 2. [PROVED] *The switch/binomial dictionary does not extend to a thermograph-level sum theorem.*

Proof. The dictionary sends one switch to one linear Newton side. Polynomial multiplication concatenates Newton slope data by Dumas additivity. But Proposition 1 shows that disjunctive sum cannot be read from thermographs alone, even before leaving the zero-temperature layer. Any extension would need extra residue data not present in the switch wall. $\square$

5 The actual candidate bridge: an associated graded game object

[CONJECTURAL] The positive object should be a temperature filtration of the game group. Let

$$F_{\leq \tau} = \{G : \text{temp}(G) \leq \tau\}$$

for $\tau \geq 0$, with the numeric games placed in the cold bottom layer. The standard temperature inequality says these are additive subgroups. The associated graded candidate is

$$\text{gr}_T(\text{Games}) = \bigoplus_{\tau} F_{\leq \tau} / F_{< \tau}.$$

The leading thermic residue of G would be the class of G in the quotient at $\tau = \text{temp}(G)$, not the pair $(\text{mean}(G), \text{temp}(G))$.

[TESTED] At $\tau = 0$, this is already visible. The thermograph collapses $*$, $\uparrow$, $\downarrow$, and $*2$ to the same constant-zero object. Atomic weight, implemented in `src/games/atomic_weight.rs`, recovers one additive residue on all-small games:

$$\text{aw}(\uparrow) = 1, \quad \text{aw}(\downarrow) = -1, \quad \text{aw}(\ast) = 0.$$

But atomic weight still has a kernel containing nimber-like residues such as $*$, and $\ast + \ast = 0$ shows that this kernel matters. So even the first graded piece is not “the mast”. It is a genuine residual game object.

Remark 4. This is the closest analogue to the valued-field picture. The Newton polygon records valuations and forgets residues; the initial forms in $\text{gr}_v(K[x])$ restore them. The thermograph records temperatures and walls but forgets thermic residues; a graded game object would restore them. The difference is that $\text{gr}_v(K)$ is a graded ring, while short games under disjunctive sum are only an abelian group.

6 Where this pass hit the wall

[OPEN] I do not currently have a product on $\text{gr}_T(\text{Games})$ that would make it the literal peer of the graded ring of a valued field. The obvious ambient product is absent: full short games do not form a ring, and the repo correctly refuses to pretend otherwise. Norton multiplication or overheating might supply a product on restricted classes, but neither structure is present in the codebase, and this pass did not derive a canonical one from the thermograph recursion.

[OPEN] I also do not have a nontrivial thermograph hyperoperation. The scalar temperature hyperoperation is clear and useful:

$$a \boxplus b = \begin{cases} \max(a, b), & a \neq b, \\ \{\text{temperatures} \leq a\}, & a = b, \end{cases}$$

with the usual convention that colder numbers lie below 0. But lifting this from scalar temperatures to full thermographs without adding residues becomes either false (Proposition 1) or tautological (return “all thermographs of all sums of representatives”).

[OPEN] The next solvable formulation is therefore not “find a binary operation on thermographs”. It is:

Define the temperature filtration quotient $F_{\leq \tau} / F_{< \tau}$ for a manageable class of games, compute its first pieces, and decide whether Norton multiplication/overheating gives those pieces a product compatible with cooling.

If yes, the answer to the open problem is “one tropical object, but only after passing to a residue-enriched associated graded”. If no, the answer is “two tropical objects sharing only the scalar hyperfield shadow”.

A Checks run in this pass

The code-level facts used above were checked against the live repo with:

- `cargo test tropical -- --nocapture`: 15 relevant tests passed, including tropical semiring laws and `thermograph_via_tropical`.
- `cargo test newton -- --nocapture`: 8 Newton-polygon tests passed, including Eisenstein slopes and Dumas additivity.
- `cargo test springer -- --nocapture`: 50 Springer/local-global tests passed, including the Newton-polygon/Springer-shadow check.
- A Python probe through the installed `.venv` binding checked 324 small-game sums for the temperature inequality and the off-diagonal equality pattern, and printed the witnesses in Proposition 1.

References

- [1] Berlekamp, Conway, and Guy. *Winning Ways for your Mathematical Plays*.
- [2] Aaron Siegel. *Combinatorial Game Theory*.
- [3] Elwyn Berlekamp. The economist’s view of combinatorial games. In *Games of No Chance*, 1996.
- [4] Oleg Viro. Hyperfields for tropical geometry I.
- [5] Diane Maclagan and Bernd Sturmfels. *Introduction to Tropical Geometry*.