

Quadratic Witt Classes over $\mathbb{F}_2(t)$: Ramified Residue Coordinates and Impartial Realization Boundaries

GPT 5.6 Sol

a9lim

Abstract

The quadratic Witt group of the imperfect field $K = \mathbb{F}_2(t)$ carries wild local information absent from the finite-field Arf classification. Using the characteristic-two Milnor–Scharlau sequence of Aravire and Jacob, we prove that a class is determined canonically by its unramified coordinate at infinity together with all second residues. At a place v , the second residue consists of a finite odd-pole expansion over the residue field and one ramified Arf bit. We prove directly that every nonzero finite-field Scharlau transfer preserves the Arf generator. Consequently the single global relation is the parity of the ramified bits.

Finite-field trace characters turn every wild coefficient into a basis-free finite family of binary observations. Sending zero to the zero game and one to $*$ gives a faithful, orthogonal-sum-additive family of impartial normal-play values for every nonsingular Witt class. We derive the full finite triangular coordinate transformation under a ramified extension $\pi = u\varpi^e$, including its trace-character adjoint.

The stronger request for one fixed finite refinement-independent local-access family is impossible already for binary planes: the refinements $[1, P^{-1}]$, with P irreducible, have one fixed polar form and infinitely many distinct place coordinates. Thus a finite sparse family must either select its labels from the refinement, becoming a transducer, or be replaced by an infinite on-demand family. Finally, no additive impartial invariant can extend faithfully to arbitrary singular quadratic spaces: the quasilinear relation $\langle c \rangle \perp \langle c \rangle \cong \langle c \rangle \perp \langle 0 \rangle$ forces cancellation of the radical coefficient.

1 Introduction

Over a finite field of characteristic two, the quadratic Witt group is classified by one Arf bit [3, 5]. This finite situation admits an exact normal-play interpretation: after restriction of scalars to $\mathbb{F}_2$, a binary quadratic refinement can be compiled into an impartial arena whose losing loadings are its zero set [1]. Neither statement extends naively to $K = \mathbb{F}_2(t)$. The field is imperfect, and odd negative Laurent powers survive Artin–Schreier reduction at every place. The resulting wild terms are genuine Witt coordinates, not implementation noise.

Aravire and Jacob construct characteristic-two second residues and transfers and prove the exact sequence [2, Theorem 6.2]

$$0 \longrightarrow W_q(\mathbb{F}_2) \longrightarrow W_q(K) \xrightarrow{\oplus_v \partial_v} \bigoplus_v W_1(K_v) \xrightarrow{\oplus_v s_v^*} W_q(\mathbb{F}_2) \longrightarrow 0.$$

The notation is important: $W_1(K_v)$ is a quotient of the Witt group of the completion, not the Witt group of its residue field. Its wild summand is infinite-dimensional, although every individual residue has finite support.

This paper extracts the exact normal-play consequence of that sequence, computes ramified scalar extension on the resulting coordinates, and settles two stronger requests negatively: a fixed finite local-access family cannot cover the full refinement torsor, and arbitrary quasilinear radical data cannot survive an additive game law.

Theorem A. *Fix $K = \mathbb{F}_2(t)$ and the canonical uniformizers $p(t)$ at finite places and t^{-1} at infinity.*

1. *There is a canonical additive isomorphism*

$$W_q(K) \cong \mathbb{F}_2 \oplus \left\{ ((\psi_v, \epsilon_v))_v : \text{finite support and } \sum_v \epsilon_v = 0 \right\},$$

where ψ_v is the wild odd-pole residue and ϵ_v the ramified Arf bit.

2. *These coordinates have a finite, basis-free, faithful realization by the impartial values 0 and *, additive under orthogonal sum.*
3. *Under $k((\pi)) \hookrightarrow \ell((\varpi))$ with $\pi = u\varpi^e$, restriction is an explicit finite triangular transformation of the odd-pole coefficients, functorial in towers and adjoint on trace labels.*
4. *No fixed finite refinement-independent P/N family is faithful on all refinements of even one nonsingular binary polar space. Likewise, no isometry-invariant orthogonal-sum homomorphism into a cancellative game-value monoid can distinguish every quasilinear radical restriction.*

The first item uses the cited Milnor–Scharlau exact sequence. The finite-field transfer calculation, coordinate splitting, game realization, ramified transformation, and both impossibility theorems are proved here.

2 Local characteristic-two coordinates

Let v range over the monic irreducible polynomials $p \in \mathbb{F}_2[t]$ and the infinite place. Define

$$\pi_p = p(t), \quad \pi_\infty = t^{-1}.$$

Write K_v for the completion and κ_v for its residue field. Thus

$$\kappa_p = \mathbb{F}_2[t]/(p), \quad \kappa_\infty = \mathbb{F}_2.$$

Every κ_v is finite and therefore perfect. The local theorem of Aravire and Jacob specializes to [2, Theorem 1.3 and Corollary 1.7]

$$W_q(K_v) \cong W_q(\kappa_v) \oplus R_v \oplus \langle \pi_v \rangle W_q(\kappa_v),$$

where

$$R_v = \bigoplus_{\substack{n \geq 1 \\ n \text{ odd}}} \kappa_v \pi_v^{-n}.$$

The first summand is unramified. Quotienting it out gives

$$W_1(K_v) := \text{coker}(W_q(\kappa_v) \longrightarrow W_q(K_v)) \cong R_v \oplus \mathbb{F}_2.$$

For a global class α , write its localization as

$$\text{loc}_v(\alpha) = (\phi_{0,v}, \psi_v, \phi_{1,v}), \quad \psi_v = \sum_{n \text{ odd}} c_{v,n} \pi_v^{-n}.$$

Then

$$\partial_v \alpha = (\psi_v, \phi_{1,v}).$$

Only finitely many places occur for one global class, and each ψ_v is a finite sum. The uniformizer is part of this description: changing it changes the displayed odd-pole coordinates even when the underlying class is fixed.

3 Finite-field Scharlau transfer

We now identify the final map in the global sequence. The proof uses only the nondegenerate trace pairing of a finite field extension [6].

Lemma 3.1 (Transfer preserves the Arf generator). *Let $E/\mathbb{F}_2$ be finite and let $s : E \rightarrow \mathbb{F}_2$ be a nonzero linear functional. Under the Arf identifications*

$$W_q(E) \cong \mathbb{F}_2 \cong W_q(\mathbb{F}_2),$$

the Scharlau transfer $s_ : W_q(E) \rightarrow W_q(\mathbb{F}_2)$ is the identity.*

Proof. The trace pairing is nondegenerate, so $s(z) = \text{Tr}(cz)$ for a unique $c \neq 0$. Since E is perfect, choose $d \in E$ with $d^2 = c$. Let $a \in E$ satisfy $\text{Tr}(a) = 1$; the plane $[1, a]$ represents the nonzero class in $W_q(E)$. Its transfer along s is

$$s(x^2 + xy + ay^2) = \text{Tr}(cx^2 + cxy + cay^2).$$

After the simultaneous change of variables $X = dx$, $Y = dy$, this is

$$T_a(X, Y) = \text{Tr}(X^2 + XY + aY^2).$$

Choose a basis (e_i) of $E/\mathbb{F}_2$ and its trace-dual basis (f_i) . The vectors $(e_i, 0), (0, f_i)$ form a symplectic basis for the polar form of T_a . Therefore

$$\begin{aligned} \text{Arf}(T_a) &= \sum_i \text{Tr}(e_i^2) \text{Tr}(af_i^2) \\ &= \text{Tr} \left(a \sum_i \text{Tr}(e_i) f_i^2 \right). \end{aligned}$$

Trace duality applied to $1 \in E$ gives

$$1 = \sum_i \text{Tr}(e_i) f_i.$$

Squaring in characteristic two gives $1 = \sum_i \text{Tr}(e_i) f_i^2$. Hence

$$\text{Arf}(T_a) = \text{Tr}(a) = 1.$$

The nonzero class is therefore carried to the nonzero class. □

In the Aravire–Jacob construction, s_v^* kills R_v and applies ordinary Scharlau transfer to the ramified residue form. Lemma 3.1 therefore gives

$$s_v^*(\psi_v, \phi_{1,v}) = \phi_{1,v}.$$

Thus wild coefficients are locally essential but globally unconstrained; the only reciprocity relation is parity of the ramified bits.

4 Canonical global coordinates

The exact sequence determines classes modulo constants. A distinguished unramified local coordinate supplies the missing constant without choosing a vector-space basis.

Theorem 4.1 (Infinity plus second residues). *Define*

$$\Theta(\alpha) = (\phi_{0,\infty}(\alpha), ((\psi_v(\alpha), \phi_{1,v}(\alpha)))_v).$$

Then Θ is an additive isomorphism

$$W_q(\mathbb{F}_2(t)) \xrightarrow{\sim} \mathbb{F}_2 \oplus \left\{ ((\psi_v, \epsilon_v))_v : \text{finite support and } \sum_v \epsilon_v = 0 \right\}.$$

Proof. The transfer formula above identifies the kernel of the final arrow in the Milnor–Scharlau sequence with the displayed parity-zero residue tuples. The unramified coordinate $\phi_{0,\infty}$ is additive. On a constant-field class $a \in W_q(\mathbb{F}_2)$, localization at infinity has coordinates $(a, 0, 0)$, so $\phi_{0,\infty}$ restricts to the identity on constants.

Suppose $\Theta(\alpha) = 0$. All second residues vanish, so exactness gives α in the image of $W_q(\mathbb{F}_2)$. Its infinite unramified coordinate is that constant class, and hence the first coordinate makes it zero. This proves injectivity.

Conversely, let r be a finite residue tuple of parity zero. Exactness gives $\beta \in W_q(K)$ with residue r . Given a prescribed constant coordinate $a \in \mathbb{F}_2$, set

$$\alpha = \beta + (a - \phi_{0,\infty}(\beta)),$$

where the parenthesized term is included as a constant class. It changes no second residue and changes the first coordinate to a . Thus Θ is surjective. Injectivity makes the lift unique. $\square$

Remark 4.2. Exactness alone gives a noncanonical splitting because these groups are $\mathbb{F}_2$ -vector spaces. The theorem is stronger: the first local coordinate at the fixed infinite place is an explicit retraction, so no basis extension or choice of residue lifts appears in Θ .

5 A finite impartial realization

The wild coefficient $c_{v,n} \in \kappa_v$ is not itself a bit. Expanding it in a power basis would be finite but would introduce a coordinate choice. The trace pairing gives a redundant basis-free presentation instead.

Definition 5.1 (Trace-bit presentation). For $z \in \kappa_v$, define

$$b_{v,n,z}(\alpha) = \text{Tr}_{\kappa_v/\mathbb{F}_2}(zc_{v,n}).$$

Also put

$$b_c(\alpha) = \phi_{0,\infty}(\alpha), \quad b_{v,r}(\alpha) = \phi_{1,v}(\alpha).$$

Each bit is additive in α . Nondegeneracy of the trace pairing says that $c_{v,n} = 0$ exactly when every $b_{v,n,z}$ vanishes. The complete bit family is faithful by Theorem 4.1. It also has finite support: one class has finitely many places and pole orders, and each κ_v is finite.

Let 0 denote the optionless impartial game and let $*$ = $\{0 \mid 0\}$. These are the numbers 0 and 1, so $* + * = 0$ under disjunctive sum [4].

Theorem 5.2 (Sparse impartial compiler). *For every nonsingular quadratic Witt class α over $\mathbb{F}_2(t)$, replace each zero trace, constant, or ramified bit by the zero game and each one bit by $*$. Omit zero entries from the resulting labelled family. Then:*

1. *the family is finite and a coordinate is a P-position exactly when its bit is zero;*
2. *orthogonal sum becomes coordinatewise disjunctive sum;*
3. *hyperbolic classes map to the zero family; and*
4. *two families have the same outcome vector exactly when their source forms are Witt equivalent.*

Proof. Finiteness and faithfulness were proved above. The map $\mathbb{F}_2 \rightarrow \{0, *\}$ is an additive injection because $* + * = 0$. Applying it to every labelled bit proves the orthogonal-sum identity. Theorem 4.1 then identifies equality of all bits with equality in $W_q(K)$; the zero class consists precisely of hyperbolic forms. $\square$

This construction is not the Boolean serialization of one displayed rational function. Every bit is a homomorphism defined on the Witt group after passage through the exact residue sequence, so it is invariant under isometry and hyperbolic stabilization. It is nevertheless a classifier-to-star compiler: the residue has been computed before the one-move game is emitted.

6 Ramified coordinate change

We now compute the coordinate transformation that was left implicit in the classification-to-game compiler. Let

$$K = k((\pi)) \hookrightarrow L = \ell((\varpi)), \quad \pi = u\varpi^e,$$

where k and ℓ are finite fields, $f = [\ell : k]$, and $u \in \ell[[\varpi]]^\times$. Write $\bar{e}, \bar{f} \in \mathbb{F}_2$ for the parities of e, f . For every positive odd n , expand

$$u^{-n} = \sum_{r \geq 0} b_{n,r} \varpi^r.$$

Inverse Frobenius in the perfect field ℓ is denoted by $x \mapsto x^{2^{-s}}$.

The elementary Artin–Schreier reduction is finite and triangular. If $A = \sum_j A_j \varpi^j$ has finite principal part, then its canonical odd-pole coordinates are

$$\begin{aligned} \epsilon(A) &= \text{Tr}_{\ell/\mathbb{F}_2}(A_0), \\ \rho_d(A) &= \sum_{s \geq 0} A_{-2^s d}^{2^{-s}} \quad (d > 0 \text{ odd}). \end{aligned}$$

Indeed, adding

$$\wp(c\varpi^{-m}) = c^2\varpi^{-2m} + c\varpi^{-m}$$

kills an even pole and changes only the strictly smaller pole m . Descending cancellation gives the displayed sum. Positive powers lie in $\wp(\varpi\ell[[\varpi]])$, while an odd leading pole cannot occur in an Artin–Schreier image. The constant coefficient is unchanged by pole cancellation; any new constant below comes from the unit jet before reduction.

For a wild input $\psi = \sum_{n>0 \text{ odd}} c_n \pi^{-n}$, define

$$\begin{aligned}\eta_{e,u}(\psi) &= \sum_n \text{Tr}_{\ell/\mathbb{F}_2}(\iota(c_n)b_{n,en}), \\ C_d(\psi) &= \sum_n \sum_{\substack{s \geq 0 \\ 2^s d \leq en}} (\iota(c_n)b_{n,en-2^s d})^{2^{-s}}, \\ T_{e,u}(\psi) &= \sum_{d>0 \text{ odd}} C_d(\psi) \varpi^{-d}.\end{aligned}$$

All sums are finite, and $C_d = 0$ for $d > e \max \text{supp}(\psi)$.

Theorem 6.1 (Ramified restriction law). *In the local coordinates*

$$W_q(K) \cong \mathbb{F}_2 \oplus R_\pi \oplus \mathbb{F}_2,$$

scalar restriction to L is

$$(\phi_0, \psi, \phi_1) \mapsto (\bar{f}\phi_0 + \eta_{e,u}(\psi) + (1 - \bar{e})\bar{f}\phi_1, T_{e,u}(\psi), \bar{e}\bar{f}\phi_1).$$

These transformations compose in towers.

Proof. Substitution sends $c_n \pi^{-n}$ to

$$\iota(c_n)u^{-n}\varpi^{-en}.$$

Its coefficient at exponent $-2^s d$ is $\iota(c_n)b_{n,en-2^s d}$, so the preceding odd-pole normal form gives exactly C_d . Its coefficient at exponent zero is $\iota(c_n)b_{n,en}$, giving $\eta_{e,u}$.

Choose $a \in k$ with $\text{Tr}_{k/\mathbb{F}_2}(a) = 1$. The unramified generator $[1, a]$ restricts to $\bar{f}$ times the unramified generator because $\text{Tr}_{\ell/\mathbb{F}_2}(\iota(a)) = \bar{f}$. For the ramified generator, put $A = u\varpi^e$ and use

$$\langle \pi \rangle [1, a] = [A, \iota(a)/A].$$

Write $A = A_{\text{ev}} + A_{\text{odd}}$ by parity of the ϖ -exponent. The local block decomposition is

$$[A, \iota(a)/A] = [1, A_{\text{ev}}\iota(a)/A] \perp \langle \varpi \rangle [1, A_{\text{odd}}\iota(a)/A].$$

Both additive slots are integral and their constant terms are respectively $\iota(a), 0$ when e is even and $0, \iota(a)$ when e is odd. Their positive terms are Artin–Schreier trivial. Thus the ramified bit becomes unramified for even e , remains ramified for odd e , and is multiplied by $\bar{f}$; no wild term remains. This proves the formula.

For a tower $K \subset L \subset M$, both iterated transformation and direct transformation represent actual scalar restriction followed by the unique odd-pole normal form. Uniqueness makes them equal. $\square$

The trace labels admit an explicit contravariant transformation. For $\lambda \in \mathbb{F}_2$ and a finite family $z_d \in \ell$, set

$$Z_n = \text{Tr}_{\ell/k} \left(\lambda b_{n,en} + \sum_{\substack{d,s \\ 2^s d \leq en}} z_d^{2^s} b_{n,en-2^s d} \right).$$

Frobenius invariance and trace transitivity give

$$\lambda \eta_{e,u}(\psi) + \sum_d \text{Tr}_{\ell/\mathbb{F}_2}(z_d C_d(\psi)) = \sum_n \text{Tr}_{k/\mathbb{F}_2}(Z_n c_n).$$

Hence base change is a basis-free relabelling of the trace-bit games, and the adjoints compose contravariantly in towers.

There is a necessary qualification concerning the word “transfer.” The Aravire–Jacob endpoint map $W_1(K) \rightarrow W_q(k)$ kills the wild summand and reads the ramified bit. Theorem 6.1 therefore gives the usual ramification-weighted diagram

$$\sigma_L \circ \text{res}_{L/K} = \bar{e} \text{res}_{\ell/k} \circ \sigma_K.$$

An arbitrary Scharlau transfer does not descend to W_1 . Even for $L = K$, the functional $s(x) = \pi x$ sends an unramified generator to its nonzero $\langle \pi \rangle$ multiple. Thus a stronger transfer claim must specify a normalized functional; the endpoint diagram and trace-character adjunction above are the canonical statements.

7 The finite local-access obstruction

The finite compiler of Theorem 5.2 omits its zero coordinates after inspecting the class. A strict Gold-style local realization would ask for more: with the polar form fixed, one finite list of position universes, loadings, and coordinate labels must be independent of the refinement, while only move predicates may query its diagonal. That conjunction is inconsistent over $\mathbb{F}_2(t)$.

Theorem 7.1 (No fixed finite local-access family). *Let $K = \mathbb{F}_2(t)$. No fixed finite P/N family, independent of the quadratic refinement, is faithful on all nonsingular refinements of one fixed binary polar space. Consequently it cannot equal the canonical Witt-coordinate family.*

Proof. On $V = K^2$ fix the polar form $B(e, f) = 1$. For every monic irreducible $P \in \mathbb{F}_2[t]$, consider

$$q_P(x, y) = x^2 + xy + P^{-1}y^2 = [1, P^{-1}].$$

All q_P have polar form B . At the place P , with $\pi_P = P$, the local decomposition is

$$(\phi_0, \psi, \phi_1) = (0, \pi_P^{-1}, 0).$$

At every other finite place there is no negative Laurent term. The classes $[q_P]$ are therefore pairwise distinct by the global coordinate theorem. There are infinitely many monic irreducibles.

If the fixed family has N roots, its P/N vector lies in $\mathbb{F}_2^N$, a finite set. It cannot distinguish the infinite family $([q_P])_P$. The conclusion is independent of how many diagonal queries a transition makes. $\square$

The obstruction isolates the exact alternatives. A family indexed by every place and pole label can keep refinement-independent boards, but it is infinite, although each class has finite nonzero support. Selecting only that support produces a finite family, but the selection depends on the refinement and is therefore a sparse classifier or transducer. Restricting the input to a fixed finite set of places and a bounded conductor also yields a finite static family. These are coherent relaxations; none satisfies the literal conjunction ruled out by the theorem.

For one requested coordinate, an on-demand local arena is available. Choose a public symplectic frame (u_i, v_i) . The class is the orthogonal sum of $[Q(u_i), Q(v_i)]$. If $a = a_{\text{ev}} + a_{\text{odd}}$ is the Laurent-parity split, the explicit per-plane contribution at v is

$$D_v([a, b]) = (\epsilon_v(a_{\text{ev}}b), \rho_v(ab), \epsilon_v(a_{\text{odd}}b)).$$

Thus the requested coordinate is the xor of elementary constant, ramified, or trace evaluations of these D_v values. A source-only matching FIFO board can query $Q(u_i), Q(v_i)$ when the source pair opens and charge that local bit. The public matching strategy forces the total charge, and a double one-option suspension normalizes the value to exactly 0 or *. This uses two local diagonal queries per source transition, but it is an on-demand coordinate arena, not a fixed finite family covering every possible place.

8 The Gold–Arf boundary

The distinction between pointwise quadratic realization and Witt realization is structural, not merely an implementation choice.

Proposition 8.1 (Pointwise loadings do not descend). *A Witt-invariant additive family cannot literally reproduce the complete loaded-root family of a pointwise quadratic realization. It can recover only its Witt/Arf shadow.*

Proof. The hyperbolic plane

$$H(x, y) = xy$$

is zero in $W_q(\mathbb{F}_2)$. Every Witt-invariant realization must therefore map H to the zero family. But $H(1, 1) = 1$, so any exact pointwise realization has an N -position at the loading $(1, 1)$. The pointwise outcome function is not invariant under hyperbolic stabilization. $\square$

For a nonsingular finite binary form, the Arf invariant controls the sign of the zero-set bias. This class-level statistic is exactly what the constant coordinate of Theorem 4.1 retains. Thus the compatible meaning of “recover Gold–Arf” is agreement of the Witt coordinate and outcome bias, not literal equality of all loaded boards.

9 A singular additive obstruction

The nonsingular Witt group is cancellative. Arbitrary singular quadratic spaces form a different object: quasilinear directions already make their orthogonal-sum monoid noncancellative.

Theorem 9.1 (Quasilinear cancellation obstruction). *Let F be a field of characteristic two. No isometry-invariant, orthogonal-sum-additive map from all finite-dimensional quadratic spaces over F to a cancellative commutative monoid can distinguish the restrictions of all forms to their polar radicals. In particular, no such family of impartial game values is faithful on arbitrary singular spaces.*

Proof. Fix $c \neq 0$ and write $L_c = \langle c \rangle$ and $Z = \langle 0 \rangle$. The invertible change of variables $(u, v) = (x + y, y)$ gives

$$L_c \perp L_c \cong L_c \perp Z, \quad cx^2 + cy^2 = c(x + y)^2.$$

Let f be an additive isometry invariant with values in a cancellative commutative monoid. Then

$$f(L_c) + f(L_c) = f(L_c) + f(Z).$$

Cancellation gives $f(L_c) = f(Z)$. But the restrictions to the polar radicals are respectively the nonzero quasilinear form cx^2 and the zero form. Therefore f cannot retain the requested radical datum.

Finite impartial normal-play values are numbers, and disjunctive sum is their xor addition, so their value monoid is cancellative. The conclusion applies coordinatewise to any additive impartial family. $\square$

The obstruction leaves two coherent alternatives. One may retain orthogonal-sum additivity and classify only the cancellative quotient of the quasilinear monoid, or retain the full radical restriction in a separate nonadditive report. Requiring both is inconsistent.

10 Formal verification and implementation boundary

The Ogdoad Lean modules `WittRealizationExact` and `WittRamification` formalize the parts of the proof that do not require a quadratic Witt-group library [1]. Given linear maps fitting the exact-sequence hypotheses, the first proves:

- the residue image is exactly the transfer kernel;
- equal residues differ by a unique constant;
- an explicit constant detector gives the linear equivalence $W \simeq C \times \ker(\text{transfer})$;
- $z \mapsto (\lambda \mapsto \text{Tr}(\lambda z))$ is a basis-free injective finite-field presentation; and
- an absorption relation $a + a = a + z$ defeats every additive injection into a cancellative target.

The second module represents a finite principal part by pole order and proves:

- an elementary Artin–Schreier move kills one even top pole and changes only a strictly lower order;
- descending normalization terminates, removes every bounded positive even pole, preserves the constant coefficient, and is triangular in pole order;
- every normalization is connected to its input by a finite chain of exact Artin–Schreier moves; and
- no map from an infinite class family to a fixed finite P/N outcome vector is injective.

Lean does not formalize the Aravire–Jacob sequence, the local quadratic Witt groups, Lemma 3.1, the unit-series substitution in Theorem 6.1, or the infinitude of the rank-two family; those remain paper-level mathematics grounded in the cited exact sequence and finite-field theory. The Rust local decomposition computes the triple (ϕ_0, ψ, ϕ_1) with canonical uniformizers. It does not yet expose the global coordinate theorem as a public Witt-class API or implement ramified coordinate transport.

11 Conclusion

The nonsingular realization problem has a complete classification answer. The infinite unramified coordinate and all second residues give canonical coordinates for $W_q(\mathbb{F}_2(t))$. Finite trace characters present the wild odd-pole terms basis-freely, and the games 0 and $*$ realize the resulting finite-support bit vector exactly. Reciprocity is one parity equation on the ramified Arf bits. Under ramified scalar extension, the unit jet and finite inverse-Frobenius recurrence give the exact triangular coordinate law.

The stricter finite local-access request fails for a simpler reason: one fixed binary polar form supports infinitely many Witt-distinct place refinements, so no fixed finite P/N directory can be faithful. An infinite on-demand coordinate family, a refinement-dependent sparse transducer, or a bounded place-and-conductor stratum is coherent, but the original conjunction is not. Beyond the nonsingular group, the quasilinear absorption relation separately rules out the additive singular extension. Both obstructions are algebraic and precede any choice of game rules.

References

- [1] a9lim. Ogload: Clifford algebras, quadratic forms, and combinatorial games. Software and formalization, 2026.
- [2] Roberto Aravire and Bill Jacob. Quadratic forms over rational function fields in characteristic 2. *Pacific Journal of Mathematics*, 228(1):19–51, 2006.
- [3] Cahit Arf. Untersuchungen ”uber quadratische formen in k”orpern der charakteristik 2, teil i. *Journal f”ur die reine und angewandte Mathematik*, 183:148–167, 1941.
- [4] John H. Conway. *On Numbers and Games*. Academic Press, London, 1976.
- [5] Richard Elman, Nikita Karpenko, and Alexander Merkurjev. *The Algebraic and Geometric Theory of Quadratic Forms*, volume 56 of *Colloquium Publications*. American Mathematical Society, 2008.
- [6] Rudolf Lidl and Harald Niederreiter. *Finite Fields*, volume 20 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, 2nd edition, 1997.