

STT 3850 Midterm Study Guide

Andrew Thorp

October 9, 2017

Statistics

Characterizing a set of data (3 S's)

- Shape: how the data is distributed
- Low outliers make a dataset skewed to the Right
- High outliers make a dataset skewed to the Left
- Normal distributions have fairly even outliers on either side

Center: Where the data is centered around

- Normal: If the dataset has a normal distribution (shape) this can be calculated using the `mean($data)` function.
- Skewed (left or right): The mean will be misrepresent the center. Calculate a skewed center using `median($data)`.

Spread: How far the data differs from the center

- Normal: If the dataset has a normal distribution, then the standard deviation applies to both sides of the data and so it represents the spread.
- Skewed (left or right): If the dataset is skewed on either side, the deviation above and below the center will not be the same, so you must calculate it using `IQR($data)` for the interquartile range.

Hypothesis testing (5 step procedure)

- Z-Score: The number of Standard deviations an element is from the mean.
 - P-Score:
 - $\bar{X} = \text{mean}$
1. Specify the Null and Alternative hypothesis
 - Null hypothesis notated as $H_0 : M = \text{value}$ or $\bar{M}_1 - \bar{M}_2 = 0$
 - Alternative hypothesis notated as $H_A : M \neq 0$
 2. Test your statistic using the Z-test or T-test
 - `t.test(variable~categoricalVariable, data=DF)` will perform the t-test on a set of data. If the data is not tidy you might want to use `dplyr` to tidy it up first.

Example:

```
DF <- ChickWeight %>% # Imports data frame
  filter(Diet %in% c(3:4)) # Then removes all columns except for 3 and 4
                             # Weight~Diet means the weight across Diets value.
t.test(weight~Diet, data=DF) # Quickly gives us what we need to know
```

3. Determine rejection region

- Rejection region is the percent of data points on either end of a data set.
- Percentile is represented by alpha (α)

- Unless otherwise specified, assume rejection region to be 5% (0.05).

4. Statistical conclusion.

You can either

- Reject the null hypothesis H_0 if $P < \alpha$

or

- Fail to reject the null hypothesis H_0 if $P \geq \alpha$

Based on the output from the previous example pictured below, $P = 0.45$, and we assume $\alpha = 0.05$. P is far greater than α , so we **fail to reject the null hypothesis H_0** .

```
##
## Welch Two Sample t-test
##
## data: weight by Diet
## t = 0.75908, df = 226.16, p-value = 0.4486
## alternative hypothesis: true difference in means is not equal to 0
## 95 percent confidence interval:
## -12.26840 27.64298
## sample estimates:
## mean in group 3 mean in group 4
## 142.9500 135.2627
```

5. English conclusion (or Spanish, if that's your thing)

- The statistical conclusion deals with the null hypothesis H_0 , but the English conclusion only deals with things in terms of the Alternative Hypothesis H_A (written out)

In the case of our example from above, we failed to reject the null hypothesis, so we **failed to find evidence supporting a weight difference in chicks between Diets 3 and 4**.

- The English hypothesis must ALWAYS be written this way. You either find evidence to support H_A or you fail to find evidence to support H_A .

Cohens'd (effect size)

- The difference between a control group's mean, and an experiment group's mean, measured in standard deviations.
- Measured using the following formula: $d = \frac{|M_{control} - M_{experimental}|}{SD}$
- How large is the effect size?
 - small if $|d| < 0.20$
 - medium if $|d| < 0.5$
 - large if $|d| > 0.5$
- Compute using R with `lsr.cohensD(post_data, pre_data, method = "paired")`

```
Diet_3 <- filter(ChickWeight, Diet == 3)
Diet_4 <- filter(ChickWeight, Diet == 4)
#method should normally be "paired", but the sample size was different
cohensD(Diet_4$weight, Diet_3$weight, method = "unequal")
```

Markdown Dplyr Ggplot2