@apaji_x ADHD claims: evidence check (second run)

Date checked: 11 Oct 2026, 8:06 AM IST; re-evaluated 11 Oct 2026

Scope: posts from 11 Oct 2026, 2:45 AM–3:59 AM IST (11 original posts in 2 self-reply threads; 32 reposts not checked)

5 Supported	4 Partially supported	1 Insufficient evidence	0 Unsupported

Section 1 · Claims

Supported 5

Claim #A1 · Supported · High confidence

Research Quality: Claim: iron is a cofactor for tyrosine hydroxylase, the enzyme that makes dopamine. This is textbook biochemistry (high certainty for the mechanism only) [1]. The post doesn't claim everyday iron levels change ADHD symptoms, so no red flags apply.

X Post Link · Citations for this claim

RESEARCHERS ARE INVESTIGATING THE IRON + ADHD CONNECTION. 1/6

1/6 – RESEARCHERS ARE INVESTIGATING THE IRON + ADHD CONNECTION. 1/6

2/6 – 1. IRON HELPS YOUR BRAIN MAKE DOPAMINE.

Dopamine plays a role in motivation, attention, and reward processing. Your body uses an iron-dependent enzyme called tyrosine hydroxylase to produce it. That's one reason researchers are interested in how iron status might relate to brain function. 2/6

3/6 – 2. FERRITIN MEASURES YOUR STORED IRON.

You can have depleted iron stores before developing obvious iron-deficiency anemia. Ferritin is one of the main blood tests clinicians use to assess iron stores, although inflammation can sometimes make ferritin appear higher than expected. 3/6

4/6 – 3. SOME STUDIES HAVE FOUND LOWER FERRITIN LEVELS IN CHILDREN WITH ADHD.

Researchers have reported lower average ferritin levels in some groups of children with ADHD compared with children without ADHD. However, not every study finds the same result, and these findings don't establish that iron deficiency causes ADHD. 4/6

5/6 – 4. LOW IRON CAN CAUSE SYMPTOMS THAT OVERLAP WITH ADHD.

Iron deficiency can contribute to fatigue, weakness, and difficulty concentrating. These symptoms may resemble or worsen some attention-related difficulties, but they can also have many other explanations. 5/6

6/6 – 5. THIS DOESN'T MEAN EVERYONE WITH ADHD NEEDS IRON SUPPLEMENTS.

Iron is essential, but taking extra iron when you aren't deficient won't necessarily improve attention or dopamine function. Too much iron can be harmful, so testing and treatment should be guided by a healthcare professional. 6/6

Claim #A2 · **Supported** · High confidence

Research Quality: Claim: ferritin reflects iron stores, which can run low before anaemia, and inflammation raises ferritin. All three points match the WHO 2020 guideline [1] and a Lancet seminar [2] (GRADE moderate, strong consensus). The only caveat is that cut-offs vary by guideline and age.

X Post Link · Citations for this claim

RESEARCHERS ARE INVESTIGATING THE IRON + ADHD CONNECTION. 1/6

1/6 – RESEARCHERS ARE INVESTIGATING THE IRON + ADHD CONNECTION. 1/6

2/6 – 1. IRON HELPS YOUR BRAIN MAKE DOPAMINE.

Dopamine plays a role in motivation, attention, and reward processing. Your body uses an iron-dependent enzyme called tyrosine hydroxylase to produce it. That's one reason researchers are interested in how iron status might relate to brain function. 2/6

3/6 – 2. FERRITIN MEASURES YOUR STORED IRON.

You can have depleted iron stores before developing obvious iron-deficiency anemia. Ferritin is one of the main blood tests clinicians use to assess iron stores, although inflammation can sometimes make ferritin appear higher than expected. 3/6

4/6 – 3. SOME STUDIES HAVE FOUND LOWER FERRITIN LEVELS IN CHILDREN WITH ADHD.

Researchers have reported lower average ferritin levels in some groups of children with ADHD compared with children without ADHD. However, not every study finds the same result, and these findings don't establish that iron deficiency causes ADHD. 4/6

5/6 – 4. LOW IRON CAN CAUSE SYMPTOMS THAT OVERLAP WITH ADHD.

Iron deficiency can contribute to fatigue, weakness, and difficulty concentrating. These symptoms may resemble or worsen some attention-related difficulties, but they can also have many other explanations. 5/6

6/6 – 5. THIS DOESN'T MEAN EVERYONE WITH ADHD NEEDS IRON SUPPLEMENTS.

Iron is essential, but taking extra iron when you aren't deficient won't necessarily improve attention or dopamine function. Too much iron can be harmful, so testing and treatment should be guided by a healthcare professional. 6/6

Claim #A3 · Supported · High confidence

Research Quality: Claim: some studies find lower ferritin in ADHD, not all do, and this doesn't prove causation. Three Level 1 observational meta-analyses agree on a small association [1] [2] [3] (GRADE low), and the post's hedging matches that. Red flags: high heterogeneity, and in Tseng the iron-deficiency odds ratio lost significance after correcting for publication bias. Prospective cohort or Mendelian-randomisation studies would strengthen it.

X Post Link · Citations for this claim

RESEARCHERS ARE INVESTIGATING THE IRON + ADHD CONNECTION. 1/6

1/6 – RESEARCHERS ARE INVESTIGATING THE IRON + ADHD CONNECTION. 1/6

2/6 – 1. IRON HELPS YOUR BRAIN MAKE DOPAMINE.

Dopamine plays a role in motivation, attention, and reward processing. Your body uses an iron-dependent enzyme called tyrosine hydroxylase to produce it. That's one reason researchers are interested in how iron status might relate to brain function. 2/6

3/6 – 2. FERRITIN MEASURES YOUR STORED IRON.

You can have depleted iron stores before developing obvious iron-deficiency anemia. Ferritin is one of the main blood tests clinicians use to assess iron stores, although inflammation can sometimes make ferritin appear higher than expected. 3/6

4/6 – 3. SOME STUDIES HAVE FOUND LOWER FERRITIN LEVELS IN CHILDREN WITH ADHD.

Researchers have reported lower average ferritin levels in some groups of children with ADHD compared with children without ADHD. However, not every study finds the same result, and these findings don't establish that iron deficiency causes ADHD. 4/6

5/6 – 4. LOW IRON CAN CAUSE SYMPTOMS THAT OVERLAP WITH ADHD.

Iron deficiency can contribute to fatigue, weakness, and difficulty concentrating. These symptoms may resemble or worsen some attention-related difficulties, but they can also have many other explanations. 5/6

6/6 – 5. THIS DOESN'T MEAN EVERYONE WITH ADHD NEEDS IRON SUPPLEMENTS.

Iron is essential, but taking extra iron when you aren't deficient won't necessarily improve attention or dopamine function. Too much iron can be harmful, so testing and treatment should be guided by a healthcare professional. 6/6

Claim #A5 · Supported · High confidence

Research Quality: Claim: not everyone with ADHD needs iron, extra iron without deficiency won't help, and testing should come first. A Level 1 review of 3 small RCTs found no significant benefit and flagged fatal overdose risk [1], and benefits disappear without deficiency [2] (GRADE low, strong consensus on "test, then

treat"). The cautious claim matches this. The often-cited pilot (n=23, 5 on placebo) is at high risk of bias [3]; larger RCTs in iron-deficient children with ADHD would help.

X Post Link · Citations for this claim

RESEARCHERS ARE INVESTIGATING THE IRON + ADHD CONNECTION. 1/6

1/6 – RESEARCHERS ARE INVESTIGATING THE IRON + ADHD CONNECTION. 1/6

2/6 – 1. IRON HELPS YOUR BRAIN MAKE DOPAMINE.

Dopamine plays a role in motivation, attention, and reward processing. Your body uses an iron-dependent enzyme called tyrosine hydroxylase to produce it. That's one reason researchers are interested in how iron status might relate to brain function. 2/6

3/6 – 2. FERRITIN MEASURES YOUR STORED IRON.

You can have depleted iron stores before developing obvious iron-deficiency anemia. Ferritin is one of the main blood tests clinicians use to assess iron stores, although inflammation can sometimes make ferritin appear higher than expected. 3/6

4/6 – 3. SOME STUDIES HAVE FOUND LOWER FERRITIN LEVELS IN CHILDREN WITH ADHD.

Researchers have reported lower average ferritin levels in some groups of children with ADHD compared with children without ADHD. However, not every study finds the same result, and these findings don't establish that iron deficiency causes ADHD. 4/6

5/6 – 4. LOW IRON CAN CAUSE SYMPTOMS THAT OVERLAP WITH ADHD.

Iron deficiency can contribute to fatigue, weakness, and difficulty concentrating. These symptoms may resemble or worsen some attention-related difficulties, but they can also have many other explanations. 5/6

6/6 – 5. THIS DOESN'T MEAN EVERYONE WITH ADHD NEEDS IRON SUPPLEMENTS.

Iron is essential, but taking extra iron when you aren't deficient won't necessarily improve attention or dopamine function. Too much iron can be harmful, so testing and treatment should be guided by a healthcare professional. 6/6

Claim #B5 · Supported · Moderate confidence

Research Quality: Claim: simple tracking, such as a check mark, helps maintain routines. A Level 1 meta-analysis of 138 randomised studies (N=19,951) found progress monitoring improves goal attainment (d=0.40), more so when recorded physically [1] (GRADE moderate, strong consensus). Red flags: general population, not ADHD, and bias tests unverified. Self-monitoring trials in adults with ADHD would strengthen it.

X Post Link · Citations for this claim

Consistency is the hardest part of ADHD, and it is not a character flaw. Here is how to keep showing up for your food and routines without burning out or quitting. 1/5

1/5 – Consistency is the hardest part of ADHD, and it is not a character flaw. Here is how to keep showing up for your food and routines without burning out or quitting. 1/5

2/5 – Fix 1: Celebrate small wins. Ate breakfast, drank water or packed a snack? That counts. Noticing wins gives your brain the reward it craves and keeps the habit alive. 2/5

3/5 – Fix 2: Rotate things to keep it fresh. Swap a snack, try a new topping or change your breakfast every couple of weeks. A little novelty keeps the routine interesting. 3/5

4/5 – Fix 3: Use a never miss twice rule. Skipping one day is fine. Just restart at the next meal instead of waiting for a perfect Monday. 4/5

5/5 – Fix 4: Track simply. A check mark on a calendar or a quick phone note is enough. Make tracking so easy that it takes five seconds or less. 5/5

Partially supported 4

Claim #A4 · Partially supported · Moderate confidence

Research Quality: Claim: iron deficiency can cause fatigue, weakness and poor concentration resembling ADHD. A Level 1 meta-analysis of RCTs found iron improved fatigue (d=0.34) and memory in non-anaemic people, but not attention [1] (GRADE low), so the concentration part is only partly supported; fatigue is well recognised [2]. Red flag: bias tests couldn't be verified (no free full text). RCTs in iron-deficient adults with ADHD measuring attention would strengthen it.

X Post Link · Citations for this claim

RESEARCHERS ARE INVESTIGATING THE IRON + ADHD CONNECTION. 1/6

1/6 – RESEARCHERS ARE INVESTIGATING THE IRON + ADHD CONNECTION. 1/6

2/6 – 1. IRON HELPS YOUR BRAIN MAKE DOPAMINE.

Dopamine plays a role in motivation, attention, and reward processing. Your body uses an iron-dependent enzyme called tyrosine hydroxylase to produce it. That's one reason researchers are interested in how iron status might relate to brain function. 2/6

3/6 – 2. FERRITIN MEASURES YOUR STORED IRON.

You can have depleted iron stores before developing obvious iron-deficiency anemia. Ferritin is one of the main blood tests clinicians use to assess iron stores, although inflammation can sometimes make ferritin appear higher than expected. 3/6

4/6 – 3. SOME STUDIES HAVE FOUND LOWER FERRITIN LEVELS IN CHILDREN WITH ADHD.

Researchers have reported lower average ferritin levels in some groups of children with ADHD compared

with children without ADHD. However, not every study finds the same result, and these findings don't establish that iron deficiency causes ADHD. 4/6

5/6 – 4. LOW IRON CAN CAUSE SYMPTOMS THAT OVERLAP WITH ADHD.

Iron deficiency can contribute to fatigue, weakness, and difficulty concentrating. These symptoms may resemble or worsen some attention-related difficulties, but they can also have many other explanations. 5/6

6/6 – 5. THIS DOESN'T MEAN EVERYONE WITH ADHD NEEDS IRON SUPPLEMENTS.

Iron is essential, but taking extra iron when you aren't deficient won't necessarily improve attention or dopamine function. Too much iron can be harmful, so testing and treatment should be guided by a healthcare professional. 6/6

Claim #B1 · **Partially supported** · Moderate confidence

Research Quality: Claim: inconsistency is the hardest part of ADHD and not a character flaw. A Level 1 meta-analysis of 319 studies, corrected for publication bias, shows greater performance variability in ADHD (GRADE moderate) [1]. "The hardest part" is an unsupported ranking, and adults with ADHD didn't differ from clinical controls. Daily-life studies linking variability to routine adherence would strengthen it.

X Post Link · Citations for this claim

Consistency is the hardest part of ADHD, and it is not a character flaw. Here is how to keep showing up for your food and routines without burning out or quitting. 1/5

1/5 – Consistency is the hardest part of ADHD, and it is not a character flaw. Here is how to keep showing up for your food and routines without burning out or quitting. 1/5

2/5 – Fix 1: Celebrate small wins. Ate breakfast, drank water or packed a snack? That counts. Noticing wins gives your brain the reward it craves and keeps the habit alive. 2/5

3/5 – Fix 2: Rotate things to keep it fresh. Swap a snack, try a new topping or change your breakfast every couple of weeks. A little novelty keeps the routine interesting. 3/5

4/5 – Fix 3: Use a never miss twice rule. Skipping one day is fine. Just restart at the next meal instead of waiting for a perfect Monday. 4/5

5/5 – Fix 4: Track simply. A check mark on a calendar or a quick phone note is enough. Make tracking so easy that it takes five seconds or less. 5/5

Claim #B2 · **Partially supported** · Moderate confidence

Research Quality: Claim: celebrating small wins supplies the reward the ADHD brain needs. Level 1 meta-analyses show steeper delay discounting (d=0.43, no publication bias) [1] and weaker reward anticipation [2] in ADHD, which supports the mechanism (GRADE low). Self-celebration itself is untested (indirectness). RCTs of immediate self-reward in ADHD habit programmes would strengthen it.

X Post Link · Citations for this claim

Consistency is the hardest part of ADHD, and it is not a character flaw. Here is how to keep showing up for your food and routines without burning out or quitting. 1/5

1/5 – Consistency is the hardest part of ADHD, and it is not a character flaw. Here is how to keep showing up for your food and routines without burning out or quitting. 1/5

2/5 – Fix 1: Celebrate small wins. Ate breakfast, drank water or packed a snack? That counts. Noticing wins gives your brain the reward it craves and keeps the habit alive. 2/5

3/5 – Fix 2: Rotate things to keep it fresh. Swap a snack, try a new topping or change your breakfast every couple of weeks. A little novelty keeps the routine interesting. 3/5

4/5 – Fix 3: Use a never miss twice rule. Skipping one day is fine. Just restart at the next meal instead of waiting for a perfect Monday. 4/5

5/5 – Fix 4: Track simply. A check mark on a calendar or a quick phone note is enough. Make tracking so easy that it takes five seconds or less. 5/5

Claim #B4 · Partially supported · Moderate confidence

Research Quality: Claim: missing one day doesn't break a habit ("never miss twice"). A single prospective study (n=82 analysed) found one missed opportunity didn't materially affect habit formation [1] (Level 3, GRADE low). Red flags: unreplicated and no ADHD participants. Replication in ADHD would strengthen it.

X Post Link · Citations for this claim

Consistency is the hardest part of ADHD, and it is not a character flaw. Here is how to keep showing up for your food and routines without burning out or quitting. 1/5

1/5 – Consistency is the hardest part of ADHD, and it is not a character flaw. Here is how to keep showing up for your food and routines without burning out or quitting. 1/5

2/5 – Fix 1: Celebrate small wins. Ate breakfast, drank water or packed a snack? That counts. Noticing wins gives your brain the reward it craves and keeps the habit alive. 2/5

3/5 – Fix 2: Rotate things to keep it fresh. Swap a snack, try a new topping or change your breakfast every couple of weeks. A little novelty keeps the routine interesting. 3/5

4/5 – Fix 3: Use a never miss twice rule. Skipping one day is fine. Just restart at the next meal instead of waiting for a perfect Monday. 4/5

5/5 – Fix 4: Track simply. A check mark on a calendar or a quick phone note is enough. Make tracking so easy that it takes five seconds or less. 5/5

Insufficient evidence 1

Claim #B3 · Insufficient evidence · Low confidence

Research Quality: Claim: rotating foods every couple of weeks adds novelty that keeps routines going. Nothing tests this; habit research (narrative review and one cohort) suggests repeating the same behaviour with the same cue builds habits [1] [2] (GRADE very low). It is an untested idea, probably harmless if the time and cue stay fixed. Trials comparing variety and fixed routines in ADHD would be needed.

X Post Link · Citations for this claim

Consistency is the hardest part of ADHD, and it is not a character flaw. Here is how to keep showing up for your food and routines without burning out or quitting. 1/5

1/5 – Consistency is the hardest part of ADHD, and it is not a character flaw. Here is how to keep showing up for your food and routines without burning out or quitting. 1/5

2/5 – Fix 1: Celebrate small wins. Ate breakfast, drank water or packed a snack? That counts. Noticing wins gives your brain the reward it craves and keeps the habit alive. 2/5

3/5 – Fix 2: Rotate things to keep it fresh. Swap a snack, try a new topping or change your breakfast every couple of weeks. A little novelty keeps the routine interesting. 3/5

4/5 – Fix 3: Use a never miss twice rule. Skipping one day is fine. Just restart at the next meal instead of waiting for a perfect Monday. 4/5

5/5 – Fix 4: Track simply. A check mark on a calendar or a quick phone note is enough. Make tracking so easy that it takes five seconds or less. 5/5

Unsupported 0

None.

Section 2 · Citations

One group per claim, in the same order as Section 1. Numbers match the [n] markers in each claim's Research Quality note. A source cited by several claims is repeated in each group.

Claim #A1 – RESEARCHERS ARE INVESTIGATING THE IRON + ADHD CONNECTION. 1/6

↑ Back to claim

1. Daubner SC, Le T, Wang S (2011). Tyrosine hydroxylase and regulation of dopamine synthesis. *Archives of Biochemistry and Biophysics*. doi.org/10.1016/j.abb.2010.12.017
   Biochemistry review n/a · Textbook-level enzymology; says nothing about clinical iron status.

Claim #A2 – RESEARCHERS ARE INVESTIGATING THE IRON + ADHD CONNECTION. 1/6

↑ Back to claim

1. World Health Organization (2020). WHO guideline on use of ferritin concentrations to assess iron status in individuals and populations. *WHO, Geneva*. www.who.int/publications/i/item/9789240000124
   International guideline n/a · Formal WHO guideline process; not indexed in Crossref (checked on WHO site).
2. Pasricha SR, Tye-Din J, Muckenthaler MU, et al. (2021). Iron deficiency. *The Lancet*. doi.org/10.1016/S0140-6736(20)32594-0
   Clinical seminar (review) n/a · Authoritative narrative review.

Claim #A3 – RESEARCHERS ARE INVESTIGATING THE IRON + ADHD CONNECTION. 1/6

↑ Back to claim

1. Wang Y, Huang L, Zhang L, et al. (2017). Iron status in attention-deficit/hyperactivity disorder: a systematic review and meta-analysis. *PLOS ONE*. doi.org/10.1371/journal.pone.0169145
   Meta-analysis (observational) 11 studies; ferritin: 10 studies, 2,191 children · Newcastle-Ottawa mostly high quality; Egger non-significant (ferritin p=0.42) but ≤10 studies limits power; no COI.
2. Tseng PT, Cheng YS, Yen CF, et al. (2018). Peripheral iron levels in children with attention-deficit hyperactivity disorder: a systematic review and meta-analysis. *Scientific Reports*. doi.org/10.1038/s41598-017-19096-x
   Meta-analysis (observational) 17 studies; ferritin 1,560 vs 4,691 · Ferritin g=−0.25, I^2=83%, Egger p=0.43; the ADHD–iron-deficiency OR (1.64) lost significance after trim-and-fill (OR 1.22, CI 0.81–1.82), a publication-bias signal.
3. Gedfie S, Gashaw M, Adisu A, et al. (2026). Serum ferritin in children with attention deficit hyperactivity disorder (ADHD): a systematic review and meta-analysis. *PLOS ONE*. doi.org/10.1371/journal.pone.0359790
   Meta-analysis (observational) 21 studies, 4,058 ADHD cases · PROSPERO-registered; high heterogeneity; some funnel asymmetry but Egger p=0.30; no COI.

Claim #A5 – RESEARCHERS ARE INVESTIGATING THE IRON + ADHD CONNECTION. 1/6

↑ Back to claim

1. Elliott SD, Vickers ML, McKeon G, et al. (2024). Iron supplementation in management of neurodevelopmental disorders: systematic review, meta-analysis, and qualitative synthesis. *Journal of Neuropsychiatry and Clinical Neurosciences*. doi.org/10.1176/appi.neuropsych.20230081
   Systematic review & meta-analysis 3 RCTs, 124 participants · GRADE very low; differences not significant; flags fatal overdose risk; no COI.
2. Fiani D, Chahine S, Zaboube M, et al. (2025). Psychiatric and cognitive outcomes of iron supplementation in non-anemic children, adolescents, and menstruating adults: a meta-analysis and systematic review. *Neuroscience & Biobehavioral Reviews*. doi.org/10.1016/j.neubiorev.2025.106372
   Meta-analysis (12 RCTs + 6 pre-post) 18 studies, 1,408 participants · RCT effects small to moderate (fatigue d=0.34) and no effect on attention; benefits absent without iron deficiency; bias tests not verifiable (no free full text).
3. Konofal E, Lecendreux M, Deron J, et al. (2008). Effects of iron supplementation on attention deficit hyperactivity disorder in children. *Pediatric Neurology*. doi.org/10.1016/j.pediatrneurol.2007.08.014
   Pilot RCT n=23 (18 iron vs 5 placebo) · Tiny 3:1 placebo arm, within-group comparisons only, parent/teacher scales non-significant; 'comparable to stimulants' is an overreach (RoB 2: high).

Claim #B5 – Consistency is the hardest part of ADHD, and it is not a character flaw. Here is how to keep showing up for your food and routines without burning out or quitting. 1/5

↑ Back to claim

1. Harkin B, Webb TL, Chang BPI, et al. (2016). Does monitoring goal progress promote goal attainment? A meta-analysis of the experimental evidence. *Psychological Bulletin*. doi.org/10.1037/bul0000025
   Meta-analysis of RCTs 138 studies, N=19,951 · Large, randomised; general population (not ADHD); bias tests not verifiable (no free full text).

Claim #A4 – RESEARCHERS ARE INVESTIGATING THE IRON + ADHD CONNECTION. 1/6

↑ Back to claim

1. Fiani D, Chahine S, Zaboube M, et al. (2025). Psychiatric and cognitive outcomes of iron supplementation in non-anemic children, adolescents, and menstruating adults: a meta-analysis and systematic review. *Neuroscience & Biobehavioral Reviews*. doi.org/10.1016/j.neubiorev.2025.106372
   Meta-analysis (12 RCTs + 6 pre-post) 18 studies, 1,408 participants · RCT effects small to moderate (fatigue d=0.34) and no effect on attention; benefits absent without iron deficiency; bias tests not verifiable (no free full text).
2. Pasricha SR, Tye-Din J, Muckenthaler MU, et al. (2021). Iron deficiency. *The Lancet*. doi.org/10.1016/S0140-6736(20)32594-0
   Clinical seminar (review) n/a · Authoritative narrative review.

Claim #B1 – Consistency is the hardest part of ADHD, and it is not a character flaw. Here is how to keep showing up for your food and routines without burning out or quitting. 1/5

↑ Back to claim

1. Kofler MJ, Rapport MD, Sarver DE, et al. (2013). Reaction time variability in ADHD: a meta-analytic review of 319 studies. *Clinical Psychology Review*. doi.org/10.1016/j.cpr.2013.06.001
   Meta-analysis 319 studies · Effects corrected for measurement unreliability and publication bias; ADHD adults not distinguishable from clinical controls.

Claim #B2 – Consistency is the hardest part of ADHD, and it is not a character flaw. Here is how to keep showing up for your food and routines without burning out or quitting. 1/5

↑ Back to claim

1. Jackson JNS, MacKillop J (2016). Attention-deficit/hyperactivity disorder and monetary delay discounting: a meta-analysis of case-control studies. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*. doi.org/10.1016/j.bpsc.2016.01.007
   Meta-analysis (case-control) 25 comparisons, N=3,913 · No publication bias (Egger p=0.53; trim-and-fill d 0.43→0.40); minimal heterogeneity.
2. Plichta MM, Scheres A (2014). Ventral-striatal responsiveness during reward anticipation in ADHD and its relation to trait impulsivity in the healthy population: a meta-analytic review of the fMRI literature. *Neuroscience & Biobehavioral Reviews*. doi.org/10.1016/j.neubiorev.2013.07.012
   Meta-analysis (fMRI) 8 studies, N=340 · Medium effect (d=0.48); few, small studies (authors note limited power); no COI.

Claim #B4 – Consistency is the hardest part of ADHD, and it is not a character flaw. Here is how to keep showing up for your food and routines without burning out or quitting. 1/5

↑ Back to claim

1. Lally P, van Jaarsveld CHM, Potts HWW, et al. (2010). How are habits formed: modelling habit formation in the real world. *European Journal of Social Psychology*. doi.org/10.1002/ejsp.674
   Prospective observational 96 volunteers, 82 analysed · Single, unreplicated study of self-chosen habits; not ADHD; widely cited.

Claim #B3 – Consistency is the hardest part of ADHD, and it is not a character flaw. Here is how to keep showing up for your food and routines without burning out or quitting. 1/5

↑ Back to claim

1. Wood W, Rünger D (2016). Psychology of habit. *Annual Review of Psychology*. doi.org/10.1146/annurev-psych-122414-033417
   Narrative review n/a · Authoritative review of habit research; not ADHD-specific.
2. Lally P, van Jaarsveld CHM, Potts HWW, et al. (2010). How are habits formed: modelling habit formation in the real world. *European Journal of Social Psychology*. doi.org/10.1002/ejsp.674
   Prospective observational 96 volunteers, 82 analysed · Single, unreplicated study of self-chosen habits; not ADHD; widely cited.

About this report

10 claims re-evaluated with the Scientific Evidence Evaluator framework. The iron thread is careful and well hedged. The consistency tips are sensible but rest on general habit research, not ADHD trials. One verdict changed: A4 (iron and concentration) moved to Partially supported because RCTs show no effect of iron on attention.

How claims were evaluated. Evidence hierarchy: Systematic reviews and meta-analyses (Level 1), RCTs (2), cohort (3), case-control (4), cross-sectional (5), case reports and expert opinion (6). Guidelines and clinical reviews are named as such. **GRADE certainty:** Our rating of the body of evidence for this specific claim: randomised evidence starts High and observational starts Low. Ratings go down for risk of bias, inconsistency, indirectness (e.g. non-ADHD or lab-only samples), imprecision and publication bias, and up for large effects. **Verdicts:** Supported: strong, consistent, high-quality evidence. Partially supported: some solid evidence, but the claim overgeneralises or has important caveats. Insufficient evidence: too little good-quality research to judge. Unsupported: evidence is absent for the claim as stated, or contradicts it (flagged as contradicted). Confidence is our confidence in the verdict, which can be high even when GRADE is low (e.g. a well-hedged claim).

Verdict changes: 1 changed in substance; 4 relabelled onto the four-category scale with no change in substance (Not supported or Contradicted → Unsupported; Partly → Partially supported).

Claim	Previous	Now	GRADE
#A4	Supported	Partially supported	Low

Method. Posts were pulled with the X API. Self-reply parts were reconstructed via conversation_id and reply links and quoted verbatim, numbered in order. Reference metadata, corrections and retractions were checked against Crossref and PubMed, and open-access full texts were read. GRADE ratings are our own per-claim judgement unless stated.

Not medical advice. Verdicts reflect evidence as of the check date and are group-level. Talk to a qualified clinician about diagnosis or treatment.